

## Subject Index

- A1330V polymorphism, in LRP5, as a risk factor of hypercholesterolemia in Japanese males, 751-757
- AA supplementation, *see* Amino acid (AA) supplementation
- Acanthosis nigricans, AT, liver and SM S<sub>i</sub> in extremely obese subjects with, 1658-1663
- Acarbose  
effect on fibrinolytic activity with IGT, 935-939  
effect on LDL atherogenicity with IGT, 946-952  
effects on age-related glucose intolerance and pancreatic atrophy in rats, 533-540
- ACC $\beta$ , *see* Acetyl-coenzyme A carboxylase  $\beta$
- Acetaldehyde, in moderation of long-term alcohol exposure on HSP gene expression in rats, 843-851
- Acetylcholine, neurochemicals on awake state, S2:16
- Acetyl-coenzyme A carboxylase  $\beta$  (ACC $\beta$ ), phosphorylation, and  $\alpha$ 2 AMPK activity and phosphorylation, effects of AICAR in skeletal muscle, 304-305
- Acolbifene, hypocholesterolemic action, in intact and ovariectomized rats with diet-induced hypercholesterolemia, 605-613
- Acromegaly, active, effect on spontaneous PTH pulsatility, 736-740
- ACTH, *see* Adrenocorticotrophic hormone
- Adenosine, neurochemicals on awake state, S2:17
- Adenosine monophosphate-activated protein kinase (AMPK)  
activity and subunit expression in rat VAT and liver, long-term exercise on stimulation of, 1122-1128  
 $\alpha$ 2 isoform-specific activation by AICAR at physiological levels, effects on glucose transport and GLUT4 in mouse skeletal muscle, 300-308
- Adenosine triphosphate (ATP), role in metabolic mechanism of wakefulness and sleep, S2:24-29
- S-Adenosylhomocysteine (SAH), plasma, relation to GFR in children, 252-257
- Adipocyte precursor cells, human, effect of troglitazone on TNF- $\alpha$  and TGF- $\beta$  expression and action in primary culture of, 309-316
- Adipocytes  
3T3-L1  
differentiated, effects of insulin and leptin on fatty acid uptake in, 8-12
- Adipocytes (continued)  
gliclazide protection of, against IR-induced hydrogen peroxide, 722-730  
cultured, short- and long-term insulin-like effects of MAO and SSAO substrates, 1397-1405  
differentiation  
inhibition by TNF- $\alpha$  and TGF- $\beta$ , effect of troglitazone, 312  
and PPAR $\gamma$  mRNA expression, effect of troglitazone, 311-312  
effect of Trf and Fe on IR of glucose transport in, 1042-1045  
versus non-fat cells, IL-1Ra, cathepsin S, MIF, NGF and IL-18 release by, 1113-1121
- Adipocytokines  
circulating, effect of spironolactone, in NIDDM nephropathy patients, 1645-1652  
effects of angiotensin II infusion in rats, 483
- Adipokines  
and insulin, association with maternal body size and prediction of birth weight, 339-344  
and niacin treatment, 1283-1285  
in nondiabetic hypercholesterolemic patients, 918-922
- Adiponectin  
circulating plasma, and IR, CV disease risk, abdominal adiposity and, association with SNP276>T versus SNP45T>G adiponectin gene polymorphism in nonobese/nondiabetic Korean men, 59-66  
concentrations, beta cell-function, IR and, impact of rosiglitazone combined with glimepiride, 20-25  
effect on artery stiffness, in NIDDM treated with pioglitazone and metformin, 996-1001  
higher levels in American than Japanese men despite obesity, 1561-1563  
and its correlates of CV risk, in young adults, 1551-1557  
and leptin ratio, relationship with IR parameters, in subjects without hyperglycemia, 1248-1254  
in middle-aged postmenopausal women, in association with lifestyle habits, hormones and inflammatory markers, 1630-1636  
plasma, *see* Plasma adiponectin  
relation to HDL-C and plasma insulin concentration, in healthy children, 1171-1176

## Adiponectin (continued)

serum, *see* Serum adiponectin

and stress hormone, responses to maximal sculling after volume-extended training season in elite rowers, 13-19

in T cell-mediated hepatitis in mice, role of TNF- $\alpha$  in modulation of, 555-559

Adiponectin receptors, expression in lean rats and in obese Zucker rats, 396-401

## Adipose tissue (AT)

fenofibrate and lipid metabolism in rats, 731-735

increased glyceroneogenesis in rats adapted to high-protein, CHO-free diet, role of dietary FAs, 84-89

liver and SM S<sub>i</sub>, in extremely obese subjects with acanthosis nigricans, 1658-1663

mass, and glucose tolerance, effect of pregnancy and aging in rats, 409-414

percentage of, and plasma glucose, prediction of sialic acid concentration in NIDDM, 1165-1170

subcutaneous abdominal, gene expression of adiponectin, leptin, IL-6 and TNF- $\alpha$ , effect of aerobic training, in obese women, 1375-1381

white, mRNA profiles, of female Zucker diabetic fatty rats after topiramate treatment, 1411-1419

Adiposity, abdominal, and IR, CV disease risk and circulating plasma adiponectin, association with SNP276>T versus SNP45T>G adiponectin gene polymorphism in nonobese/nondiabetic Korean men, 59-66

ADMA, *see* Asymmetric dimethylarginine

## Adolescents

Asian Indians, simple anthropometric measures in fasting hyperinsulinemia and clustering of CV risk factors in, 1569-1573

with PCOS versus regular cycling adolescents, free testosterone, obesity and CVD risk factors in, 508-514

Turkish, MS in, 1002-1006

Adrenal glands, transient induction of variant hepatic lipase mRNA by corticotropin hormone in rats, 467-477

Adrenocorticotrophic hormone (ACTH), and equivalent dose of cortisol, effects on serum concentrations of lipids, Lp and Apo, 1083-1087

Adrenocorticotropin, effect of NEFAs, in healthy young men, 1022-1028

## Adults

with normal glucose regulation, association of MS and IR with cardiovascular risk in, 445-452

normal-weight, obese, and overweight, BSA in, 515-524

## Advanced glycation end products (AGEs)

association with RAGEs, in nondiabetic subjects, 1227-1231

association with thrombogenic markers in humans, 912-917

## Advanced glycation end products (AGEs) (continued)

in healthy women and women with PCOS, short-term effects of orlistat, 494-500

methylglyoxal-derived hydroimidazolone, increased serum levels, association with retinopathy in NIDDM, 232-236

NADPH (reduced form) activation by, linked oxidative stress to altered retinal VEGF, 1516-1523

Adverse drug effects, to NIDDM, a hurdle to overcome for achieving glycemic control, S1:7

Aerobic capacity, *see* VO<sub>2max</sub> during exercise

## Aerobic exercise

effects on CRP, body composition and maximum oxygen consumption, in adults, 1500-1507

and PSE supplementation, effects on blood lipid levels, 541-549

and S<sub>i</sub>, effects of repeated prolonged whole-body low-intensity exercise, 217-223

weight-bearing, effect on bone formation, during short-term weight loss in overweight obese men and women, 1616-1618

African Americans, *see* Race and ethnicity

Afro-Caribbeans, *see* Race and ethnicity

## Age

differential effects on the association of *KLOTHO* polymorphism with CAD, 1344-1351

glucose intolerance related to, effects of acarbose on, 533-540

*see also* Middle-aged women

AGEs, *see* Advanced glycation end products

Aging, effect on pregnancy-induced adaptations in glucose tolerance in rats, 409-414

252A/G polymorphism, and 804C/A, of LTA, relationship with TNF- $\alpha$  –308G/A polymorphism, in Japanese subjects with NIDDM retinopathy, 1406-1410

AGT, *see* Angiotensinogen

AICAR, *see* 5-Aminoimidazole-4-carboxamide-1- $\beta$ -D-ribose

Ala54Thr variant, of FABP2 gene, association with MS and hypertriglyceridemia, in urban South Indians, 1222-1226

Alanine aminotransferase (ALT), increased, association with NAFLD metabolic factors, 1604-1609

Albuminuria, stages of, association with Mn-SOD gene polymorphism (V16A) in Korean NIDDM patients, 1-7

## Alcohol

chronic consumption, effect on CS, PMS and LMS, in rat liver, 803-810

effect on SM HSP gene expression in rats, 843-851 exposure, and fasting, skeletal muscle and liver oxysterols during, 119-127

ingestion, effect on peptide YY serum levels and on total and octanoylated ghrelin levels in healthy subjects, 1625-1629

- Aldosterone  
and free cortisol, urinary excretion of, in response to LoBAG diet in men with NIDDM, 248  
urinary, and free cortisol, and plasma renin, in response to LoBAG diet in men with NIDDM, 248
- Alendronate, combined with HRT, effects on osteoporotic postmenopausal Chinese women, 741-747
- Allostatic load, on sleep deprivation, S2:20-23
- Alpha ( $\alpha$ )-amino nitrogen, plasma, in response to LoBAG diet in men with NIDDM, 247
- Alpha ( $\alpha$ )-cyclodextrin, effects on weight gain and selected blood parameters in rats, 195-202
- Alpha ( $\alpha$ )-glucosidase inhibitor  
ascarbose, effects on age-related glucose intolerance and pancreatic atrophy in rats, 533-540  
prevention of CHO-induced upregulation in intestinal sugar transport, in diabetic mice, 832-841  
voglibose, reduction of oxidative stress markers and soluble ICAM-1, in obese NIDDM patients, 786-793
- Alpha ( $\alpha$ )-tocopherol, and ascorbic acid, effect on down-regulation of Apo A-I gene expression in HepG2 and Caco-2 cell lines, 159-167
- ALT, *see* Alanine aminotransferase
- American Diabetes Association, newly recommended criteria for fasting hyperglycemia, relation of IR and serum high-sensitivity CRP to FBG subgroups categorized by, in 10 059 healthy Koreans, 183-187
- American subjects, men with obesity, levels of adiponectin in, 1561-1563
- Amino acid (AA) supplementation, effects on hormonal responses to resistance training overreaching, 282-291
- 5-Aminoimidazole-4-carboxamide-1- $\beta$ -D-ribonucleoside (AICAR)  
activation of  $\alpha$ 2 AMPK at physiological levels, effects on glucose transport and GLUT4 in mouse skeletal muscle, 300-308  
and moderate-intensity exercise, extent of  $\alpha$ 2 AMPK activation by, 303-304
- Ammonemia, associated with pregnancies and deliveries in LPI, 224-231
- Amniotic fluid, mid-trimester, Met concentrations in, as predictor of body weight and length, 1186-1191
- AMPK, *see* 5' Adenosine monophosphate-activated protein kinase
- Androgen, deficiency, prevalence and risk factors, in middle-aged men in Hong Kong, 1488-1494
- Anemia, associated with pregnancies and deliveries in LPI, 224-231
- Angiotensin II, infusion of, decreases plasma adiponectin levels via its type 1 receptor in rats, 478-488
- Angiotensinogen (AGT), gene polymorphism (Met235Thr), influence on visceral obesity and IR, in obese Japanese women, 819-824
- Anoxia, nocturnal, pathophysiology of, S2:47
- Anthropometric indices  
in fasting hyperinsulinemia and clustering of CV risk factors in Asian Indian adolescents, 1569-1573  
intra-abdominal fat and 3 nonobese metabolic risk factors, CT scans of, 1337-1343
- Antiepileptic drug, N-glucuronidation of, results from human UGTs, liver and kidney of Crigler-Najjar type II, 711-721
- Antioxidant response element (ARE), of Apo A-I, effects of antioxidants on, 163-164
- Antioxidants, effects on Apo A-I mRNA, protein levels, promoter activity and ARE, 162-164
- Antiretroviral therapy (ART), lipodystrophy and MS in HIV-infected patients treated with, 940-945
- Apheresis, regular LDL, and statins, effect of ezetimibe on LDL subtype distribution in patients treated by, 599-604
- Apo, *see* Apolipoprotein(s)
- APOA5, haplotype and polymorphism, longitudinal analysis, associated with variation in serum TGs, 1574-1581
- Apo A-I, *see* Apolipoprotein A-I
- APOC3 haplotype and polymorphism, longitudinal analysis, associated with variation in serum TGs, 1574-1581
- Apo C-I, *see* Apolipoprotein C-I
- Apo E, *see* Apolipoprotein E
- Apolipoprotein(s) (Apo)  
altered in IBD, as a consequence on cholesterol efflux, 980-988  
serum concentrations, effects of ACTH and equivalent dose of cortisol, 1083-1087  
*see also specific apolipoproteins*
- Apolipoprotein A-I (Apo A-I), gene expression in HepG2 and Caco-2 cell lines, effect of ascorbic acid and  $\alpha$ -tocopherol on down-regulation of, 159-167
- Apolipoprotein C-I (Apo C-I), effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26-34
- Apolipoprotein E (Apo E)  
enriched with TG-rich lipoproteins, accumulation in patients with CAD, 662-668  
isoforms, differential effects on lipolysis of VLDL TGs, 1129-1134
- ARE, *see* Antioxidant response element
- Arg121Trp* mutation, in *PAX4* gene, association with beta-cell dysfunction in Japanese NIDDM subjects, 213-216
- ART, *see* Antiretroviral therapy
- Arterial calcification, lower-extremity, as a predictor of coronary artery calcification, 1689-1696
- Arterial distensibility, estrogen treatment on, in postmenopausal women with NIDDM, 953-959
- Arterial stiffness  
carotid, association with plasma adiponectin levels, 587-592

- Arterial stiffness (continued)  
  effect of adiponectin, in NIDDM treated with  
    pioglitazone and metformin, 996-1001  
  effect of weight loss, in overweight adults, 907-911
- Ascorbic acid, and  $\alpha$ -tocopherol, effect on down-regulation  
  of Apo A-I gene expression in HepG2 and Caco-2  
  cell lines, 159-167
- Asian Indians, *see* Race and ethnicity
- Asymmetric dimethylarginine (ADMA)  
  in citrullinemia, elevated plasma concentrations of  
    endogenous NOS, 1599-1603  
  and hemodynamic regulation, in middle-aged men,  
    771-777
- AT, *see* Adipose tissue
- Athletes, elite rowers, adiponectin and stress hormone  
  responses to maximal sculling after  
  volume-extended training season, 13-19
- Atorvastatin, lowers plasma LDL-C and CRP in Japanese  
  NIDDM patients, 67-71
- ATP, *see* Adenosine triphosphate
- Atrophy  
  differential changes in respiratory capacity and  
    ischemia tolerance of mitochondria of heart with,  
      1097-1106  
  glucocorticoid-induced SM, Gln in prevention of,  
    associated with myostatin suppression, 1239-1247  
  pancreatic, effects of acarbose on, 533-540
- Autoimmunity, and glucose tolerance, family history of  
  diabetes,  $S_{\text{I}}$ , C-peptide and inflammation, in relation  
  to serum adiponectin in population sample of  
  64-year-old women, 188-194
- Autonomic nervous system, *see* Sympathetic nervous  
  system
- BDNF, *see* Brain-derived neurotrophic factor
- Beta<sub>3</sub> ( $\beta_3$ )-adrenergic receptor, polymorphisms, in GDM  
  and normal pregnancy, 1451-1456
- Beta ( $\beta$ )-adrenergic agonists, combined with PPAR, effects  
  on Lp composition in healthy moderately obese  
  subjects, 26-34
- Beta ( $\beta$ ) cell(s) (islets of Langerhans)  
  damage, induced by oxidative stress, protective role of  
    gliclazide on, S1:13  
  dysfunction, association with *Arg121Trp* variant in *PAX4*  
    gene in Japanese NIDDM subjects, 213-216  
  exhaustion, sulfonylureas in, S1:23-24  
  function, and IR, adiponectin concentrations, impact of  
    rosiglitazone combined with glimepiride, 20-25  
  isolated, dexamethasone suppresses PLC activation and  
    insulin secretion in, 35-42  
  loss of function, sulfonylureas for, S1:3-4  
  in mice, effect of SVS on glyburide-induced  
    desensitization, 1674-1680  
  protective effect of BDNF, in obese diabetic mice,  
    1286-1292  
  responses to hypoglycemia, in NIDDM, 593-598
- Beta ( $\beta$ ) cell(s) (islets of Langerhans) (continued)  
  skeletal muscle, liver, fat and, gene expression for energy  
    metabolism after ER in Zucker diabetic fatty rats,  
      43-52
- 11Beta ( $\beta$ )-hydroxysteroid dehydrogenase 2 (11 $\beta$ -HSD2),  
  physiologic roles in kidney, 1352-1357
- 11 $\beta$ -HSD2, *see* 11Beta ( $\beta$ )-hydroxysteroid dehydrogenase 2
- Bioaccumulation, short-term, of vanadium when ingested  
  with tea decoction in STZ-DM rats, 263-270
- Birth length, mid-trimester amniotic fluid Met  
  concentrations as predictor of, 1186-1191
- Birth weight  
  in adult twins, interrelation with regional lipid deposition,  
    561-562  
  mid-trimester amniotic fluid Met concentrations as  
    predictor of, 1186-1191  
  prediction of, and maternal body size, association with  
    insulin and adipokines, 339-344
- Bleeding complications, associated with pregnancies and  
  deliveries in LPI, 224-231
- Blood glucose  
  decreased concentration, and increased blood lactate  
    concentration, increased 2DG transport activity  
    accompanied by, effects of AICAR in skeletal  
      muscle, 305  
  and serum insulin and C-peptide, in PM women and  
    age-matched men with NIDDM, 334
- Blood lactate, increased concentration, and decreased blood  
  glucose concentration, increased 2DG transport  
  activity accompanied by, effects of AICAR in  
  skeletal muscle, 305
- Blood lipids, effects of PSE supplementation and aerobic  
  exercise training, 541-549
- Blood parameters, selected, and weight gain, effects of new  
  soluble dietary fiber in rats, 195-202
- Blood pressure (BP)  
  ambulatory, sleep-related clinical observations on,  
    S2:46  
  and body composition, in response to weight reduction  
    after diet and physical activity, 361  
  and counterregulatory hormones, in response to brief  
    hypoglycemia in PM women and age-matched men  
    with NIDDM, 334-336  
  and inflammatory marker and glucose metabolism,  
    relations to IMT in childhood obesity, 113-118
- BMI, *see* Body mass and body mass index
- Body composition  
  and BP, in response to weight reduction after diet and  
    physical activity, 361  
  effects of aerobic exercise, in adults, 1500-1507  
  effects of FBCx in high fat-fed rats, 198  
  and metabolic effects of diet and exercise, on obese  
    HIV-infected women, 1327-1336  
  offspring, and insulin axis, programming of maternal  
    protein intake on pregnant rat, 642-649  
  sex-specific, impact on cardiovascular risk factors, 563-569



- Body fat, *see* Adipose tissue
- Body mass and body mass index (BMI)  
 effects of HF feeding in Sprague-Dawley rat, 209  
 hip and waist circumference relative to, impact on  
 cardiovascular risk factors, 563-569  
 sequencing of ghrelin 5' flanking region in relation to,  
 1420-1425
- Body size, maternal, and prediction of birth weight,  
 association with and adipokines, 339-344
- Body surface area (BSA), in normal-weight, overweight  
 and obese adults, 515-524
- Body temperature, fluid, electrolyte and nitrogen homeostasis,  
 on sleep-related vascular disorders, S2:47
- Bodyweight (BW)  
 at birth, *see* Birth weight  
 change in, correlation with change in serum adiponectin  
 levels in Japanese population, 324-330  
 Cockcroft-Gault formula is biased by, in diabetic patients  
 with renal impairment, 108-112  
 effects of FCBx in high fat-fed rats, 198  
 and food intake, effects of NO-1886 (ibrolipim) in  
 ovariectomized rats, 154  
 gain, sulfonylureas in, S1:22-23  
 and liver relative weight, liver glycogen concentration and  
 plasma parameters in LETO and OLETF rats, 319  
 loss of, *see* Weight loss  
 normal, and overweight and obese, BSA in adults with,  
 515-524
- Bone metabolism, in PM Japanese women, effects of  
 exercise and isoflavones, 423-433
- BP, *see* Blood pressure
- Brain-derived neurotrophic factor (BDNF)  
 increased, associated with obesity in newly diagnosed  
 female with NIDDM, 852-857  
 protective effect on pancreatic islets, in obese diabetic  
 mice, 1286-1292
- BSA, *see* Body surface area
- BW, *see* Body weight
- C, *see* Cholesterol
- C1429T polymorphism, in GNB3, as a risk factor of  
 hypercholesterolemia in Japanese males, 751-757
- Caco-2 cell line, and HepG2 cell line, effect of ascorbic  
 acid and  $\alpha$ -tocopherol on down-regulation of Apo  
 A-I gene expression in, 159-167
- CAD, *see* Coronary artery disease
- Canola oil, high-oleic, and PS mixed with MTGs, effect on  
 plasma lipids in overweight men, 391-395
- CAP, *see* Cbl-associated protein
- 804C/A polymorphism, and 252A/G, of LTA, relationship  
 with TNF- $\alpha$  -308G/A polymorphism, in Japanese  
 subjects with NIDDM retinopathy, 1406-1410
- Carbohydrate(s) (CHO)  
 liquid, and EAA, ingestion, during short-term resistance  
 exercise, effect on myofibrillar protein degradation,  
 570-577
- Carbohydrate(s) (CHO) (continued)  
 upregulation induced by,  $\alpha$ -glucosidase inhibitors in  
 prevention of, in intestinal sugar transport in diabetic  
 mice, 832-841
- Cardiomyocytes, HL-1, effects of short- and long-term  
 leptin treatment on glucose and FA uptake and  
 metabolism, 1067-1075
- Cardiovascular (CV) disease  
 with NIDDM, correlation of PON1 and HCTL and  
 antioxidant capacity of HDL with severity of,  
 1201-1206  
 and postprandial hyperglycemia, 1264-1266  
 risk, *see* Risk, for CV disease  
 risk factors, and free testosterone and obesity in  
 adolescents with PCOS versus regular cycling  
 adolescents, 508-514
- Cardiovascular responses, to hypoglycemia and  
 epinephrine in IDDM, 377
- Carnitine, and FFAs and TGs, levels in liver and serum of  
 SSADDH knockout mice, 355-356
- Carotid intima-media thickness, and metabolic parameters,  
 effects of lifestyle modification, in NIDDM,  
 1053-1059
- CAT, *see* Catalase
- Catalase (CAT)  
 and CuZnSOD, MnSOD, GCLc, Gpx1 and CypB,  
 mRNA expressions and protein levels in limited  
 amounts of tissue from rat retina, 168-174  
 expression in rat retinal ischemia after reperfusion,  
 892-898  
 oxidative stress with diet-induced MS due to, 928-934
- Cathepsin S, adipocytes versus non-fat cells in release of,  
 1113-1121
- Caucasians, *see* Race and ethnicity
- Cbl-associated protein (CAP)/Cbl signaling cascade,  
 components of, effects of HF feeding in  
 Sprague-Dawley rat skeletal muscle, 203-212
- CD40 ligand (sCD40L), elevated, and disease severity, MS  
 independent association with, in symptomatic CAD,  
 1029-1034
- Cerebral infarction, silent  
 hyperhomocysteinemia a significant risk factor for, in  
 CRF patients undergoing HD, 656-661  
 progression in patients with type 2 diabetes mellitus,  
 relation to CRP and soluble adhesion molecules,  
 461-466
- Cerivastatin, TNF- $\alpha$ -induced production of PAI-1 in  
 regulation of, in nonmalignant hepatocyte cell line,  
 1464-1472
- CHD, *see* Coronary heart disease
- Chennai Urban Rural Epidemiology Study (CURES),  
 serum levels of IL-6, CRP, VCAM-1 and MCP-1 in  
 relation to IR and glucose tolerance, 1232-1238
- Children  
 healthy, adiponectin in relation to HDL-C and plasma  
 insulin concentration in, 1171-1176

## Children (continued)

## obese

prepubertal, correlation of hyperhomocysteinemia with IR and low-grade systemic inflammation in, 72-77  
relations of IMT to inflammatory marker, glucose metabolism and BP in, 113-118

overweight, effect of short-term diet and exercise intervention on MS in, 871-878

relationship between plasma SAH concentration and GFR in, 252-257

Turkish, MS in, 1002-1006

## Chinese subjects

with IGT, eNOS G894T polymorphism in prediction of glycemic status in a 5-year study, 1155-1158

older, association of vascular risk factors with increasing glycemia and normoglycemia in, 1035-1041

older than 30 years, with NIDDM, IDF definition of MS in, 1088-1096

osteoporotic postmenopausal women, effects of alendronate combined with HRT, 741-747

CHO, *see* Carbohydrate(s)

## Cholesterol (C)

plasma levels, and hepatic synthesis and intestinal absorption, effects of stigmasterol in rat, 292-299

total, effects of PSE supplementation and aerobic exercise training, 541-549

Cholesterol efflux, serum capacity using Fu5AH cell system, altered lipid, Apo and Lp in IBD as consequences on, 980-988

Cholesterol ester transfer protein, effects of PSE supplementation and aerobic exercise training, 541-549

Choline, and folate, different effects on plasma kinetics of methionine and homocysteine, in rats, 899-906

Chomczynski method, simultaneous extraction of RNA and protein, for detection of enzymes in limited amounts of tissue from rat retina, 168-174

Chronic renal failure (CRF), undergoing HD, hyperhomocysteinemia a significant risk factor for silent cerebral infarction with, 656-661

## Circadian rhythm

genetics of sleep-wake cycle, S2:7-8

and mood disorders, S2:10-11

Circulating plasma adiponectin, and IR, CV disease risk, abdominal adiposity and, association with SNP276>T versus SNP45T>G adiponectin gene polymorphism in nonobese/nondiabetic Korean men, 59-66

CIRS, *see* Collège International de Recherche Servier

Citrullinemia, elevated plasma concentrations of endogenous NOS ADMA in, 1599-1603

Cockcroft-Gault formula, biased by body weight in diabetic patients with renal impairment, 108-112

Colesevelam hydrochloride, with ezetimibe, Lp effects of, versus ezetimibe alone, in hypercholesterolemic subjects, 1697-1703

Collagen, production, effect of physiologic concentrations of leptin by human hepatic stellate cells on, 1317-1322

Collège International de Recherche Servier (CIRS), foreword, S2:1

Compliance, to medication, on overcoming hurdles for achieving glycemic control in NIDDM, S1:8-9

Computed tomography (CT), scans of intra-abdominal fat, anthropometric measurements and 3 nonobese metabolic risk factors, 1337-1343

Copper/zinc superoxide dismutase (CuZnSOD) and CAT, MnSOD, GCLc, Gpx1 and CypB, mRNA expressions and protein levels in limited amounts of tissue from rat retina, 168-174

expression in rat retinal ischemia after reperfusion, 892-898

oxidative stress with diet-induced MS due to, 928-934

Coronary artery calcification, lower-extremity calcification as a predictor of, 1689-1696

## Coronary artery disease (CAD)

accumulation of apo E-enriched TG-rich lipoproteins in patients with, 662-668

differential effects of age on the association of *KLOTHO* polymorphism with, 1344-1351

symptomatic, MS independent association with elevated sCD40L and disease severity in, 1029-1034

Coronary artery stenosis, multiple, association of hypoadiponectinemia in men with, 1653-1657

## Coronary heart disease (CHD)

early onset, association of hypoadiponectinemia in men with, 1653-1657

risk factors, association with increasing glycemia, in older Chinese population, 1035-1041

Corticosterone, impact of maternal folic acid

supplementation, in adult offsprings, 628-634

Corticotropin hormone, in rat adrenals, transient induction by variant hepatic lipase mRNA, 467-477

## Cortisol (hydrocortisone)

effects on glucocorticoid receptor in malnutrition and acute infection, 550-554

equivalent dose of, and ACTH, effects on serum concentrations of lipids, Lp and Apo, 1083-1087

in men, effect of NEFAs, 1022-1028

response to maximal sculling after volume-extended training season in elite rowers, 13-19

## Counterregulatory hormones

and BP, in response to brief hypoglycemia in PM women and age-matched men with NIDDM, 334-336

responses, effects of glimepiride and glyburide, 81

## C-peptide

and blood glucose and serum insulin, in PM women and age-matched men with NIDDM, 334

and glucose tolerance, family history of diabetes, autoimmunity, S<sub>i</sub> and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188-194

secretion rate, and minimal model analysis, in determining clinical phenotype of Japanese NIDDM subjects with *Arg121Trp* mutation, 215-216

- C-reactive protein (CRP)  
 association with metabolic risk factors among young and middle-aged Koreans, 415-421  
 in biracial younger adults, distribution of MS variables in correlation with, 699-705  
 decrease concentration after diet- and physical activity-induced weight reduction, association with changes in plasma lipids, IL-6 and adiponectin, 359-365  
 effects of aerobic exercise, in adults, 1500-1507  
 effects of lifestyle modifications, 825-831  
 genotypes, effects on exercise training-induced changes in  $S_i$ , 453-460  
 high-sensitivity  
 associated with relative sympathetic overactivity, in NIDDM with MS, 1014-1021  
 versus IR, relation to FBG subgroups categorized by newly recommended criteria for fasting hyperglycemia in 10 059 healthy Koreans, 183-187  
 and plasma LDL-C, effect of atorvastatin in Japanese NIDDM patients, 67-71  
 relation to IR and glucose intolerance, 1232-1238  
 and serum adiponectin, in nonobese subjects with impaired glucose tolerance, effects of rosiglitazone, 439-444  
 and soluble adhesion molecules, relation to progression of SCI in patients with type 2 diabetes mellitus, 461-466  
 Creatinine clearance, effect of rigorous exercise, 106  
 CRF, *see* Chronic renal failure  
 Crigler-Najjar syndrome, type II, results of N-glucuronidation of retigabine from human UGTs, liver and kidney of, 711-721  
 CRP, *see* C-reactive protein  
 CS, *see* Cytosolic sialidase  
 CT, *see* Computed tomography  
 CURES, *see* Chennai Urban Rural Epidemiology Study  
 CuZnSOD, *see* Copper/zinc superoxide dismutase  
 ( $\alpha$ )-Cyclodextrin, new soluble dietary fiber, effects on weight gain and selected blood parameters in rats, 195-202  
 Cyclophilin B (CypB), and CAT, CuZnSOD, MnSOD, GCLc and GPx1, mRNA expressions and protein levels in limited amounts of tissue from rat retina, 168-174  
 CypB, *see* Cyclophilin B  
 Cytokines  
 proinflammatory, relationship with serum resistin, in obese women with IGT and normal glucose tolerance, 1495-1499  
 on sleep and energy balance, S2:33  
 Cytosolic sialidase (CS), effect of chronic ethanol consumption, in rat liver, 803-810  
 Dehydroepiandrosterone (DHEA), oral, endocrine effects of, in HIV-infected men, 858-870  
 Delivery, obstetric, and pregnancies, in LPI, hazards associated with, 224-231  
 2-Deoxyglucose (2DG) transport, increase in, accompanied by increased blood lactate concentration and decreased blood glucose concentration, effects of AICAR in skeletal muscle, 305  
 Deoxyribonucleic acid (DNA), damage, as oxidative stress marker in diabetes, S1:12  
 Dexamethasone, suppresses PLC activation and insulin secretion from isolated rat islets, 35-42  
 2DG, *see* 2-Deoxyglucose  
 DHA, *see* Docosahexaenoic acid  
 DHEA, *see* Dehydroepiandrosterone  
 Diabetes mellitus (DM)  
 in African Americans, utility of fasting plasma glucose in detection of progression to, 434-438  
 development in patients with distal pancreatectomy, major role of insulin secretory defect, 135-141  
 family history of, and glucose tolerance, autoimmunity,  $S_i$ , C-peptide and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188-194  
 $\alpha$ -glucosidase inhibitors in prevention of CHO-induced upregulation in intestinal sugar transport in mice with, 832-841  
 lipid and Lp profiles in Ethiopian patients with, 706-710  
 type 2, relation of CRP and soluble adhesion molecules to SCI progression in patients with, 461-466  
 Diabetic nephropathy, *see* Non-insulin dependent diabetes mellitus nephropathy  
 Diabetic retinopathy, *see* Non-insulin dependent diabetes mellitus retinopathy  
 Diet, 1582-1589  
 and exercise  
 body composition and metabolic effects of, on obese HIV-infected women, 1327-1336  
 in older adults with impaired glucose tolerance,  $S_i$  assessment indices versus euglycemic-hyperinsulinemic clamp data after, 525-532  
 hypercholesterolemia induced by, hypocholesterolemic action of acolbifene in intact and ovariectomized rats with, 605-613  
 maternal, for postnatal growth in rat, effect of Met imbalance in, 763-770  
 MS induced by, oxidative stress and NADPH oxidase dysregulation and antioxidant enzymes in, 928-934  
 and physical activity, weight reduction induced by, decrease CRP concentration after, association with changes in plasma lipids, IL-6 and adiponectin, 359-365  
 short-term, effect on MS in overweight children, 871-878  
 very low-energy changes induced by, among Korean female subjects, effects of UCP-3 haplotypes, 578-586

- Dietary fat, differential modulation of mRNA abundance of LDLR, HMG-CoA and MTP, in Golden-Syrian hamsters, 635-641
- Dietary fatty acid(s) (FAs), role in increased glyceroneogenesis in rats adapted to high-protein, CHO-free diet, 84-89
- Dietary fiber, new soluble, effects on weight gain and selected blood parameters in rats, 195-202
- Dietary glycotoxins, in healthy women and women with PCOS, short-term effects of orlistat, 494-500
- Dietary proteins  
     impact on WBPTO at rest in endurance-trained men, 501-507  
     maternal intake, programming of pregnant rat insulin axis and offspring body composition, 642-649  
     metabolized, in response to LoBAG diet in men with NIDDM, 247
- Dietary sesamin, and DHA and EPA, synergistic effect on gene expression of enzymes involved in hepatic peroxisomal fatty acid oxidation in rats, 381-390
- Differentiation, association with biologic events, in monocyte-derived macrophages, gliclazide inhibition of, 778-785
- 4,5-Dihydroxyhexanoic acid, and D-2 hydroxyglutaric acid, conversion of GBH to, in SSADH knockout mice, 356
- Dimethylarginine, *see* Asymmetric dimethylarginine
- Dimethylbiguanide, *see* Metformin
- DM, *see* Diabetes mellitus
- Docosahexaenoic acid (DHA), and dietary sesamin and EPA, synergistic effect on gene expression of enzymes involved in hepatic peroxisomal fatty acid oxidation in rats, 381-390
- N-6 Docosapentaenoic acid, increased flux to, in PLs of pancreas from *cfr*<sup>-/-</sup> knockout mice, 1192-1200
- Dopamine, neurochemicals on awake state, S2:16-17
- Dynein, apparent role in G6PDase trafficking in neutrophils from pregnant women, 279-281
- EAA, *see* Essential amino acid
- Eicosapentaenoic acid (EPA)  
     and dietary sesamin and DHA, synergistic effect on gene expression of enzymes involved in hepatic peroxisomal fatty acid oxidation in rats, 381-390  
     effects on early stage of NIDDM nephropathy, in KKA<sup>Y</sup>/Ta mice, 1590-1598
- Elderly subjects, *see* Older subjects
- Endothelial function, markers of, effect of atorvastatin in Japanese NIDDM patients, 69-70
- Endothelial nitric oxide synthase (eNOS), G894T polymorphism, prediction of glycemic status, in a 5-year study of Chinese subjects with IGT, 1155-1158
- Energy balance, and sleep, S2:30-35
- Energy (caloric) restriction (ER), prevents development of NIDDM, coordinated patterns of gene expression for energy metabolism determined by oligonucleotide microarray, 43-52
- Energy expenditure, effects of NO-1886 (ibrolipim) in obese ovariectomized rats, 154
- Energy metabolism, gene expression in insulin-sensitive tissues determined by oligonucleotide microarray, role of ER in development of NIDDM, 43-52
- Energy surplus, interactions with short-term exercise, on glucose and insulin responses in healthy people with induced, mild S<sub>i</sub>, 402-408
- EPA, *see* Eicosapentaenoic acid
- Epidemiology, sulfonylureas on risk of MI, S1:17-18
- Epididymal fat pad mass, effects of HF feeding in Sprague-Dawley rat, 209
- Epinephrine, increased lipolysis in IDDM, effect of antecedent hypoglycemia, 371-380
- 8-Epi-prostaglandin F<sub>2α</sub>, leptin as determinant of, 918-922
- ER, *see* Energy restriction
- Erectile dysfunction, *see* Impotence
- E-selectin, and CRP, relation to progression of SCI in patients with type 2 diabetes mellitus, 461-466
- Essential amino acid (EAA), liquid CHO, ingestion, during short-term resistance exercise, effect on myofibrillar protein degradation, 570-577
- Essential fatty acid (EFA), brain, impact of maternal folic acid supplementation, in adult offsprings, 628-634
- Essential hypertension, *see* Hypertension
- Estrogen, treatment on arterial distensibility, fibrinolysis and metabolic profile, in postmenopausal women with NIDDM, 953-959
- Estrogen receptor modulators, acolbifene, hypocholesterolemic action, in intact and ovariectomized rats with diet-induced hypercholesterolemia, 605-613
- Ethanol, *see* Alcohol
- Ethiopian subjects, with DM, lipid and Lp profiles in, 706-710
- Euglycemic-hyperinsulinemic clamp data, versus S<sub>i</sub> assessment indices, after dietary and exercise intervention in older adults with impaired glucose tolerance, 525-532
- Exercise  
     adiponectin and stress hormone responses to maximal sculling after volume-extended training season in elite rowers, 13-19  
     and diet  
         effects on body composition and metabolism of obese HIV-infected women, 1327-1336  
         in older adults with impaired glucose tolerance, S<sub>i</sub> assessment indices versus euglycemic-hyperinsulinemic clamp data after, 525-532  
         weight loss from, effects on LDL particle size and distribution in obese women, 1302-1307



- Exercise (continued)  
 differential effects of palmitic and oleic acid dietary intake on oxygen consumption during and after, 1215-1221  
 effect on resting MR, in postmenopausal African American and Caucasian women, 1358-1364  
 effects on MS in overweight children, 871-878  
 insulin-stimulated IRS-2–associated PI3-K in SM after, 1046-1052  
 and isoflavones, effects on bone and lipid metabolism in PM Japanese women, 423-433  
 long-term, stimulation of AMPK activity and subunit expression, in VAT and liver, 1122-1128  
 moderate intensity, and AICAR, extent of  $\alpha 2$  AMPK activation by, 303-304  
 in NIDDM subjects, influence of glycemic control on pulmonary function and heart rate in response to, 1532-1537  
 rigorous, plasma concentrations and urinary excretion of purine bases and oxypurinol after, 103-107  
 short-term, interactions with energy surplus, on glucose and insulin responses in healthy people with induced, mild  $S_i$ , 402-408  
 $S_i$  changes induced by, effects of CRP genotypes on, 453-460
- Exercise training  
 effect on endothelial function and biochemical vascular markers, in first-degree relatives of NIDDM patients, 1508-1515  
 and PSE supplementation, effects on blood lipid levels, 541-549
- Ezetimibe  
 effect on subtype distribution in patients treated by LDL apheresis and statins, 599-604  
 Lp effects of, alone, and with colessevelam hydrochloride, in hypercholesterolemic subjects, 1697-1703
- FA, *see* Fatty acid(s)
- FABP2, *see* Fatty acid–binding protein 2
- Family history, of diabetes, and glucose tolerance, autoimmunity,  $S_i$ , C-peptide and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188-194
- FAs, *see* Fatty acid(s)
- Fasting  
 and alcohol exposure, skeletal muscle and liver oxysterols during, 119-127  
 effect on NEFA turnover in mice, 1481-1487  
 and refeeding in rats,  $\beta$ -adrenergic activity in modulation of GLUT4 during, 1538-1545
- Fasting blood glucose (FBG)  
 categorized by newly recommended criteria for fasting hyperglycemia, relation to IR and serum high-sensitivity CRP levels in 10 059 healthy Koreans, 183-187  
 elevated 2-hour postload plasma glucose as alternative to, criterion for MS, 1323-1326
- Fasting blood glucose (FBG) (continued)  
 impaired, lower limit in Korea, 489-493
- Fasting hyperglycemia, FBG subgroups categorized by newly recommended criteria for, relation to IR and serum high-sensitivity CRP levels in 10 059 healthy Koreans, 183-187
- Fasting plasma glucose, utility in detection of prediabetes in African Americans, 434-438
- Fat accumulation, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154
- Fatty acid(s) (FAs)  
 dietary, role in increased glyceroneogenesis in rats adapted to high-protein, CHO-free diet, 84-89  
 uptake and metabolism, glucose and, effects of short- and long-term leptin treatment on, in HL-1 cardiomyocytes, 1067-1075
- Fatty acid–binding protein 2 (FABP2), gene, association of MS and hypertriglyceridemia on Ala54Thr variant of, 1222-1226
- Fatty acid (FA) transporter enzymes, mRNA levels of, effects of NO-1886 in ovariectomized rats, 154
- Fatty acid oxidation, hepatic peroxisomal, synergistic effect of dietary sesamin, DHA and EPA on gene expression of enzymes involved in, 381-390
- Fatty acid uptake, effects of insulin and leptin in differentiated 3T3-L1 adipocytes, 8-12
- FBCx, new soluble dietary fiber, effects on weight gain and selected blood parameters in rats, 195-202
- FBG, *see* Fasting blood glucose
- Fecal lipid content, effects of FBCx in high fat-fed rats, 199
- Fenofibrate, and lipid metabolism, in AT of rats, 731-735
- FFAs, *see* Free fatty acids
- Fibrinolysis  
 estrogen treatment on, in postmenopausal women with NIDDM, 953-959  
 with IGT, effect of acarbose, 935-939
- FoF1-ATPase mRNA levels, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154-155
- Folate, and choline, different effects on plasma kinetics of methionine and homocysteine, in rats, 899-906
- Folic acid, maternal supplementation, impact on brain fatty acid levels in adult offspring, 628-634
- Food intake  
 and BW, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154  
 effects of FBCx in high fat-fed rats, 198
- Forearm, effects of hypoglycemia and epinephrine in IDDM, 376-377
- Free cortisol  
 and aldosterone, urinary excretion of, in response to LoBAG diet in men with NIDDM, 248  
 urinary, and aldosterone, and plasma renin, in response to LoBAG diet in men with NIDDM, 248
- Free fatty acids, *see* Nonesterified fatty acids

- Free levorotatory thyroxine, serum, in response to LoBAG diet in men with NIDDM, 248
- Fu5AH cell system, altered lipid, Apo and Lp in IBD as consequences on cholesterol efflux capacity using, 980-988
- Fungata study, correlation between change in BW rather than current BW and change in serum adiponectin levels in Japanese population, 324-330
- G6PDase, *see* Glucose-6-phosphate dehydrogenase
- G894T polymorphism, eNOS, prediction of glycemic status, in a 5-year study of Chinese subjects with IGT, 1155-1158
- GADA, *see* Glutamic acid decarboxylase autoantibody
- Gamma ( $\gamma$ )-hydroxybutyric acid (GHB)  
 metabolic transformations intragastrically administered to baboons, 356  
 metabolism to D-2-hydroxyglutarate in mammals, evidence for D-2-hydroxyglutarate transhydrogenase, 353-358
- 308G/A polymorphism, of TNF- $\alpha$ , relationship with LTA 804C/A and 252A/C polymorphism, in Japanese subjects with NIDDM retinopathy, 1406-1410
- GCLc, *see* Glutamylcysteine ligase, catalytic subunit
- GCT, *see* Glucose challenge test
- Gender, *see* Sex
- Gene polymorphism, *see* Polymorphism
- Genetics  
 sleep disorders, S2:8  
 of sleep-wake cycle and its disorders, S2:7-12
- Genotypes, of CRP, effects on exercise training–induced changes in S<sub>i</sub>, 453-460
- Gestational diabetes mellitus (GDM), IRS-1 and  $\beta_3$ -ADR genes in, 1451-1456
- GFR, *see* Glomerular filtration rate
- GH, *see* Growth hormone
- GHB, *see* Gamma ( $\gamma$ )-hydroxybutyric acid
- Ghrelin  
 5' flanking region, sequencing analysis of, in relations between sequence variants and BMI, 1420-1425  
 total and octanoylated levels, effect of alcohol ingestion, in healthy levels, 1625-1629
- Gliclazide  
 inhibition of differentiation-associated biologic events, in monocyte-derived macrophages, 778-785  
 protection of 3T3L1 adipocytes against IR–induced hydrogen peroxide, with restoration of GLUT4 translocation, 722-730  
 protective role on oxidative stress-induced beta-cell damage, S1:13  
 on vascular complications of diabetes, effects of, S1:10-13  
 in vitro and in vivo antioxidant action of, S1:12-13
- Glimepiride  
 and glyburide, effects on glucose counterregulation and recovery from hypoglycemia, 78-83
- Glimepiride (continued)  
 with rosiglitazone, impact on beta-cell function, IR, and adiponectin concentrations, 20-25
- Gln, *see* Glutamine
- Glomerular filtration rate (GFR)  
 Cockcroft-Gault formula is biased by body weight in diabetic patients with renal impairment, 108-112  
 relation to plasma SAH concentration in children, 252-257
- Glucocorticoid receptor (GR)  
 in malnutrition and acute infection, effects of cortisol, 550-554  
 selective antagonism of, reduction of HGP, 1255-1262
- Glucocorticoids (GCs), SM atrophy induced by, Gln in prevention of, 1239-1247
- Glucophage, diet, relationship between changes in PAI-1 activity and first-trimester miscarriage in women with PCOS on, 349
- Glucose  
 blood, *see* Blood glucose  
 counterregulation, and recovery from hypoglycemia, effects of glimepiride and glyburide, 78-83  
 and FA uptake and metabolism, effects of short- and long-term leptin treatment on, in HL-1 cardiomyocytes, 1067-1075  
 impaired fasting, lower limit in Korea, 489-493  
 ingestion, effect on NF- $\kappa$ B, I $\kappa$ B and TNF- $\alpha$  mRNA in MNCs, in healthy subjects, 1177-1185  
 and insulin, in response to epinephrine in IDDM, 375-376  
 metabolism, effect of telmisartan, in nondiabetic insulin-resistant subjects, 1149-1154  
 plasma, *see* Plasma glucose  
 in response to weight reduction after diet and physical activity, 361-362  
 use, and insulin content of dexamethasone-pretreated islets, 39
- Glucose-6-phosphate dehydrogenase (G6PDase),  
 trafficking, in neutrophils from pregnant women, apparent role of dynein, 279-281
- Glucose challenge test (GCT), association between insulin and adipokines and maternal body size and prediction of birth weight, 339-344
- Glucose intolerance  
 age-related, effects of acarbose on, 533-540  
 serum levels of IL-6, CRP, VCAM-1 and MCP-1 in relation to, 1232-1238
- Glucose metabolism, and inflammatory marker and BP, relations to IMT in childhood obesity, 113-118
- Glucose regulation, normal, association of MS and IR with cardiovascular risk in adults with, 445-452
- Glucose responses, and insulin responses, in healthy people with induced, mild S<sub>i</sub>, interactions between energy surplus and short-term exercise on, 402-408
- Glucose tolerance (GT)  
 factors responsible for deterioration in newly diagnosed Japanese NIDDM men, 53-58

- Glucose tolerance (GT) (continued)  
 and family history of diabetes, autoimmunity,  $S_i$ , C-peptide and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188-194  
 impaired, *see* Impaired glucose tolerance  
 pregnancy-induced adaptations in, effect of aging in rats, 409-414
- Glucose transport  
 in adipocytes, effect of Trf and Fe on IR of, 1042-1045  
 effect of troglitazone, 313-314  
 and GLUT4, effects of activation of  $\alpha 2$  AMPK by AICAR at physiological levels in mouse skeletal muscle, 300-308  
 in skeletal muscle, effects of HF feeding in Sprague-Dawley rat, 209  
 whole-body, and insulin tolerance, effect of repeated AICAR injection, 306
- Glucose transporter 4 (GLUT4)  
 content in arm and leg muscle, and  $S_i$ , effects of repeated prolonged whole-body low-intensity exercise, 217-223  
 expression, and glucose transport, effects of activation of  $\alpha 2$  AMPK by AICAR at physiological levels in mouse skeletal muscle, 300-308  
 during fasting and refeeding in rats,  $\beta$ -adrenergic activity in modulation of, 1538-1545  
 translocation, gliclazide in restoration of, 722-730
- GLUT4, *see* Glucose transporter 4
- Glutamic acid decarboxylase autoantibody (GADA), clinical and biochemical characteristics of nonobese NIDDM patients with, 1107-1112
- Glutamine (Gln)  
 neurochemicals on awake state, S2:15-16  
 prevention of GC-induced SM atrophy associated with myostatin suppression, 1239-1247
- Glutamylcysteine ligase, catalytic subunit (GCLc)  
 and CAT, CuZnSOD, MnSOD, Gpx1 and CypB, mRNA expressions and protein levels in limited amounts of tissue from rat retina, 168-174
- Glutamylcysteine ligase (GCLc)  
 catalytic subunit, expression in rat retinal ischemia after reperfusion, 892-898
- Glutathione peroxidase (GPx1)  
 and CAT, CuZnSOD, MnSOD, GCLc and CypB, mRNA expressions and protein levels in limited amounts of tissue from rat retina, 168-174  
 oxidative stress with diet-induced MS due to, 928-934
- Glyburide, and glimepiride, effects on glucose counter-regulation and recovery from hypoglycemia, 78-83
- Glycemia, increasing, association of vascular risk factors with, in older Chinese population, 1035-1041
- Glycemic control of NIDDM  
 influence of chromium chloride-containing milk, 923-927  
 influence on pulmonary function and heart rate, in response to exercise, 1532-1537  
 overcoming hurdles for, S1:6-9
- Glyceroneogenesis, increased in AT from rats adapted to high-protein, CHO-free diet, role of dietary FAs, 84-89
- Glycogen, liver concentration, in LETO and OLETF rats, 319
- Glycotoxins, dietary, in healthy women and women with PCOS, short-term effects of orlistat, 494-500
- G-protein  $\beta 3$  subunit (GNB3), C1429T polymorphism in, as a risk factor of hypercholesterolemia in Japanese males, 751-757
- GPx1, *see* Glutathione peroxidase
- GR, *see* Glucocorticoid receptor
- Granulocytes, oxidizing and reducing responses, from NIDDM, 1426-1428
- Growth hormone (GH)  
 and IGF-1, serum levels, in response to LoBAG diet in men with NIDDM, 247-248  
 responses to hypoglycemia, in NIDDM, 593-598  
 response to maximal sculling after volume-extended training season in elite rowers, 13-19
- GT, *see* Glucose tolerance
- HAART, *see* Highly active antiretroviral therapy
- HCTL, *see* Homocysteine thiolactonase
- Hcy, *see* Homocysteine
- HDL, *see* High-density lipoprotein(s)
- HDL-C, *see* High-density lipoprotein-cholesterol
- Heart  
 atrophy and hypertrophy, differential changes in respiratory capacity and ischemia tolerance of mitochondria from, 1097-1106  
 individual sulfonylureas and, S1:16-17
- Heart disease, *see* Coronary heart disease
- Heart rate (pulse rate; HR)  
 in NIDDM, *see* Heart rate in NIDDM  
 in NIDDM, influence of glycemic control pulmonary function and, in response to exercise, 1532-1537  
 sleep-related clinical observations on, S2:46-47
- Heat shock protein (HSP), SM, effect of alcohol, 843-851
- Heme oxygenase-2 (HO-2), oxidative stress with diet-induced MS due to, 928-934
- Hemodialysis (HD), hyperhomocysteinemia a significant risk factor for silent cerebral infarction in CRF patients undergoing, 656-661
- Hemodynamics, regulation, and ADMA, in middle-aged men, 771-777
- Hepatic enzymes, involved in peroxisomal fatty acid oxidation, synergistic effect of dietary sesamin, DHA and EPA on gene expression of, 381-390  
 activity of, 382-386
- Hepatic glucose production (HGP), selective antagonism of GR in reduction of, 1255-1262
- Hepatic lipase  
 effects of PSE supplementation and aerobic exercise training, 541-549  
 variant mRNA of, transient induction by corticotropin hormone in rat adrenals, 467-477

- Hepatic lipase gene (LIPC), variation, association with Lp subclass and particle size differences in Afro-Caribbeans, African Americans and white Americans, 96-102
- Hepatic pyruvate dehydrogenase kinases, *see* Pyruvate dehydrogenase kinases
- Hepatic stellate cells, non-immortalized human, effect of physiologic concentrations of leptin on collagen by, 1317-1322
- Hepatic synthesis, and plasma cholesterol levels and intestinal absorption, effects of stigmaterol in rat, 292-299
- Hepatitis, T cell-mediated, role of TNF- $\alpha$  in modulation of adiponectin levels in mice, 555-559
- Hepatocytes, nonmalignant human cell line, TNF- $\alpha$ -induced production of PAI-1 in regulation, in nonmalignant hepatocyte cell line, 1464-1472
- HepG2 cell line, and Caco-2 cell line, effect of ascorbic acid and  $\alpha$ -tocopherol on down-regulation of Apo A-I gene expression in, 159-167
- HF diet, *see* High-fat diet
- HF/LCD, *see* High-fat/low carbohydrate diet
- High-density lipoprotein(s) ( $\alpha$ -lipoprotein; HDL)  
antioxidant capacity of, correlation with severity of CV disease with NIDDM, 1201-1206  
effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26-34  
subclasses, relationship between TC/HDL-C and TG/HDL-C, 1141-1148
- High-density lipoprotein-cholesterol (HDL-C;  $\alpha$ -lipoprotein-cholesterol)  
adiponectin in relation to, in healthy children, 1171-1176  
effects of PSE supplementation and aerobic exercise training, 541-549  
HIV infection, effect of disease versus effect of treatment on, 90-95  
ratio with TC and TG, and HDL subclasses, relationship between, 1141-1148
- High-fat (HF) diet, effects on components of CAP/Cb1 signaling cascade in Sprague-Dawley rat skeletal muscle, 203-212
- High-fat/low carbohydrate diet (HF/LCD), regulation of glucose metabolism via long-term transcriptional loop, 1457-1463
- Highly active antiretroviral therapy (HAART), for HIV infection, effect of disease versus effect of treatment on HDL-C levels, 90-95
- High oleic canola (HOC) oil, and PS mixed with M TGs, effect on plasma lipids in overweight men, 391-395
- High-protein, carbohydrate-free diet, role of dietary FAs in increased glyceroneogenesis in rats adapted to, 84-89
- High-protein, low carbohydrate (CHO) diet, metabolic response in men with NIDDM, 243-251
- Hip circumference, relative to BMI, impact on cardiovascular risk factors, 563-569
- Histamine, neurochemicals on awake state, S2:16
- HIV, *see* Human immunodeficiency virus
- HOC oil, *see* High-oleic canola oil
- Homeostasis, interactive, on sleep and energy balance, S2:30-35
- Homocysteine (Hcy)  
association with insulin, in Swedish population, 1007-1013  
clinical value in hospitalized elderly, 620-627  
correlation with IR and low-grade systemic inflammation in obese prepubertal children, 72-77  
different effects choline and folate on plasma kinetics, in rats, 899-906  
plasma, relation to plasma SAH concentration and GFR in children, 252-257  
*see also* Hyperhomocysteinemia
- Homocysteine thiolactonase (HCTL), correlation with severity of CV disease with NIDDM, 1201-1206
- Hong Kong, prevalence and risk factors for androgen deficiency in middle-aged men, 1488-1494
- Hong Kong Cardiovascular Risk Factor Study, impact of sex-specific body composition on cardiovascular risk factors, 563-569
- Hormonal responses  
to brief hypoglycemia in PM women and age-matched men with NIDDM, 331-338  
to resistance training overreaching, effects of AA supplementation, 282-291
- Hormone replacement therapy (HRT), combined with alendronate, effects on osteoporotic postmenopausal Chinese women, 741-747
- HPA, *see* Hypothalamic-pituitary-adrenal axis
- HRT, *see* Hormone replacement therapy
- HSP, *see* Heat shock protein
- HTG, *see* Hypertriglyceridemia
- HTN, *see* Hypertension
- Human immunodeficiency virus (HIV)  
body composition and metabolic effects of diet and exercise in obese women with, 1327-1336  
men with, endocrine effects of DHEA, 858-870  
treated by ART, lipodystrophy and MS in, 940-945
- Human immunodeficiency virus (HIV) infection  
effect of disease versus effect of treatment on HDL-C levels, 90-95  
with lipodystrophy, association between TNF- $\alpha$  and insulin-mediated suppression of free fatty acids and net lipid oxidation in, 175-182
- Hydrogen peroxide, IR-induced, gliclazide in protection of 3T3L1 adipocytes against, 722-730
- Hydroimidazolone, methylglyoxal-derived, increased serum levels, association with retinopathy in NIDDM, 232-236
- 8-Hydroxy-2'-deoxyguanosine, as oxidative stress marker in diabetes, S1:12
- 3-Hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase, mRNA abundance of, effect of dietary fatty acids, 635-641



- D-2-Hydroxyglutarate, metabolism from GHB in mammals, evidence for D-2-hydroxyglutarate transhydrogenase, 353-358
- D-2-Hydroxyglutarate transhydrogenase, metabolism of GHB to D-2-hydroxyglutarate in mammals, evidence for, 353-358
- Hypercholesterolemia  
 diet-induced, hypocholesterolemic action of acolbifene in intact and ovariectomized rats with, 605-613  
 effect of atorvastatin in Japanese NIDDM patients, 67-71  
 IR, adipokines and oxidative stress in, 918-922  
 in Japanese males, GNB3 polymorphism C1429T and LRP5 polymorphism A1330V as risk factors for, 751-757  
 Lp effects of ezetimibe alone, and with colesvelam hydrochloride, in subjects with, 1697-1703  
 use of LDL apheresis and statins for, effect of ezetimibe on LDL subtype distribution in, 599-604
- Hyperglycemia  
 acute, effect on PMN functions, in healthy subjects, 811-818  
 altered release of TNF- $\alpha$  from MNCs of obese reproductive-age women during, 271-276  
 association with IR, in STZ-DM pigs, 960-971  
 and diabetic complications, sulfonylureas for, S1:2-3  
 factors responsible for, in newly diagnosed Japanese NIDDM men, 53-58  
 postprandial  
   CV disease and, 1264-1266  
   mechanisms of, 1264  
   mechanisms of vascular damage by, 1266-1267  
   targeting of, 1263-1281  
   treatment approaches, 1267-1270  
 progressive, sulfonylureas for, S1:2-3  
 relationship between adiponectin-leptin ratio and IR parameters in subjects without, 1248-1254
- Hyperhomocysteinemia  
 correlation with IR and low-grade systemic inflammation in obese prepubertal children, 72-77  
 link with MS, association between insulin and Hcy leading to, 1007-1013  
 as a risk factor in erectile dysfunction, 1564-1568  
 as significant risk factor for silent cerebral infarction in CRF patients undergoing HD, 656-661
- Hyperinsulinemia  
 acute, effect on PMN functions, in healthy subjects, 811-818  
 fasting, simple anthropometric measures, in Asian Indian adolescents, 1569-1573
- Hypertension (HTN)  
 effect of telmisartan on S<sub>i</sub> in nondiabetics with, 1159-1164  
 IR in, quantified by OGTT, comparison of methods, 143-150  
 offspring of parents with, metabolic characteristics of soleus muscle in relation to insulin action in, 1388-1396
- Hypertension (HTN) (continued)  
 relation to IR, angiotensin II infusion decreases plasma adiponectin via its type 1 receptor in rats, 478-488
- Hypertriglyceridemia (HTG)  
 association with Ala54Thr variant of FABP2 gene, in urban South Indians, 1222-1226  
 effects of raloxifene on lipid and Lp in postmenopausal women with and without, 972-979  
 mild, modification of LDL features during oral fat load in subjects and LDLR binding affinity with, 1308-1316
- Hypertriglyceridemic clamp technique, for long-chain and structured TG emulsions, in healthy subjects, 1443-1450
- Hypertrophy, differential changes in respiratory capacity and ischemia tolerance of mitochondria of heart with, 1097-1106
- Hypoadiponectinemia  
 association with CV risk factors, in nonobese healthy adults, 1546-1550  
 association with early onset CHD and multiple coronary stenoses, in men, 1653-1657
- Hypoalphalipoproteinemia, in HIV infection, effect of disease versus effect of treatment, 90-95
- Hypocholesterolemia, as an effect of acolbifene, in intact and ovariectomized rats with diet-induced hypercholesterolemia, 605-613
- Hypocretins  
 biochemistry, S2:37  
 and narcolepsy, S2:36-39  
 role in metabolic mechanism of wakefulness and sleep, S2:24-29  
 on sleep and energy balance, S2:32-33
- Hypocretins-orexins, neurochemicals on awake state, S2:16
- Hypoglycemia  
 antecedent, effect on increased epinephrine-induced lipolysis in IDDM, 371-380  
 brief, hormonal, subjective and neurocognitive responses to, in PM women and age-matched men with NIDDM, 331-338  
 multiple defects in counterregulation of, in advanced NIDDM, 593-598  
 recovery from, and glucose counterregulation, effects of glimepiride and glyburide, 78-83  
 sulfonylurea-induced, S1:21-22
- Hypothalamic-pituitary-adrenal (HPA) axis, activity in NIDDM, role of autonomic balance in, 1135-1140
- Hypothalamus  
 mRNA profiles, of female Zucker diabetic fatty rats after topiramate treatment, 1411-1419  
 neurophysiology and essential brain region for sleep  
   brain stem, S2:4-5  
   lateral, S2:4  
   pineal gland, S2:4  
   suprachiasmatic nucleus, S2:4  
   tuberomammillary nucleus, S2:4  
   ventrolateral preoptic nucleus, S2:4

- Hypoxanthine, and oxypurinol, plasma concentrations and urinary excretion after rigorous exercise, 103-107
- Hypoxia  
acute, persistent suppression of REE after, 669-675  
nocturnal, pathophysiology of, S2:47
- Ibrolipim, NO-1886, effects on expression of UCP3 mRNA and obesity in ovariectomized rats, 151-158
- ICAM-1, *see* Intercellular adhesion molecule-1
- IDDM, *see* Insulin-dependent diabetes mellitus
- IDF, *see* International Diabetes Federation
- IGF-1, *see* Insulin-like growth factor-1
- IGT, *see* Impaired glucose tolerance
- I $\kappa$ B, *see* Inhibitor  $\kappa$ B
- IL-1Ra, *see* Interleukin-1 receptor antagonist
- IL-6, *see* Interleukin-6; Plasma interleukin-6
- IL-18, *see* Interleukin-18
- IMCL, *see* Intramyocellular lipid
- Impaired glucose tolerance (IGT)  
effect of acarbose  
on fibrinolytic activity with, 935-939  
on LDL atherogenicity with IGT, 946-952  
effects of rosiglitazone on serum adiponectin and CRP in nonobese subjects with, 439-444  
eNOS G894T polymorphism in prediction of glycemic status in a 5-year study of Chinese subjects with, 1155-1158  
and NIDDM and MS, soluble P-selectin and sCD40L levels in subjects with (Chennai Urban Rural Epidemiology Study), 237-242  
relationship between serum resistin and proinflammatory cytokines in obese women with, 1495-1499  
S<sub>i</sub> assessment indices versus euglycemic-hyperinsulinemic clamp data after diet and exercise intervention in older adults with, 525-532
- Impotence, hyperhomocysteinemia as a risk factor for, 1564-1568
- IMT, *see* Intima-media thickness
- Infection, acute, and malnutrition, effects of cortisol on glucocorticoid receptor in, 550-554
- Inflammation, and glucose tolerance, family history of diabetes, autoimmunity, S<sub>i</sub> and C-peptide, in relation to serum adiponectin in population sample of 64-year-old women, 188-194
- Inflammatory bowel disease (IBD), altered lipid, Apo and Lp in, 980-988
- Inflammatory markers  
correlation with hyperhomocysteinemia in obese prepubertal children, 72-77  
effect of atorvastatin in Japanese NIDDM patients, 69-70  
and glucose metabolism and BP, relations to IMT in childhood obesity, 113-118
- Inhibitor  $\kappa$ B (I $\kappa$ B), in MNCs, effect of glucose ingestion, in healthy subjects, 1177-1185
- Inositol phosphate (IP), accumulation in dexamethasone-pretreated islets, 39
- Insomnia  
psychiatric causes, S2:42  
symptoms, durations and causes, S2:41-42
- Insulin  
and adipokines, association with maternal body size and prediction of birth weight, 339-344  
content, and glucose use of dexamethasone-pretreated islets, 39  
effect on intracellular oxidative stress in patients with NIDDM, 691-695  
and glucose, in response to epinephrine in IDDM, 375-376  
and glucose, responses in healthy people with induced, mild S<sub>i</sub>, interactions between energy surplus and short-term exercise on, 402-408  
glucose transport stimulated by, effect of repeated AICAR injection, 305-306  
and leptin, effects on fatty acid uptake in differentiated 3T3-L1 adipocytes, 8-12  
plasma, *see* Plasma insulin  
in response to weight reduction after diet and physical activity, 361-362  
serum, *see* Serum insulin  
suppression of free fatty acids and net lipid oxidation mediated by, association with TNF- $\alpha$  in HIV-infected patients with lipodystrophy, 175-182
- Insulin absorption, as a major factor in apparent IR and control of NIDDM, 614-619
- Insulin action  
in lean and reduced-obese subjects, effects of short-term overfeeding, 1207-1214  
metabolic characteristics of soleus muscle in relation to, in offspring of parents with HTN, 1388-1396
- Insulin-dependent diabetes mellitus (IDDM; type 1 diabetes mellitus)  
early, sympathetic sudomotor disturbance in, linked to lipid peroxidation, 1524-1531  
effect of antecedent hypoglycemia on increased epinephrine-induced lipolysis in, 371-380  
with renal impairment, Cockcroft-Gault formula is biased by body weight in, 108-112
- Insulin-like growth factor-1 (IGF-1), and GH, in response to LoBAG diet in men with NIDDM, 247-248
- Insulin receptor substrate-1 (IRS-1), polymorphisms, in GDM and normal pregnancy, 1451-1456
- Insulin receptor substrate-2 (IRS-2), insulin stimulated, associated with PI3-K, in SM after exercise, 1046-1052
- Insulin release and secretion  
defect, major role in development of diabetes mellitus in patients with distal pancreatectomy, 135-141  
effects of glimepiride and glyburide on rate of, 80-81  
factors responsible for deteriorating glucose tolerance in Japanese NIDDM men, 53-58  
PLC activation and, effect of dexamethasone in isolated rat islets, 35-42

**Insulin resistance (IR)**

- apparent, insulin absorption as a major factor in, 614-619
- association with hyperglycemia, in STZ-DM pigs, 960-971
- CV disease risk, abdominal adiposity circulating plasma adiponectin and, association with SNP276>T versus SNP45T>G adiponectin gene polymorphism in nonobese/nondiabetic Korean men, 59-66
- gliclazide in protection of 3T3L1 adipocytes against hydrogen peroxide induced by, 722-730
- glucose transport in adipocytes, effect of Trf and Fe, 1042-1045
- in hypertension, quantified by OGTT, comparison of methods, 143-150
- and IL-6, adiponectin and leptin in nonobese Japanese NIDDM patients, 258-262
- and impaired GT, NIDDM and MS, soluble P-selectin and sCD40L levels in subjects with (Chennai Urban Rural Epidemiology Study), 237-242
- influence of AGT polymorphism (Met235Thr), in obese Japanese women, 819-824
- and low sympathetic nerve activity in Tzamura Suzuki obese diabetic mouse, as a new model for NIDDM and obesity, 1664-4669
- and MS, association with cardiovascular risk in adults with normal glucose regulation, 445-452
- in NIDDM, beta cell-function, adiponectin concentrations and, impact of rosiglitazone combined with glimepiride, 20-25
- in nondiabetic hypercholesterolemic patients, 918-922
- nondiabetic subjects with, effect on telmisartan on glucose and lipid metabolism, 1149-1154
- in obesity, and low-grade systemic inflammation, correlation with hyperhomocysteinemia in prepubertal children, 72-77
- parameters, relationship with adiponectin-leptin ratio, in subjects without hyperglycemia, 1248-1254
- relationship with LDL particle size, in nondiabetic Koreans, 1610-1615
- relation to hypertension, angiotensin II infusion decreases plasma adiponectin via its type 1 receptor in rats, 478-488
- versus serum high-sensitivity CRP levels, relation to FBG subgroups categorized by newly recommended criteria for fasting hyperglycemia in 10 059 healthy Koreans, 183-187
- serum levels of IL-6, CRP, VCAM-1 and MCP-1 in relation to, 1232-1238
- sustainability by metformin-diet sustainability by metformin-diet over 4 years in women with PCOS, 1582-1589

**Insulin sensitivity ( $S_i$ )**

- in adult twins, interrelation with birth weight, 561-562
- effect of telmisartan, in nondiabetics with HTN, 1159-1164

**Insulin sensitivity ( $S_i$ ) (continued)**

- effect of weight loss, in overweight adults, 907-911
- effects of activation of  $\alpha 2$  AMPK by AICAR in mouse skeletal muscle, 300-308
- exercise training—induced changes in, effects of CRP genotypes on, 453-460
- factors responsible for deteriorating glucose tolerance in Japanese NIDDM men, 53-58
- and glucose effectiveness during SBT, versus minimal model analysis of IVGTT, 676-690
- and glucose tolerance, family history of diabetes, autoimmunity, C-peptide and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188-194
- induced, mild, interactions between energy surplus and short-term exercise on glucose and insulin responses in healthy people with, 402-408
- and limb muscle adaptations, effects of repeated prolonged whole-body low-intensity exercise, 217-223
- major role of insulin secretory defect in development of diabetes mellitus in patients with distal pancreatectomy, 135-141
- in nonobese subjects with impaired glucose tolerance, effects of rosiglitazone on, 439-444
- in SM, AT and liver of extremely obese subjects with acanthosis nigricans, 1658-1663
- Insulin sensitivity indices (ISI)
  - versus euglycemic-hyperinsulinemic clamp data, after dietary and exercise intervention in older adults with impaired glucose tolerance, 525-532
  - in hypertension, quantified by OGTT, comparison of methods, 143-150
- Insulin signaling, effects of HF feeding in Sprague-Dawley rat skeletal muscle, 209-210
- Insulin therapy
  - intensive, in reduction of small dense LDL particles in patients with NIDDM, 879-884
  - sulfonylureas plus, for NIDDM, S1:4
  - see also* Glycemic control of NIDDM
- Insulin tolerance, and whole-body glucose tolerance, effect of repeated AICAR injection, 306
- Interleukin-1 receptor antagonist (IL-1Ra), adipocytes versus non-fat cells in release of, 1113-1121
- Interleukin-6 (IL-6)
  - plasma
    - elevation in PCOS independent of obesity and sleep apnea, 1076-1082
    - and AT gene expression, effect of aerobic training, in obese women, 1375-1381
    - and lipids and adiponectin, association with decrease CRP concentration after diet- and physical activity-induced weight reduction, 359-365
  - serum
    - and adiponectin, leptin and IR in nonobese Japanese NIDDM patients, 258-262
    - in relation to IR and glucose intolerance, 1232-1238

- Interleukin-18 (IL-18), adipocytes versus non-fat cells in release of, 1113-1121
- International Diabetes Federation (IDF), definition of MS in Chinese subjects older than 30 years with NIDDM, 1088-1096
- Intestinal absorption, and plasma cholesterol levels and hepatic synthesis, effects of stigmasterol in rat, 292-299
- Intima-media thickness (IMT)  
relationship with LDL particle size, in nondiabetic Koreans, 1610-1615  
relations to inflammatory marker, glucose metabolism and BP in childhood obesity, 113-118
- Intracellular trafficking, of G6PDase in neutrophils from pregnant women, apparent role of dynein, 279-281
- Intramycellular lipid (IMCL), relation to CV/NIDDM risk factors in healthy men versus women subjects, 128-134
- Intrauterine growth retardation, associated with pregnancies and deliveries in LPI, 224-231
- Intravenous glucose tolerance test (IVGTT), minimal model analysis of, versus SBT of  $S_i$  and glucose effectiveness, 676-690
- Inuit of Nunavik, dietary contaminants in oxidative stress in, 989-995
- IP, *see* Inositol phosphate
- IR, *see* Insulin resistance
- Iron (Fe), effect on IR of glucose transport, in adipocytes, 1042-1045
- IRS-1, *see* Insulin receptor substrate-1
- IRS-2, *see* Insulin receptor substrate-2
- Ischemia, tolerance, and respiratory capacity, differential changes in mitochondria from atrophied and hypertrophied hearts, 1097-1106
- Ischemia-reperfusion injury, antioxidant enzymes expression in rat retina induced by, 892-898
- Ischymetric mechanism, on wakefulness and sleep controlling, S2:26-27
- ISI, *see* Insulin sensitivity indices
- Islets of Langerhans, *see* Beta ( $\beta$ ) cell(s) (islets of Langerhans)
- Isoflavones, and exercise, effects on bone and lipid metabolism in PM Japanese women, 423-433
- IVGTT, *see* Intravenous glucose tolerance test
- Japanese NIDDM subjects  
*Arg121Trp* variant, in *PAX4* gene, association with beta-cell dysfunction in, 213-216  
atorvastatin lowers plasma LDL-C and CRP in, 67-71  
men  
factors responsible for deteriorating glucose tolerance in newly diagnosed, 53-58  
metabolic response to high-protein, low CHO diet in, 243-251  
nonobese, IL-6, adiponectin, leptin and IR in, 258-262  
Japanese NIDDM subjects (continued)  
with retinopathy, relationship between LTA 804C/A and 252A/C polymorphism and TNF- $\alpha$  -308G/A polymorphism, 1406-1410  
Japanese subjects  
correlation between change in BW rather than current BW and change in serum adiponectin levels, 324-330  
male, GNB3 polymorphism C1429T and LRP5 polymorphism A1330V as risk factors for hypercholesterolemia in, 751-757  
NIDDM, *see* Japanese NIDDM subjects  
obese women, influence of AGT polymorphism (Met235Thr) on visceral obesity and IR, 819-824  
PM women, effects of isoflavones and exercise on lipid and bone metabolism in, 423-433
- Kidney  
of Crigler-Najjar type II, results of N-glucuronidation of retigabine from, 711-721  
physiologic roles of 11 $\beta$ -HSD2, 1352-1357  
*see also* Nephropathy and entries beginning with: Renal  
*KLOTHO* polymorphism, with CAD, differential effects of age on the association of, 1344-1351
- Korean subjects  
clinical and biochemical characteristics of nonobese NIDDM patients with GADA, 1107-1112  
with distal pancreatectomy, major role of insulin secretory defect in development of diabetes mellitus, 135-141  
healthy, relation of IR and serum high-sensitivity CRP to FBG subgroups categorized by newly recommended criteria for fasting hyperglycemia in, 183-187  
with NIDDM  
association of Mn-SOD gene polymorphism V16A with diabetic macular edema, 1681-1688  
association of Mn-SOD gene polymorphism (V16A) with stages of albuminuria, 1-7  
nondiabetic, relationship of LDL particle size to IR and IMT, 1610-1615  
nonobese/nondiabetic men, association of SNP276>T versus SNP45T>G adiponectin gene polymorphism with IR, CV disease risk, abdominal adiposity and circulating plasma adiponectin, 59-66  
overweight females, effects of UCP-3 haplotypes on obesity phenotypes and very low-energy diet-induced changes, 578-586  
young and middle-aged, association between CRP and metabolic risk factors in, 415-421
- Lactate, increased concentration in blood, and decreased blood glucose concentration, increased 2DG transport activity accompanied by, effects of AICAR in skeletal muscle, 305
- Lactic acid, plasma concentration, effect of rigorous exercise, 106



- LDL, *see* Low-density lipoprotein(s)
- LDLR, *see* Low-density lipoprotein receptor
- Leptin  
 and adiponectin ratio, relationship with IR parameters, in subjects without hyperglycemia, 1248-1254  
 correlation with hyperhomocysteinemia in obese prepubertal children, 72-77  
 as determinant of 8-epi-prostaglandin F<sub>2α</sub>, 918-922  
 and insulin, effects on fatty acid uptake in differentiated 3T3-L1 adipocytes, 8-12  
 in middle-aged postmenopausal women, in association with lifestyle habits, hormones and inflammatory markers, 1630-1636  
 physiologic concentrations by hepatic stellate cells, effect on collagen production, 1317-1322
- plasma  
 and AT gene expression, effect of aerobic training, in obese women, 1375-1381  
 and glucose and insulin, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154  
 response to maximal sculling after volume-extended training season in elite rowers, 13-19  
 serum, and IL-6, adiponectin and IR in nonobese Japanese NIDDM patients, 258-262  
 short- and long-term, effects on glucose and FA uptake and metabolism, in HL-1 cardiomyocytes, 1067-1075
- Leptin receptor, expression in differentiated and undifferentiated 3T3-L1 adipocytes, 10
- LETO, *see* Long-Evans Tokushima Otsuka
- Leu72Met polymorphism, preproghrelin, association with NIDDM, 366-370
- Lifestyle  
 factors, modification of hormonal situation on lipids by, in middle-aged Swedish women, 1060-1066  
 and glycemic control in NIDDM, S1:7-8  
 modification, effects on metabolic parameters and carotid intima-media thickness, in NIDDM, 1053-1059  
 modifications, associated with weight loss and improved aerobic capacity, effects on CRP, 825-831  
 relation to leptin and adiponectin in middle-aged postmenopausal women, 1630-1636
- Limb muscle adaptations, and S<sub>1</sub>, effects of repeated prolonged whole-body low-intensity exercise, 217-223
- LIPC, *see* Hepatic lipase gene
- Lipid deposition, regional, in adult twins, interrelation with birth weight, 561-562
- Lipid metabolism  
 effect of telmisartan, in nondiabetic insulin-resistant subjects, 1149-1154  
 in PM Japanese women, effects of exercise and isoflavones, 423-433
- Lipid oxidation, net, insulin-mediated suppression of, association with TNF- $\alpha$  in HIV-infected patients with lipodystrophy, 175-182
- Lipid peroxidation  
 as oxidative stress marker in diabetes, S1:11  
 sympathetic sudomotor disturbance in early type IDDM linked to, 1524-1531
- Lipid(s)  
 altered, in IBD, as a consequence on cholesterol efflux, 980-988  
 blood, effects of PSE supplementation and aerobic exercise training, 541-549  
 effect of raloxifene, in postmenopausal women with and without HTG, 972-979  
 influence by lifestyle, modification of hormonal situation on, in middle-aged Swedish women, 1060-1066
- plasma  
 effect of PS mixed with MTGs and HOC oil in overweight men, 391-395  
 and HIV status, 92-93  
 and IL-6 and adiponectin, association with decrease CRP concentration after diet- and physical activity-induced weight reduction, 359-365
- profiles in Ethiopian patients with DM, 706-710
- serum  
 effects of ACTH and equivalent dose of cortisol, 1083-1087  
 and IL-6, adiponectin, leptin and IR in nonobese Japanese NIDDM patients, 258-262  
 and inflammatory marker, glucose metabolism and BP, relations to IMT in childhood obesity, 113-118
- Lipodystrophy  
 HIV-infected patients with, association between TNF- $\alpha$  and insulin-mediated suppression of free fatty acids and net lipid oxidation in, 175-182  
 and MS, in HIV-infected patients treated with ART, 940-945
- Lipolysis  
 epinephrine-induced, increased in IDDM induced by, effect of antecedent hypoglycemia, 371-380  
 of VLDL TGs, differential effects of Apo E isoforms, 1129-1134
- Lipoprotein A-I (LpA-I), effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26-34
- Lipoprotein lipase (LPL), 1481-1487  
 effects of PSE supplementation and aerobic exercise training, 541-549
- Lipoprotein lipase (LPL)–promoting agent, NO-1886 (ibrolipim), effects on expression of UCP3 mRNA and obesity in ovariectomized rats, 151-158
- Lipoprotein(s) (Lp)  
 altered, in IBD, as a consequence on cholesterol efflux, 980-988  
 composition, effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26-34  
 effect of raloxifene, in postmenopausal women with and without HTG, 972-979

## Lipoprotein(s) (Lp) (continued)

- effects of ezetimibe alone, and with colesvelam hydrochloride, in hypercholesterolemic subjects, 1697-1703
  - profiles in Ethiopian patients with DM, 706-710
  - serum, effects of ACTH and equivalent dose of cortisol, 1083-1087
  - subclass and particle size differences in Afro-Caribbeans, African Americans and white Americans, association with LIPC variation, 96-102
- Liver
- long-term exercise on stimulation of AMPK activity and subunit expression in, 1122-1128
  - and microsomal membranes, of Crigler-Najjar type II, results of N-glucuronidation of retigabine from, 711-721
  - mRNA profiles, of female Zucker diabetic fatty rats after topiramate treatment, 1411-1419
  - and serum levels of FFAs, TGs and carnitine in SSADH knockout mice, 355-356
  - and skeletal muscle
    - and fat, and pancreatic islets, gene expression for energy metabolism after ER in Zucker diabetic fatty rats, 43-52
    - oxysterols in, during fasting and alcohol exposure, 119-127
  - and AT and SM, S<sub>i</sub> in extremely obese subjects with acanthosis nigricans, 1658-1663
  - weight of
    - effects of FBCx in high fat-fed rats, 199
    - and glycogen concentration, in LETO and OLETF rats, 319
- LMS, *see* Lysosomal membrane sialidase
- LoBAG diet, *see* Low biologically available glucose diet
- Long-Evans Tokushima Otsuka (LETO), rats, body weight, liver relative weight, liver glycogen concentration and plasma parameters in, 319
- Low biologically available glucose (LoBAG) diet, metabolic response in men with NIDDM, 243-251
- Low-density lipoprotein(s) ( $\beta$ -lipoprotein; LDL)
- atherogenicity, with IGT, effect of acarbose, 946-952
  - effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26-34
  - modification of features, during oral fat load in mild hypertriglyceridemic and healthy subjects, 1308-1316
  - particle size
    - and distribution, effect of weight loss resulting from low-fat diet/exercise on, in obese women, 1302-1307
    - relationship to IR and IMT, in nondiabetic Koreans, 1610-1615
  - subtractions, concentration and relative distribution, in patients with MS, 885-891
  - subtype distribution in patients treated by LDL apheresis and statins, effect of ezetimibe on, 599-604

## Low-density lipoprotein-cholesterol (LDL-C)

- effects of PSE supplementation and aerobic exercise training, 541-549
  - plasma, and CRP, effect of atorvastatin in Japanese NIDDM patients, 67-71
- Low-density lipoprotein receptor (LDLR)
- binding affinity, retinyl palmitate in correlation with, 1308-1316
  - mRNA abundance of, effect of dietary fatty acids, 635-641
- Low-density lipoprotein receptor-related protein 5 (LRP5), A1330V polymorphism in, as a risk factor of hypercholesterolemia in Japanese males, 751-757
- Low-fat diet, and exercise regimen, effect of weight loss on LDL particle size and distribution resulting from, in obese women, 1302-1307
- Low-grade systemic inflammation, and IR, correlation with hyperhomocysteinemia in obese prepubertal children, 72-77
- Lp, *see* Lipoprotein(s)
- LpA-I, *see* Lipoprotein A-I
- LPI, *see* Lysinuric protein intolerance
- LPL, *see* Lipoprotein lipase
- LRP5, *see* Low-density lipoprotein receptor-related protein 5
- Lymphotoxin- $\alpha$  (LTA), 804C/A and 252A/C polymorphism, relationship with TNF- $\alpha$  -308G/A polymorphism, in Japanese subjects with NIDDM retinopathy, 1406-1410
- Lysinuric protein intolerance (LPI), hazards associated with pregnancies and deliveries in, 224-231
- Lysosomal membrane sialidase (LMS), effect of chronic ethanol consumption, in rat liver
- Macrophage migration inhibitory factor (MIF), adipocytes versus non-fat cells in release of, 1113-1121
- Macrophages, derived from monocytes, gliclazide inhibition of differentiation-associated biologic events in, 778-785
- Macular edema, diabetic, association with Mn-SOD gene polymorphism V16A, in Korean NIDDM patients, 1681-1688
- Malnutrition, and acute infection, effects of cortisol on glucocorticoid receptor in, 550-554
- Manganese superoxide dismutase (MnSOD)
- and CAT, CuZnSOD, GCLc, Gpx1 and CypB, mRNA expressions and protein levels in limited amounts of tissue from rat retina, 168-174
  - expression in rat retinal ischemia after reperfusion, 892-898
  - gene polymorphism V16A, association with Mn-SOD, in Korean NIDDM patients, 1-7, 1681-1688
  - oxidative stress with diet-induced MS due to, 928-934
- Maternal body size, and prediction of birth weight, association with and adipokines, 339-344

- Maximum oxygen consumption, *see*  $\text{VO}_{2\text{max}}$  during exercise
- MCP-1, *see* Monocyte chemotactic protein-1
- Medium-chain triglycerides (MTGs), and PS mixed with, and HOC oil, effect on plasma lipids in overweight men, 391-395
- Melatonin, on sleep-related vascular disorders, S2:47-48
- Memory, new evidence of sleep-dependent processing and learning on, S2:5
- Men
- association of hypoadiponectinemia with early onset CHD and multiple coronary stenoses, 1653-1657
  - endurance-trained, impact of dietary proteins on WBPTO at rest in, 501-507
  - Japanese NIDDM, *see* Japanese NIDDM subjects
  - nonobese/nondiabetic Korean, association of SNP276>T versus SNP45T>G adiponectin gene polymorphism with IR, CV disease risk, abdominal adiposity and circulating plasma adiponectin, 59-66
  - obese and overweight, effect of weight-bearing aerobic exercise on bone formation during short-term weight loss in, 1616-1618
  - overweight, effect of PS mixed with MTGs and HOC oil on plasma lipids in, 391-395
- Menstrual cycle, regular, free testosterone, obesity and CVD risk factors in adolescents with PCOS versus with, 508-514
- Messenger ribonucleic acid (mRNA)
- abundance of LDLR, HMG-CoA and MTP in Golden-Syrian hamsters, dietary fats on differential modulation of, 635-641
  - adiponectin receptors AdipoR1 and AdipoR2, in lean rats and obese Zucker rats, 396-401
  - of Apo A-I gene, in HepG2 and Caco-2 cell lines, effect of ascorbic acid and  $\alpha$ -tocopherol on down-regulation of, 159-167
  - for energy metabolism in skeletal muscle, liver, fat and pancreatic islets after ER in Zucker diabetic fatty rats, 43-52
  - of enzymes involved in hepatic peroxisomal fatty acid oxidation, synergistic effect of dietary sesamin, DHA and EPA in rats, 381-390
  - of FA transporter enzymes, effects of NO-1886 in ovariectomized rats, 154
  - of FoF1-ATPase, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154-155
  - and hepatic PDK2 and PDK4 proteins in LETO and OLETF rats, 320-321
  - and HSP, effect of alcohol, 843-851
  - and leptin receptor expression in differentiated and undifferentiated 3T3-L1 adipocytes, 10
- PPAR $\gamma$
- and AT differentiation, effect of troglitazone, 311-312
  - effect of TNF- $\alpha$  and TGF- $\beta$  in freshly isolated adipocytes, 313
- Messenger ribonucleic acid (mRNA) (continued)
- profiles in liver, hypothalamus, AT and SM, of female Zucker diabetic fatty rats after topiramate treatment, 1411-1419
  - and protein levels, of enzymes in limited amounts of tissue from rat retina, 168-174
- TNF- $\alpha$
- in MNCs, effect of glucose ingestion, in healthy subjects, 1177-1185
  - and TGF- $\beta$ , effect of troglitazone, 313
  - of UCP3, and obesity in ovariectomized rats, effects of NO-1886 (ibrolipim), 151-158
  - variant of hepatic lipase, transient induction by corticotropin hormone in rat adrenals, 467-477
- Met, *see* Methionine
- Met235Thr polymorphism, AGT, influence on visceral obesity and IR, in obese Japanese women, 819-824
- Metabolic rate (MR), resting, effect of exercise, in postmenopausal African American and Caucasian women, 1358-1364
- Metabolic response, to high-protein, low CHO diet in men with NIDDM, 243-251
- Metabolic syndrome (insulin resistance syndrome; MS; syndrome X)
- association with Ala54Thr variant of FABP2 gene, in urban South Indians, 1222-1226
  - concentration and relative distribution of LDL subfractions in patients with, 885-891
  - correlation with hyperhomocysteinemia in obese prepubertal children, 72-77
  - diet-induced, oxidative stress and NADPH oxidase dysregulation and antioxidant enzymes in, 928-934
  - distribution in CRP, in biracial younger adults, 699-705
  - elevated 2-hour postload plasma glucose as alternative to elevated fasting glucose as criterion for, 1323-1326
  - elevated serum UA levels in, 1293-1301
  - high sensitivity CRP in association with relative sympathetic overactivity in NIDDM with, 1014-1021
  - IDF definition of, in Chinese patients older than 30 years with NIDDM, 1088-1096
  - and impaired GT and NIDDM, soluble P-selectin and sCD40L levels in subjects with (Chennai Urban Rural Epidemiology Study), 237-242
  - independent association with elevated sCD40L and disease severity, in symptomatic CAD, 1029-1034
  - and IR, association with cardiovascular risk in adults with normal glucose regulation, 445-452
  - link with hyperhomocysteinemia, association between insulin and Hcy leading to, 1007-1013
  - and lipodystrophy, in HIV-infected patients treated with ART, 940-945
  - in overweight children, effects of short-term diet and exercise intervention, 871-878

- Metabolic syndrome (insulin resistance syndrome; MS; syndrome X) (continued)  
 in postmenopausal women, low SHBG associated with, 1473-1480  
 risk factors, association with CRP among young and middle-aged Koreans, 415-421  
 and sleep apnea, S2:48-49  
 in Turkish children and adolescents, 1002-1006
- Metformin (Dimethylbiguanide)  
 effect of adiponectin on artery stiffness in NIDDM treated with, 996-1001  
 effects on isoenergetic feeding in a type 2-diabetic pig model, 960-971  
 plus sulfonylureas, for NIDDM, S1:4  
 in sustainability of 8% weight loss, IR, and amelioration of atherogenic-metabolic risk factors over 4 years in women with PCOS, 1582-1589
- Methionine (Met)  
 concentrations in mid-trimester amniotic fluid, as predictor of body weight and length, 1186-1191  
 different effects choline and folate on plasma kinetics, in rats, 899-906  
 imbalance, on maternal diet, in reduction of postnatal growth in rat, 763-770
- 3-Methylhistidine, effect of liquid CHO/EAA ingestion on myocardial protein degradation during resistance exercise in assessment of, 570-577
- Methylmercury (MeHg), and oxidative stress, in Inuit of Nunavik, 989-995
- MI, *see* Myocardial infarction
- Microsomal triglyceride transfer protein (MTP), mRNA abundance of, effect of dietary fatty acids, 635-641
- Middle-aged women, sample population of, relation of serum adiponectin to glucose tolerance, family history of diabetes, autoimmunity, S<sub>i</sub>, C-peptide and inflammation in, 188-194
- MIF, *see* Macrophage migration inhibitory factor
- Mifepristone, blockade of dexamethasone effects in isolated rat islets, 39
- Milk, containing chromium chloride, influence on glycemic control of NIDDM, 923-927
- Minimal model analysis  
 and C-peptide secretion rate, in determining clinical phenotype of Japanese NIDDM subjects with *Arg121Trp* mutation, 215-216  
 of IVGTT, versus SBT of S<sub>i</sub> and glucose effectiveness, 676-690
- Miscarriage, first-trimester  
 correlates of, 349  
 and PAI activity and 4G5G polymorphism of PAI-1 gene in women with PCOS, 345-352
- Mitochondria, isolated, differential changes in respiratory capacity and ischemia tolerance, from atrophied and hypertrophied hearts, 1097-1106
- MNCs, *see* Mononuclear cells
- MnSOD, *see* Manganese superoxide dismutase
- Monoamine oxidase (MAO), substrates, short- and long-term insulin-like effects, in cultured adipocytes, 1397-1405
- Monocyte chemotactic protein-1 (MCP-1), serum levels of, in relation to IR and glucose intolerance, 1232-1238
- Monocytes, gliclazide inhibition of differentiation-associated biologic events in macrophages derived from, 778-785
- Mononuclear cells (MNCs)  
 altered release of TNF- $\alpha$  from, in obese reproductive-age women during hyperglycemia, 271-276  
 effect of glucose ingestion on NF- $\kappa$ B, I $\kappa$ B and TNF- $\alpha$  mRNA in, 1177-1185
- Mood disorders, and circadian rhythm, S2:10-11
- MR, *see* Metabolic rate
- mRNA, *see* Messenger ribonucleic acid
- MS, *see* Metabolic syndrome
- mTOR inhibitor, RAPA  
 disruption of TG metabolism, in guinea pigs, 794-802  
 in SM cells, effect on glucose utilization and fatty acid oxidation, 1637-1644
- Muscular performance, and hormonal responses during resistance training overreaching, effects of AA supplementation, 282-291
- Mutation(s), *PAX4 Arg121Trp*, association with beta-cell dysfunction in Japanese NIDDM subjects, 213-216
- Myocardial infarction(s) (MI), risk of, and sulfonylureas, S1:16-19
- Myocardial ischemia, sulfonylureas and, S1:16
- Myofibrillar proteins, degradation, effect of liquid CHO/EAA ingestion during short-term resistance exercise, 570-577
- Myostatins, suppression, Gln in prevention of GC-induced SM atrophy in association with, 1239-1247
- n-3 PUFAs, *see*  $\omega$ -3 Polyunsaturated fatty acids
- NADPH, *see* Nicotinamide adenine dinucleotide phosphate
- NAFLD, *see* Nonalcoholic fatty liver disease
- Narcolepsy  
 genetics of, S2:9  
 and hypocretins, S2:36-39
- National Cholesterol Education Program (NCEP),  
 concentration and relative distribution of LDL subfractions in patients with MS, 885-891
- NCEP, *see* National Cholesterol Education Program
- NEFAs, *see* Nonesterified fatty acids
- Nephropathy, non-insulin dependent diabetes mellitus  
 early stage, effects of EPA, 1590-1598  
 effect of spironolactone on circulating adipocytokines in, 1645-1652
- Nerve growth factor (NGF), adipocytes versus non-fat cells in release of, 1113-1121
- Neurobiology, of sleep, S2:2-6
- Neurochemicals, and wake centers, S2:15-17



- Neurocognitive functions, and symptoms, in response to brief hypoglycemia in PM women and age-matched men with NIDDM, 336
- Neurocognitive responses, to brief hypoglycemia in PM women and age-matched men with NIDDM, 331–338
- Neurophysiology, of sleep, essential brain regions in, S2:4–5
- Neutrophils, from pregnant women, apparent role of dynein in G6PDase trafficking in, 279–281
- NF- $\kappa$ B, *see* Nuclear factor  $\kappa$ B
- NGF, *see* Nerve growth factor
- N-Glucuronidation, of retigabine, results from human UGTs, liver and kidney of Crigler-Najjar type II, 711–721
- Niacin, adipokines and treatment with, 1283–1285
- Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase  
dysregulation, 928–934  
reduced form, activation by AGEs, linked oxidative stress to altered retinal VEGF, 1516–1523
- NIDDM, *see* Non-insulin dependent diabetes mellitus
- Nitration, and protein oxidation, as oxidative stress marker in diabetes, S1:11
- Nitric oxide synthase (NOS), endogenous, ADMA in citrullinemia, elevated plasma concentrations, 1599–1603
- NO-1886, ibrolipim, effects on expression of UCP3 mRNA and obesity in ovariectomized rats, 151–158
- Nocturnal enuresis, primary, S2:9–10
- Nocturnal terrors, and sleepwalking, S2:9
- Nonalcoholic fatty liver disease (NAFLD), association with increased ALT, 1604–1609
- Non-diabetic subjects, nonobese Korean men, association of SNP276>T versus SNP45T>G adiponectin gene polymorphism with IR, CV disease risk, abdominal adiposity and circulating plasma adiponectin, 59–66
- Nonesterified fatty acids (free fatty acids; NEFAs)  
effect on adrenocorticotropin and cortisol, in healthy young men, 1022–1028  
metabolism in NIDDM, effect of high-dose simvastatin, 758–762  
and net lipid oxidation, insulin-mediated suppression of, association with TNF- $\alpha$  in HIV-infected patients with lipodystrophy, 175–182  
role in increased expression of hepatic PDKs in young and middle-aged OLETF rats, 317–323  
and TGs and carnitine, levels in liver and serum of SSADH knockout mice, 355–356  
turnover, effect of fasting and SM LPL overexpression, 1481–1487
- Non-insulin dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus)  
association with Mn-SOD gene polymorphism V16A with diabetic macular edema in Korean patients with, 1–7, 1681–1688
- Non-insulin dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) (continued)  
association with preproghrelin Leu72Met polymorphism, 366–370  
BDNF association with obesity in newly diagnosed female with, 852–857  
correlation of PON1 and HCTL and antioxidant capacity of HDL with severity of CV disease with, 1201–1206  
and CV risk factors, relation to IMCL in healthy men versus women subjects, 128–134  
differential effects of n-3 PUFAs on metabolic control and vascular reactivity in mouse with, 1365–1374  
effect of a pedometer-based physical activity program on control of, 1382–1387  
effect of exercise training on endothelial function and biochemical vascular markers in first-degree relatives of, 1508–1515  
effect of high-dose simvastatin on NEFA metabolism in, 758–762  
effect of insulin on intracellular oxidative stress in patients with, 691–695  
effects of ER on development of, coordinated patterns of gene expression for energy metabolism determined by oligonucleotide microarray, 43–52  
effects of lifestyle modification on metabolic parameters and carotid intima-media thickness in, 1053–1059  
estrogen treatment on arterial distensibility, fibrinolysis and metabolic profile, in postmenopausal women with, 953–959  
glycemic control in, *see* Glycemic control of NIDDM  
hormonal, subjective and neurocognitive responses to brief hypoglycemia in PM women and age-matched men with, 331–338  
IDF definition of MS in Chinese subjects older than 30 years with, 1088–1096  
and impaired GT and MS, soluble P-selectin and sCD40L levels in subjects with (Chennai Urban Rural Epidemiology Study), 237–242  
insulin absorption as a major factor in control of, 614–619  
insulin therapy in reduction of small dense LDL particles in patients with, 879–884  
in Japanese subjects, *see* Japanese NIDDM subjects  
metabolic response to high-protein, low CHO diet in men with, 243–251  
modestly advanced, multiple defects in counterregulation of hypoglycemia, 593–598  
with MS, high sensitivity CRP in association with relative sympathetic overactivity in, 1014–1021  
nonobese patients, with GADA, clinical and biochemical characteristics of, 1107–1112  
with obesity, *see* Obesity, NIDDM with  
oxidizing and reducing responses of granulocytes from, 1426–1428  
pathogenesis of, in overcoming hurdles for achieving glycemic control, S1:6–7  
pathophysiologic basis for treatment, S1:21

- Non–insulin dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) (continued)  
 percentage of AT and plasma glucose in prediction of sialic acid concentration in, 1165–1170  
 place of sulfonylureas in, S1:20–26  
 postprandial TAFI and markers of endothelial dysfunction in, 1437–1442  
 relationship between serum resistin and inflammatory markers in patients with, 1670–1673  
 with renal impairment, Cockcroft–Gault formula is biased by body weight in, 108–112  
 role of autonomic imbalance of HPA activity in, 1135–1140  
 sulfonylureas for, S1:1  
 treated with pioglitazone and metformin, effect of adiponectin on artery stiffness, 996–1001
- Non–insulin dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus)) nephropathy  
 early stage, effects of EPA, 1590–1598  
 effect of spironolactone on circulating adipocytokines in, 1645–1652
- Non–insulin dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) retinopathy  
 association with increased serum levels of methylglyoxal-derived hydroimidazolone, 232–236  
 relationship between LTA 804C/A and 252A/C polymorphism and TNF- $\alpha$  –308G/A polymorphism in Japanese subjects with, 1406–1410
- Non-obese (lean) subjects  
 with impaired glucose tolerance, effects of rosiglitazone on serum adiponectin and CRP in, 439–444  
 Japanese with NIDDM, IL-6, adiponectin, leptin and IR in, 258–262  
 nondiabetic Korean men, association of SNP276>T versus SNP45T>G adiponectin gene polymorphism with IR, CV disease risk, abdominal adiposity and circulating plasma adiponectin, 59–66  
 rats, expression of adiponectin receptors AdipoR1 and AdipoR2 in, 396–401
- Noradrenaline, plasma concentration, effect of rigorous exercise, 106
- Normal glucose tolerance, relationship between serum resistin and proinflammatory cytokines in obese women with, 1495–1499
- Nuclear factor  $\kappa$ B (NF- $\kappa$ B), intranuclear, in MNCs, effect of glucose ingestion, in healthy subjects, 1177–1185
- Obese subjects  
 Zucker rats, expression of adiponectin receptors AdipoR1 and AdipoR2 in, 396–401
- Obesity  
 childhood, relations of IMT to inflammatory marker, glucose metabolism and BP in, 113–118  
 correlation of hyperhomocysteinemia with IR and low-grade systemic inflammation in prepubertal children, 72–77
- Obesity (continued)  
 and expression of UCP3 mRNA, in ovariectomized rats, effects of NO-1886 (ibrolipim), 151–158  
 extreme, with acanthosis nigricans, AT, liver and SM S<sub>i</sub> in, 1658–1663  
 and family history of diabetes, glucose tolerance, autoimmunity, S<sub>i</sub>, C-peptide and inflammation, in relation to serum adiponectin in population sample of 64-year-old women, 188–194  
 and free testosterone and CVD risk factors in adolescents with PCOS versus regular cycling adolescents, 508–514  
 levels of adiponectin in American and Japanese men with, 1561–1563  
 moderate, effects of  $\beta$ -adrenergic agonists combined with PPAR  $\gamma$  on Lp composition in, 26–64  
 in newly diagnosed female with NIDDM, BDNF associated with, 852–857  
 NIDDM with, IR and low sympathetic nerve activity in Tzumura Suzuki obese diabetic mouse as a new model for, 1664–4669  
 and normal-weight, and overweight, BSA in adults with, 515–524  
 or sleep apnea, IL-6 elevation in PCOS independent of, 1076–1082  
 phenotypes, in overweight Korean female subjects, effects of UCP-3 haplotypes, 578–586  
 premenopausal women, VAT as determinant of vWF in, 650–655  
 protective effect of BDNF on pancreatic islets in diabetic mice with, 1286–1292  
 reduced by, and lean subjects, effects of short-term overfeeding on insulin action, 1207–1214  
 reproductive-age women with, altered release of TNF- $\alpha$  from MNCs during hyperglycemia in, 271–276  
 voglibose in reduction of oxidative stress markers and soluble ICAM-1 in NIDDM patients with, 786–793
- Offspring  
 adult, impact of maternal folic acid supplementation, 628–634  
 body composition, and insulin axis, programming of maternal protein intake on pregnant rat, 642–649  
 of parents with HTN, metabolic characteristics of soleus muscle in relation to insulin action in, 1388–1396
- OGTT, *see* Oral glucose tolerance test
- Older subjects  
 Chinese, association of vascular risk factors with increasing glycemia and normoglycemic subjects, 1035–1041  
 hospitalized, clinical value of Hcy in, 620–627  
 with impaired glucose tolerance, S<sub>i</sub> assessment indices versus euglycemic-hyperinsulinemic clamp data after diet and exercise intervention in, 525–532
- Oleic acid, and palmitic acid, dietary intake of, differential effects on oxygen consumption, during and after exercise, 1215–1221

- OLETF, *see* Otsuka Long-Evans Tokushima Fatty
- Oligonucleotide microarray, ER prevents development of  
NIDDM, coordinated patterns of gene expression  
for energy metabolism determined by, 43-52
- Olmesartan, plasma adiponectin levels in rats infused with  
angiotensin II with or without, 480-483
- Oral glucose tolerance test (OGTT), IR in hypertension  
quantified by, comparison of methods, 143-150
- Orlistat, short-term effects on dietary glycotoxins in  
healthy women and women with PCOS, 494-500
- Osteoporosis  
effects of alendronate combined with HRT on  
postmenopausal Chinese women with, 741-747  
effects of raloxifene on lipid and Lp in postmenopausal  
women with, 972-979
- Otsuka Long-Evans Tokushima Fatty (OLETF), young and  
middle-aged rats  
body weight, liver relative weight, liver glycogen  
concentration and plasma parameters in, 319  
role of FFAs in increased expression of hepatic PDKs in,  
317-323
- Overweight  
effect of weight loss on  $S_i$  and arterial stiffness, 907-911  
effects of short-term diet and exercise intervention on  
MS in children, 871-878  
in Korean female subjects, effects of UCP-3 haplotypes  
on obesity phenotypes and very low-energy diet–  
induced changes in, 578-586  
men, effect of PS mixed with MTGs and HOC oil on  
plasma lipids in, 391-395  
and normal-weight, and obesity, BSA in adults with,  
515-524  
and obese men and women, effect of weight-bearing  
aerobic exercise on bone formation during  
short-term weight loss in, 1616-1618  
premenopausal women, VAT as determinant of vWF in,  
650-655
- Oxidation, fatty acid, and glucose utilization, effect of  
RAPA in SM cells, 1637-1644
- Oxidative stress  
beta-cell damage induced by, protective role of gliclazide  
on, S1:13  
and dietary contaminants, in Inuit of Nunavik, 989-995  
intracellular, effect on insulin on, in patient with NIDDM,  
691-695  
linked to altered retinal VEGF, activation of NADPH  
(reduced form) by AGEs associated with, 1516-1523  
markers, in diabetes, S1:11-12  
8-hydroxy-2'-deoxyguanosine, S1:12  
DNA damage, S1:12  
lipid peroxidation, S1:11  
protein oxidation and nitration, S1:11  
markers in obese NIDDM patients, voglibose in reduction  
of, 786-793  
and NADPH oxidase dysregulation and antioxidant  
enzymes, in diet-induced MS, 928-934  
in nondiabetic hypercholesterolemic patients, 918-922
- Oxygen consumption, differential effects of dietary intake  
of palmitic and oleic acid, during and after exercise,  
1215-1221
- Oxypurinol, and purine bases, plasma concentrations  
and urinary excretion after rigorous exercise,  
103-107
- Oxysterols, in liver and skeletal muscle during fasting and  
alcohol exposure, 119-127
- PAI, *see* Plasminogen activator inhibitor
- PAI-1, *see* Plasminogen activator inhibitor-1
- Palmitic acid, and oleic acid, dietary intake of, differential  
effects on oxygen consumption, during and after  
exercise, 1215-1221
- Pancreas, increased flux to n-6 docosapentaenoic acid in  
PLs of, from  $cfr^{-/-}$  knockout mice, 1192-1200
- Pancreatectomy, distal, major role of insulin secretory  
defect in development of diabetes mellitus in  
patients with, 135-141
- Pancreatic atrophy, effects of acarbose on, 533-540
- Pancreatic cells, *see* Beta ( $\beta$ ) cell(s) (islets of Langerhans)
- Parathyroid hormone (PTH), spontaneous, effect of active  
acromegaly on pulsatility of, 736-740
- Parent(s), with HTN, metabolic characteristics of soleus  
muscle in relation to insulin action in offspring of,  
1388-1396
- Pathogenesis of NIDDM, in overcoming hurdles for  
achieving glycemic control, S1:6-7
- Pathophysiology, of sleep-related vascular disorders,  
S2:47-49
- PAX4* gene, *Arg121Trp* mutation in  
association with beta-cell dysfunction in Japanese  
NIDDM subjects, 213-216  
determination of, 215
- PCOS, *see* Polycystic ovary syndrome
- PDC, *see* Pyruvate dehydrogenase complex
- PDK2, *see* Pyruvate dehydrogenase kinase 2
- PDK4, *see* Pyruvate dehydrogenase kinase 4
- Peroxisomal fatty acid oxidation, hepatic, synergistic effect  
of dietary sesamin, DHA and EPA on gene  
expression of enzymes involved in, 381-390
- Peroxisome proliferator-activated receptor  $\gamma$  (PPAR  $\gamma$ )  
combined with  $\beta$ -adrenergic agonists, effects on Lp  
composition in healthy moderately obese subjects,  
26-34  
expression, effect of TNF- $\alpha$  and TGF- $\beta$  in freshly  
isolated adipocytes, 313  
mRNA expression, and AT differentiation, effect of  
troglitazone, 311-312
- Pharmacology, of wakefulness, S2:13-19
- Phosphatidylinositol 3-kinase (PI3-K), insulin stimulated  
IRS-2 associated with, in SM after exercise,  
1046-1052
- Phospholipase C (PLC), activation, and insulin secretion,  
effect of dexamethasone in isolated rat islets,  
35-42

- Phospholipid(s) (PLs), of pancreas from cfr <sup>-/-</sup> knockout mice, increased flux to n-6 docosapentaenoic acid in, 1192-1200
- Physical activity  
and diet, weight reduction induced by, decrease CRP concentration after, association with changes in plasma lipids, IL-6 and adiponectin, 359-365  
effect of a pedometer-based program, in control of NIDDM, 1382-1387
- Phytosterols (PS), mixed with MTGs, and HOC oil, effect on plasma lipids in overweight men, 391-395
- PI3-K, *see* Phosphatidylinositol 3-kinase
- Pioglitazone  
effect of adiponectin on artery stiffness in NIDDM treated with, 996-1001  
TNF- $\alpha$ -induced production of PAI-1 in regulation of, in nonmalignant hepatocyte cell line, 1464-1472
- PKC, *see* Protein kinase C
- Plant stanol ester (PSE) supplementation, and aerobic exercise training, effects on blood lipid levels, 541-549
- Plant sterols, effects of stigmasterol on intestinal absorption of, 292-299
- Plasma  $\alpha$ -amino nitrogen, in response to LoBAG diet in men with NIDDM, 247
- Plasma adiponectin  
association with carotid arterial stiffness, 587-592  
decreased by angiotensin II infusion, 478-488  
and AT gene expression, effect of aerobic training, in obese women, 1375-1381  
and plasma lipids and IL-6, association with decrease CRP concentration after diet- and physical activity-induced weight reduction, 359-365
- Plasma cholesterol (C), levels, and hepatic synthesis and intestinal absorption, effects of stigmasterol in rat, 292-299
- Plasma corticosterone, impact of maternal folic acid supplementation, in adult offsprings, 628-634
- Plasma C-peptide, effects of glimepiride and glyburide, 80-81
- Plasma glucose  
elevated 2-hour postload, as alternative to elevated fasting glucose, criterion for MS, 1323-1326  
fasting, utility in detection of prediabetes in African Americans, 434-438  
and insulin  
effects of glimepiride and glyburide, 79-80  
and leptin, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154  
and percentage of AT, prediction of sialic acid concentration in NIDDM, 1165-1170
- Plasma homocysteine (Hcy), relation to plasma SAH concentration and GFR in children, 252-257
- Plasma insulin  
and glucose  
effects of glimepiride and glyburide, 79-80
- Plasma insulin (continued)  
and leptin, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154  
relation to adiponectin in healthy children, 1171-1176
- Plasma interleukin-6 (IL-6)  
elevation in PCOS independent of obesity and sleep apnea, 1076-1082  
and plasma lipids and adiponectin, association with decrease CRP concentration after diet- and physical activity-induced weight reduction, 359-365
- Plasma leptin, and glucose and insulin, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154
- Plasma lipids  
effect of PS mixed with MTGs and HOC oil in overweight men, 391-395  
and HIV status, 92-93  
and IL-6 and adiponectin, association with decrease CRP concentration after diet- and physical activity-induced weight reduction, 359-365
- Plasma low-density lipoprotein-cholesterol (LDL-C), and CRP, effect of atorvastatin in Japanese NIDDM patients, 67-71
- Plasma membrane sialidase (PMS), effect of chronic ethanol consumption, in rat liver, 803-810
- Plasma parameters, in LETO and OLETF rats, 319
- Plasma renin, and urinary aldosterone and urinary free cortisol, in response to LoBAG diet in men with NIDDM, 248
- Plasma S-adenosylhomocysteine (SAH), relation to GFR in children, 252-257
- Plasma urea nitrogen, in response to LoBAG diet in men with NIDDM, 247
- Plasminogen activator inhibitor (PAI), activity, and 4G5G polymorphism of PAI-1 gene and first-trimester miscarriage in women with PCOS, 345-352
- Plasminogen activator inhibitor-1 (PAI-1)  
changes in, on glucophage diet, relation to first-trimester miscarriage in women with PCOS, 349  
gene, 4G5G polymorphism, and PAI activity and first-trimester miscarriage in women with PCOS, 345-352  
genotypes, 349  
induced by TNF- $\alpha$  in regulation of pioglitazone and cerivastatin in nonmalignant hepatocyte cell line, 1464-1472
- PLC, *see* Phospholipase C
- PL(s), *see* Phospholipids
- PMN cells, *see* Polymorphonuclear cells
- PMS, *see* Plasma membrane sialidase
- PM women, *see* Postmenopausal (PM) women
- Polychlorinated biphenyls (PCBs), and oxidative stress, in Inuit of Nunavik, 989-995
- Polycystic ovary syndrome (PCOS)  
IL-6 elevation in, independent of obesity or sleep apnea, 1076-1082  
PAI activity, 4G5G polymorphism of PAI-1 gene and first-trimester miscarriage in women with, 345-352



- Polycystic ovary syndrome (PCOS) (continued)  
 versus regular cycling, free testosterone, obesity and CVD risk factors in adolescents with, 508-514  
 short-term effects of orlistat on dietary glycotoxins in health women and women with, 494-500  
 sustainability of 8% weight loss, IR, and amelioration of atherogenic-metabolic risk factors over 4 years by metformin-diet in women with, 1582-1589  
 universe of cases with, 348-349
- Polymorphism  
 4G5G, of PAI-1 gene, and PAI activity and first-trimester miscarriage in women with PCOS, 345-352  
 in LIPC, association with Lp subclass and particle size differences in Afro-Caribbeans, African Americans and white Americans, 96-102  
 Mn-SOD (V16A) gene, association with stages of albuminuria in Korean NIDDM patients, 1-7  
 SNP276>T versus SNP45T>G, in adiponectin gene, association with IR, CV disease risk, abdominal adiposity and circulating plasma adiponectin in nonobese/nondiabetic Korean men, 59-66
- Polymorphonuclear (PMN) cells, effect of hyperglycemia and/or hyperinsulinemia on functions of, in healthy subjects, 811-818
- $\omega$ -3 Polyunsaturated fatty acids (n-3 PUFAs)  
 differential effects on metabolic control and vascular reactivity, in NIDDM mouse, 1365-1374  
 protective effect in Inuit of Nunavik, 989-995
- PON1, *see* Serum paraoxonase
- Postmenopausal (PM) women  
 African American and Caucasian, effect of exercise on resting MR, 1358-1364  
 and age-matched men, with NIDDM, hormonal, subjective and neurocognitive responses to brief hypoglycemia in, 331-338  
 Japanese, effects of isoflavones and exercise on lipid and bone metabolism in, 423-433  
 low SHBG associated with MS in, 1473-1480  
 middle-aged, lifestyle, hormones and inflammatory markers associated with leptin and adiponectin in, 1630-1636  
 with NIDDM, estrogen treatment on arterial distensibility, fibrinolysis and metabolic profile in, 953-959  
 osteoporotic, effects of alendronate combined with HRT, 741-747  
 with and without HTG, effects of raloxifene on lipid and Lp, 972-979
- Postprandial glucose (PPG)  
 control, potential impact, 1266  
 pregnancy as prototype for targeting, 1270-1271  
 priority, 1271  
 treatment approaches, 1267-1270
- Potassium, insulin release induced by, effects of dexamethasone, 39
- PPAR $\gamma$ , *see* Peroxisome proliferator-activated receptor  $\gamma$
- PPG, *see* Postprandial glucose
- Prediabetes, and NIDDM and MS, soluble P-selectin and sCD40L levels in subjects with (Chennai Urban Rural Epidemiology Study), 237-242
- Pregnancy  
 and aging, effect on glucose tolerance in rats, 409-414  
 apparent role of dynein in G6PDase trafficking in neutrophils during, 279-281  
 and deliveries, in LPI, hazards associated with, 224-231  
 normal, IRS-1 and  $\beta_3$ -ADR genes in, 1451-1456  
 as prototype for targeting PPG, 1270-1271  
 rat, maternal protein intake programming of insulin axis and offspring body composition in, 642-649
- Premenopausal women, overweight and obese, VAT as determinant of vWF in, 650-655
- Preproghrelin Leu72Met polymorphism, association with NIDDM, 366-370
- Prepubertal children, *see* Children
- Protein glycoxidation, effect of valsartan, therapy, 1619-1624
- Protein kinase C (PKC)  
 contents of dexamethasone-pretreated islets, 39-41
- Protein oxidation, and nitration, as oxidative stress marker in diabetes, S1:11
- Proteins  
 dietary, impact on WBPTO at rest in endurance-trained men, 501-507  
 levels, and mRNA expressions, of enzymes in limited amounts of tissue from rat retina, 168-174
- Protein turnover, whole body, at rest in endurance-trained men, impact of dietary proteins, 501-507
- PSE supplementation, *see* Plant stanol ester (PSE) supplementation
- Psychiatry, sleep disorders in, S2:40-44
- PTH, *see* Parathyroid hormone
- Public health, consequence of sleep disorders, S2:41-42
- Purine bases, and oxypurinol, plasma concentrations and urinary excretion after rigorous exercise, 103-107
- Pyruvate dehydrogenase complex (PDC), hepatic, activities in LETO and OLETF rats, 319-320
- Pyruvate dehydrogenase kinase 2 (PDK2), hepatic amounts and mRNAs in LETO and OLETF rats, 320-321  
 increased expression in young and middle-aged OLETF rats, induction by elevated levels of FFA, 317-323
- Pyruvate dehydrogenase kinase 4 (PDK4), hepatic amounts and mRNAs in LETO and OLETF rats, 320-321  
 increased expression in young and middle-aged OLETF rats, induction by elevated levels of FFA, 317-323
- Race and ethnicity, 1222-1226  
 African Americans  
 and Afro-Caribbeans, and white Americans, association of Lp subclass and particle size differences with LIPC variation, 96-102  
 and Caucasians, effect of exercise on resting MR in postmenopausal women, 1358-1364

- Race and ethnicity (continued)  
utility of fasting plasma glucose in detection of prediabetes in, 434-438
- Asian Indians  
with impaired GT, NIDDM and MS, soluble P-selectin and sCD40L levels in (Chennai Urban Rural Epidemiology Study), 237-242  
simple anthropometric measures in fasting hyperinsulinemia and clustering of CV risk factors in adolescents, 1569-1573  
black and white younger adults, distribution of MS variables in correlation with different CRP levels in, 699-705  
*see also* Chinese subjects; Ethiopian subjects; Japanese subjects
- Raloxifene, effects on lipid and Lp in postmenopausal women with and without HTG, 972-979
- RAPA, *see* Rapamycin
- Rapamycin (RAPA)  
disruption of TG metabolism, in guinea pigs, 794-802  
in SM cells, effect on glucose utilization and fatty acid oxidation, 1637-1644
- Rapid eye movement (REM) sleep, and basic cycle, S2:3
- Receptor for advanced glycation end products (RAGEs), association with AGEs, in nondiabetic subjects, 1227-1231
- REE, *see* Resting energy expenditure
- Renal disease, severity, relation to plasma Hcy and SAH in children, 252-257
- Renal impairment, Cockcroft-Gault formula is biased by body weight in diabetic patients with, 108-112
- Renal insufficiency, relation to plasma Hcy and SAH in children, 252-257
- Renin, plasma, and urinary aldosterone and urinary free cortisol, in response to LoBAG diet in men with NIDDM, 248
- Reproductive-age women, altered release of TNF- $\alpha$  from MNCs during hyperglycemia in, 271-276
- Resistance exercise, short-term, effect of liquid CHO/EAA ingestion on myocardial protein degradation during, 570-577
- Resistance training, overreaching, effects of AA supplementation on hormonal responses to, 282-291
- Respiratory quotient (RQ), effects of NO-1886 (ibrolipim) in ovariectomized rats, 154
- Resting energy expenditure (REE), persistent suppression of, after acute hypoxia, 669-675
- Restless legs syndrome (RLS), and periodic limb movement, S2:10
- Retigabine, N-Glucuronidation of, results from human UGTs, liver and kidney of Crigler-Najjar type II, 711-721
- Retina, mRNA expressions and protein levels of enzymes in limited amounts of tissue from, 168-174
- Retinopathy, non-insulin dependent diabetes mellitus  
association with increased serum levels of methylglyoxal-derived hydroimidazolone, 232-236
- Retinopathy, non-insulin dependent diabetes mellitus (continued)  
relationship between LTA 804C/A and 252A/C polymorphism and TNF- $\alpha$  -308G/A polymorphism in Japanese subjects with, 1406-1410
- Retinyl palmitate, correlation with modification of LDL features and LDLR binding affinity, in mild hypertriglyceridemic and healthy subjects, 1308-1316
- Reverse transcription-polymerase chain reaction (RT-PCR), and Western blot, for detection of mRNA expressions and protein levels of enzymes in limited amounts of tissue from rat retina, 168-174
- Risk, for CV disease  
in adults with normal glucose regulation, association with MS and IR, 445-452  
and free testosterone and obesity in adolescents with PCOS versus regular cycling adolescents, 508-514  
and IR, and abdominal adiposity circulating plasma adiponectin, association with SNP276>T versus SNP45T>G adiponectin gene polymorphism in nonobese/nondiabetic Korean men, 59-66  
markers of, impact of rosiglitazone combined with glimepiride, 20-25  
and NIDDM risk factors, relation to IMCL in healthy men versus women subjects, 128-134
- Risk, for MI, and sulfonylureas, S1:16-19
- Risk factors  
for hypocholesterolemia in Japanese males, GNB3 polymorphism C1429T and LRP5 polymorphism A1330V, 751-757  
metabolic, association with CRP among young and middle-aged Koreans, 415-421  
nonobese metabolic, CT scans of intra-abdominal fat, anthropometric measurements and, 1337-1343  
and prevalence of androgen deficiency, in middle-aged men on Hong Kong, 1488-1494
- Risk factors, for CV disease  
adiponectin and its correlates with, in young adults, 1551-1557  
association with hypoadiponectinemia, in nonobese healthy adults, 1546-1550  
association with increasing glycemia, in older Chinese population, 1035-1041  
impact of sex-specific body composition, 563-569  
simple anthropometric measures, in Asian Indian adolescents, 1569-1573
- Rosiglitazone  
effects on serum adiponectin and CRP in nonobese subjects with impaired glucose tolerance, 439-444  
with glimepiride, impact on beta-cell function, IR, and adiponectin concentrations, 20-25
- RQ, *see* Respiratory quotient
- RT-PCR, *see* Reverse transcription-polymerase chain reaction

- S*-Adenosylhomocysteine (SAH)  
 plasma, *see* Plasma *S*-Adenosylhomocysteine  
 SAH, *see* *S*-Adenosylhomocysteine  
 SBT, *see* Standardized breakfast test  
 sCD40L, *see* Soluble CD40L  
 SCI, *see* Silent cerebral infarction  
 Selenium (Se), protective effect, in Inuit of Nunavik, 989-995  
 Semicarbazide-sensitive amine oxidase (SSAO),  
 substrates, short- and long-term insulin-like  
 effects, in cultured adipocytes, 1397-1405  
 Serotonin, neurochemicals on awake state, S2:16  
 Serum adiponectin  
 change in levels of, correlation with change in BW rather  
 than current BW in Japanese population, 324-330  
 and CRP, in nonobese subjects with impaired glucose  
 tolerance, effects of rosiglitazone, 439-444  
 and IL-6, leptin and IR in nonobese Japanese NIDDM  
 patients, 258-262  
 in population sample of 64-year-old women, relation to  
 glucose tolerance, family history of diabetes,  
 autoimmunity, S<sub>i</sub>, C-peptide and inflammation,  
 188-194  
 Serum free levorotatory thyroxine, in response to LoBAG  
 diet in men with NIDDM, 248  
 Serum growth hormone (GH), and serum IGF-1, in  
 response to LoBAG diet in men with NIDDM,  
 247-248  
 Serum high-sensitivity C-reactive protein (CRP), versus  
 IR, relation to FBG subgroups categorized by newly  
 recommended criteria for fasting hyperglycemia in  
 10 059 healthy Koreans, 183-187  
 Serum insulin  
 association with Hcy, in Swedish population, 1007-1013  
 and blood glucose and C-peptide, in PM women and  
 age-matched men with NIDDM, 334  
 Serum insulin-like growth factor-1 (IGF-1), and serum  
 GH, in response to LoBAG diet in men with  
 NIDDM, 247-248  
 Serum interleukin-6 (IL-6), and adiponectin, leptin and IR  
 in nonobese Japanese NIDDM patients, 258-262  
 Serum leptin, and IL-6, adiponectin and IR in nonobese  
 Japanese NIDDM patients, 258-262  
 Serum lipids  
 and IL-6, adiponectin, leptin and IR in nonobese  
 Japanese NIDDM patients, 258-262  
 and inflammatory marker, glucose metabolism and BP,  
 relations to IMT in childhood obesity, 113-118  
 Serum paraoxonase (PON1), correlation with severity of  
 CV disease with NIDDM, 1201-1206  
 Serum resistin  
 relationship with inflammatory markers, in NIDDM  
 patients, 1670-1673  
 relationship with proinflammatory cytokines, in obese  
 women with IGT and normal glucose tolerance,  
 1495-1499  
 Serum thyroid-stimulating hormone (TSH), in response to  
 LoBAG diet in men with NIDDM, 248  
 Serum total triiodothyronine, in response to LoBAG diet in  
 men with NIDDM, 248  
 Sesamin, dietary, and DHA and EPA, synergistic effect on  
 gene expression of enzymes involved in hepatic  
 peroxisomal fatty acid oxidation in rats, 381-390  
 Sex  
 differences in relation between CV/NIDDM risk factors  
 and IMCL in healthy subjects, 128-134  
 in moderation of long-term alcohol exposure on HSP  
 gene expression in rats, 843-851  
 Sex hormone-binding globulin (SHBG), low, associated  
 with MS, in postmenopausal women, 1473-1480  
 Sex hormones, in response to brief hypoglycemia in PM  
 women and age-matched men with NIDDM, 336  
 SHBG, *see* Sex hormone-binding globulin  
 S<sub>i</sub>, *see* Insulin sensitivity  
 Sialic acid, plasma concentration, percentage of AT and  
 plasma glucose in prediction of, in NIDDM,  
 1165-1170  
 Silent cerebral infarction (SCI), progression in patients with  
 type 2 diabetes mellitus, relation to CRP and soluble  
 adhesion molecules, 461-466  
 Simvastatin, high-dose, effect on NEFA metabolism, in  
 NIDDM, 758-762  
 Skeletal muscle (SM)  
 atrophy induced by GC, Gln in prevention of, 1239-1247  
 cells, effect of RAPA on glucose utilization and fatty acid  
 oxidation, 1637-1644  
 effects of activation of  $\alpha 2$  AMPK by AICAR at  
 physiological levels on glucose transport and  
 GLUT4 in, 300-308  
 effects of HF feeding on components of CAP/Cb1  
 signaling cascade in Sprague-Dawley rat, 203-212  
 and HSP, effect of alcohol, 843-851  
 human, after exercise, insulin stimulated IRS-2  
 associated PI3-K in, 1046-1052  
 and liver  
 and fat, pancreatic islets, gene expression for energy  
 metabolism after ER in Zucker diabetic fatty rats,  
 43-52  
 oxysterols in, during fasting and alcohol exposure,  
 119-127  
 LPL overexpression, effect on NEFA turnover in mice,  
 1481-1487  
 mRNA profiles in, of female Zucker diabetic fatty rats  
 after topiramate treatment, 1411-1419  
 and S<sub>i</sub>, AT and liver, in extremely obese subjects with  
 acanthosis nigricans, 1658-1663  
 and WAT, weights of, effects of NO-1886 (ibrolipim) in  
 ovariectomized rats, 154  
 Sleep  
 basic cycle and REM, S2:3  
 and energy balance, S2:30-35  
 function of, S2:5

- Sleep (continued)  
and health, S2:41-42  
historical perspectives, S2:3  
interdependence of health and, commentary, S2:50-53  
ischemic mechanism in control of, S2:26-27  
messengers of, and joint action on metabolism and feeding, S2:27-28  
neurobiology of, S2:2-6  
neurophysiology of, S2:4-5  
normal, S2:2-3  
relationship with feeding, S2:25-26  
and vascular disorders, S2:45-49  
and wakefulness, metabolic mechanism, S2:24-29
- Sleep apnea  
and MS, S2:48-49  
or obesity, IL-6 elevation in PCOS independent of, 1076-1082
- Sleep apnea syndrome (SAS), genetics of, S2:10
- Sleep deprivation  
interpretation, S2:22  
metabolic and hormonal responses to, S2:21  
neural responses to, S2:21-22  
a neurobiologic and physiologic stressor, S2:20-23  
potential metabolic consequences, S2:11
- Sleep disorders, in psychiatry, S2:40-44
- Sleep phase syndrome  
advanced, S2:8-9  
delayed, S2:9
- Sleep-wake cycle, genetics, and its disorders, S2:7-12
- Sleepwalking, and nocturnal terrors, S2:9
- SM, *see* Skeletal muscle
- SNS, *see* Sympathetic nervous system
- Soleus muscle, metabolic characteristics, in relation to insulin action, in offspring of parents with HTN, 1388-1396
- Soluble adhesion molecules, and CRP, relation to progression of SCI in patients with type 2 diabetes mellitus, 461-466
- Soluble CD40L (sCD40L), and soluble P-selectin, levels in subjects with prediabetes, NIDDM and MS (Chennai Urban Rural Epidemiology Study), 237-242
- Soluble dietary fiber, new, effects on weight gain and selected blood parameters in rats, 195-202
- Soluble intercellular molecule 1, and CRP, relation to progression of SCI in patients with type 2 diabetes mellitus, 461-466
- Soluble P-selectin, and sCD40L, levels in subjects with prediabetes, NIDDM and MS (Chennai Urban Rural Epidemiology Study), 237-242
- South Indians, *see* Race and ethnicity
- Spirolactone, effect on circulating adipocytokines in NIDDM nephropathy patients, 1645-1652
- Standardized breakfast test (SBT), of  $S_i$  and glucose effectiveness, versus minimal model analysis of IVGTT, 676-690
- Starvation, in moderation of long-term alcohol exposure on HSP gene expression in rats, 843-851
- Statins, and LDL apheresis, effect of LDL subtype distribution in patients treated by, 599-604
- Sterol regulatory element-binding protein (SREBP) 2, mRNA abundance of, effect of dietary fatty acids, 635-641
- Sterols, plant, effects of stigmasterol on intestinal absorption of, 292-299
- Stevioside (SVS), effect on glyburide-induced desensitization of the pancreatic beta-cell, in mice, 1674-1680
- Stiffness, carotid artery  
association with plasma adiponectin levels, 587-592  
effect of adiponectin, in NIDDM treated with pioglitazone and metformin, 996-1001  
effect of weight loss, in overweight adults, 907-911
- Stigmasterol, effects on plasma cholesterol levels, hepatic synthesis and intestinal absorption in rat, 292-299
- Streptozotocin-induced diabetes mellitus (STZ-DM)  
in pigs, association of IR with hyperglycemia in, 960-971  
in rats, short-term bioaccumulation of vanadium when ingested with tea decoction in, 263-270
- Stress hormone, and adiponectin, responses to maximal sculling after volume-extended training season in elite rowers, 13-19
- STZ-DM, *see* Streptozotocin-induced diabetes mellitus
- Subjective responses, to brief hypoglycemia in PM women and age-matched men with NIDDM, 331-338
- Sulfonylurea(s)  
in beta-cell exhaustion, S1:23-24  
for beta-cell function loss, S1:3-4  
in body weight gain, S1:22-23  
from experimental to clinical practice on, S1:25  
for hyperglycemia and diabetic complications, S1:2-3  
hypoglycemia induced by, S1:21-22  
long-term efficacy, S1:2-4  
place in NIDDM, S1:20-26  
plus insulin therapy, for NIDDM, S1:4  
plus metformin therapy, for NIDDM, S1:4  
for progressive hyperglycemia, S1:3  
and risk of MI, S1:16-19  
selectivity of, S1:24-25  
therapy for NIDDM, S1:1, S1:2-4
- Supplementation, *see specific supplements*
- Suprachiasmatic nucleus, on sleep and energy balance, S2:31-32
- SVS, *see* Stevioside
- Swedish subjects  
association between insulin and Hcy, 1007-1013  
modification of hormonal situation on lipid effects by lifestyle in middle-aged women, 1060-1066
- Sympathetic nervous system (SNS, Autonomic nervous system), relative overactivity, high sensitivity CRP associated with, in NIDDM with MS, 1014-1021



- Symptoms, and neurocognitive functions, in response to brief hypoglycemia in PM women and age-matched men with NIDDM, 336
- Systemic effects, of hypoglycemia and epinephrine in IDDM, 376
- TAFI, *see* Thrombin activatable fibrinolysis inhibitor
- TC, *see* Total cholesterol
- Tea decoction, short-term bioaccumulation of vanadium when ingested with, 263–270
- Telmisartan  
effect on glucose and lipid metabolism in nondiabetic insulin-resistant subjects, 1149–1154  
effect on S<sub>i</sub>, in nondiabetics with HTN, 1159–1164
- Testosterone  
free, and obesity and CVD risk factors in adolescents with PCOS versus regular cycling adolescents, 508–514  
response to maximal sculling after volume-extended training season in elite rowers, 13–19
- 12-*O*-Tetradecanoyl phorbol-13-acetate (TPA), insulin release induced by, effects of dexamethasone, 39
- TGF- $\beta$ , *see* Transforming growth factor- $\beta$
- TGs, *see* Triglyceride(s)
- 3T3-L1 adipocytes, differentiated, effects of insulin and leptin on fatty acid uptake, 8–12
- Thrombin activatable fibrinolysis inhibitor (TAFI), postprandial, and markers of endothelial dysfunction, in NIDDM, 1437–1442
- Thyroid hormones, metabolism in nonthyroidal tissues, role of thyrotropin, 748–750
- Thyroid-stimulating hormone (TSH), serum, in response to LoBAG diet in men with NIDDM, 248
- Thyrotropin, role in thyroid hormone metabolism in nonthyroidal tissues, 748–750
- T lymphocytes, hepatitis mediated by, role of TNF- $\alpha$  in modulation of adiponectin levels in mice, 555–559
- TNF- $\alpha$ , *see* Tumor necrosis factor- $\alpha$
- ( $\alpha$ )-Tocopherol, and ascorbic acid, effect on down-regulation of Apo A-I gene expression in HepG2 and Caco-2 cell lines, 159–167
- Topiramate, mRNA profiles in liver, hypothalamus, AT and SM of female Zucker diabetic fatty rats after treatment of, 1411–1419
- Total cholesterol (TC)  
effects of PSE supplementation and aerobic exercise training, 541–549  
with HDL-C ratio, relationship between TG/HDL-C ratio, and HDL subclasses, 1141–1148
- Total triiodothyronine, serum, in response to LoBAG diet in men with NIDDM, 248
- Toxemia, associated with pregnancies and deliveries in LPI, 224–231
- TPA, *see* 12-*O*-Tetradecanoyl phorbol-13-acetate
- Training, volume-extended, adiponectin and stress hormone responses to maximal sculling in elite rowers after, 13–19
- Transferrin (Trf), effect on IR of glucose transport, in adipocytes, 1042–1045
- Transforming growth factor- $\beta$  (TGF- $\beta$ ), and TNF- $\alpha$   
effect on PPAR $\gamma$  expression in freshly isolated adipocytes, 313  
expression and action, effect of troglitazone in human adipocyte precursor cells in primary culture, 309–316
- Trf, *see* Transferrin
- Triglyceride(s) (TGs)  
apo E-enriched lipoproteins rich with, accumulation in patients with CAD, 662–668  
effects of PSE supplementation and aerobic exercise training, 541–549  
and FFAs and carnitine, levels in liver and serum of SSADH knockout mice, 355–356  
with HDL-C ratio, relationship between TC/HDL-C ratio, and HDL subclasses, 1141–1148  
and IL-6, adiponectin, leptin and IR in nonobese Japanese NIDDM patients, 258–262  
lipoprotein subspecies rich with, reduction of small dense LDL particles in patients with NIDDM in relationship to, 879–884  
long-chain and structured emulsions, hypertriglyceridemic clamp technique for, in healthy subjects, 1443–1450  
medium-chain, PS mixed with, and HOC oil, effect on plasma lipids in overweight men, 391–395  
metabolism, RAPA in disruption of, in guinea pigs, 794–802  
serum level variations, longitudinal analysis of haplotypes and polymorphisms of APOA5 and APOC3 genes associated with, 1574–1581  
and VLDL, differential effects of Apo E isoforms on lipolysis of, 1129–1134
- Troglitazone  
effect on glucose transport, 313–314  
effect on TNF- $\alpha$  and TGF- $\beta$  expression and action in human adipocyte precursor cells in primary culture, 309–316
- TSH, *see* Thyroid-stimulating hormone
- Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )  
–308G/A polymorphism, relationship with LTA 804C/A and 252A/C polymorphism, in Japanese subjects with NIDDM retinopathy, 1406–1410  
altered release from MNCs of obese reproductive-age women during hyperglycemia, 271–276  
association with insulin-mediated suppression of free fatty acids and net lipid oxidation in HIV-infected patients with lipodystrophy, 175–182  
mRNA, in MNCs, effect of glucose ingestion, in healthy subjects, 1177–1185  
PAI-1 induced by, in regulation of pioglitazone and cerivastatin, in nonmalignant hepatocyte cell line, 1464–1472  
plasma levels and AT gene expression, effect of aerobic training, in obese women, 1375–1381

- Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) (continued)  
  role in modulation of adiponectin levels in T cell–mediated hepatitis in mice, 555–559  
  and TGF- $\beta$   
    effect on PPAR $\gamma$  expression in freshly isolated adipocytes, 313  
    expression and action, effect of troglitazone in human adipocyte precursor cells in primary culture, 309–316  
Turkish subjects, children and adolescents, MS in, 1002–1006  
Twins, adult, interrelation of birth weight and regional lipid deposition in, 561–562  
Type 1 diabetes mellitus, *see* Insulin-dependent diabetes mellitus  
Type 2 diabetes mellitus, *see* Non–insulin dependent diabetes mellitus  
  
UA, *see* Uric acid  
UCP3, *see* Uncoupling protein 3  
UDP-glucuronosyltransferase (UGT), heterologously expressed human, of Crigler-Najjar type II, results of N-glucuronidation of retigabine from, 711–721  
UGT, *see* UDP-glucuronosyltransferase  
UKPDS, *see* United Kingdom Diabetes Study  
Uncoupling protein 3 (UCP3)  
  effects on obesity phenotypes and very low-energy diet-induced changes in overweight Korean female subjects, 578–586  
  mRNA expression, and obesity in ovariectomized rats, effects of NO-1886 (ibrolipim), 151–158  
United Kingdom Diabetes Study (UKPDS), long-term efficacy of sulfonylureas, S1:2–4  
Urea nitrogen, plasma, in response to LoBAG diet in men with NIDDM, 247  
Uric acid (UA)  
  elevated serum levels, in MS, 1293–1301  
  and oxypurinol, plasma concentrations and urinary excretion after rigorous exercise, 103–107  
Urinary excretion  
  of aldosterone and free cortisol, and plasma renin, in response to LoBAG diet in men with NIDDM, 248  
  and plasma concentrations of purine bases and oxypurinol after rigorous exercise, 103–107  
  
V16A polymorphism, Mn-SOD, association with diabetic macular edema, in Korean NIDDM patients, 1681–1688  
Valsartan, effect on protein glycoxidation, 1619–1624  
Vanadium, short-term bioaccumulation of, when ingested with tea decoction in STZ-DM rats, 263–270  
Vascular cell adhesion molecule-1 (VCAM-1)  
  and CRP, relation to progression of SCI in patients with type 2 diabetes mellitus, 461–466  
  serum levels of, in relation to IR and glucose intolerance, 1232–1238  
Vascular endothelial growth factor (VEGF), altered retinal, oxidative stress linked to, activation of NADPH (reduced form) by AGEs associated with, 1516–1523  
Vascular reactivity, differential effects of n-3 PUFAs, in NIDDM mouse, 1365–1374  
VAT, *see* Visceral adipose tissue  
VCAM-1, *see* Vascular cell adhesion molecule-1  
VEGF, *see* Vascular endothelial growth factor  
Very low-density lipoprotein(s) (VLDL)  
  effects of  $\beta$ -adrenergic agonists combined with PPAR in healthy moderately obese subjects, 26–34  
  and TGs, differential effects of Apo E isoforms on lipolysis of, 1129–1134  
Visceral adipose tissue (VAT)  
  as determinant of vWF, in overweight and obese premenopausal women, 650–655  
  long-term exercise on stimulation of AMPK activity and subunit expression in, 1122–1128  
  skeletal muscle, liver, pancreatic islets and, gene expression for energy metabolism after ER in Zucker diabetic fatty rats, 43–52  
VLDL, *see* Very low-density lipoprotein; Very low-density lipoprotein(s)  
VO<sub>2max</sub> during exercise  
  effects of aerobic exercise, in adults, 1500–1507  
  improved, and weight loss, effects of lifestyle modifications on CRP associated with, 825–831  
Voglibose, reduction of oxidative stress markers and soluble ICAM-1, in obese NIDDM patients, 786–793  
von Willebrand factor (vWF), in overweight and obese premenopausal women, VAT as determinant of, 650–655  
  
Waist circumference (WC), relative to BMI, impact on cardiovascular risk factors, 563–569  
Wake centers and neurochemicals, S2:15–17  
Wakefulness  
  cellular and molecular characterization, S2:17  
  dependence on metabolic rate, S2:25–28  
  ischymetric mechanism in control of, S2:26–27  
  messengers of, and joint action on metabolism and feeding, S2:27–28  
  pharmacology of, S2:13–19  
  prolongation through artificial means, S2:17–18  
WAT, *see* White adipose tissue  
WBPTO, *see* Whole body protein turnover  
WC, *see* Waist circumference  
Weight gain, and selected blood parameters, effects of new soluble dietary fiber in rats, 195–202  
Weight loss  
  8% sustainability by metformin-diet over 4 years in women with PCOS, 1582–1589  
  and diet and exercise, body composition and metabolic effects of, on obese HIV-infected women, 1327–1336

## Weight loss (continued)

- effect on  $S_i$  and arterial stiffness, in overweight adults, 907-911
  - and improved aerobic capacity, effects of lifestyle modifications on CRP associated with, 825-831
  - induced by diet and physical activity, decrease CRP concentration after, association with changes in plasma lipids, IL-6 and adiponectin, 359-365
  - resulting from low-fat diet/exercise, effect on LDL particle sized and distribution, in obese women, 1302-1307
  - short-term, effect of weight-bearing aerobic exercise on bone formation during, in overweight obese men and women, 1616-1618
- Western blot, and RT-PCR, for detection of mRNA expressions and protein levels of enzymes in limited amounts of tissue from rat retina, 168-174
- White adipose tissue (WAT), and skeletal muscles, weights of, effects of NO-1886 (ibrolipim) in ovariectomized rats, 154
- White Americans, *see* Race and ethnicity
- Whole-body low-intensity exercise, repeated and prolonged, effects on  $S_i$  and limb muscle adaptations, 217-223
- Whole body protein turnover (WBPTO), at rest in endurance-trained men, impact of dietary proteins, 501-507
- Women
- 64-year-old sample population, relation of serum adiponectin to glucose tolerance, family history of diabetes, autoimmunity,  $S_i$ , C-peptide and inflammation in, 188-194
  - healthy and with PCOS, short-term effects of orlistat on dietary glycotoxins in, 494-500

## Women (continued)

- Japanese, effects of isoflavones and exercise on lipid and bone metabolism in, 423-433
  - obese, *see* Women, obese
  - with PCOS
    - PAI activity, 4G5G polymorphism of PAI-1 gene and first-trimester miscarriage in, 345-352
    - sustainability of 8% weight loss, IR, and amelioration of atherogenic-metabolic risk factors over 4 years by metformin-diet in, 1582-1589
  - postmenopausal, *see* Postmenopausal women
  - premenopausal, *see* Premenopausal women
  - Swedish, middle-aged, modification of hormonal situation on lipid effects by lifestyle, 1060-1066
  - see also* Reproductive-age women
- Women, obese
- effect of aerobic training on plasma levels and AT gene expression of adiponectin, leptin, IL-6 and TNF- $\alpha$ , 1375-1381
  - effect of weight loss resulting from low-fat diet/exercise on LDL particle size and distribution in, 1302-1307
  - HIV-infected, body composition and metabolic effects of diet and exercise, 1327-1336
  - with IGT and normal glucose tolerance, relationship between serum resistin and proinflammatory cytokines, 1495-1499
  - influence of AGT polymorphism (Met235Thr), in obese Japanese women, 819-824
  - and overweight, effect of weight-bearing aerobic exercise on bone formation during short-term weight loss in, 1616-1618
- Xanthine, and oxypurinol, plasma concentrations and urinary excretion after rigorous exercise, 103-107