

Tumor-Specific Anti-Nucleosome Antibody Improves Therapeutic Efficacy of Doxorubicin-Loaded Long-Circulating Liposomes against Primary and Metastatic Tumor in Mice

Tamer A. ElBayoumi[†] and Vladimir P. Torchilin*

Department of Pharmaceutical Sciences and Center for Pharmaceutical Biotechnology and Nanomedicine, Northeastern University, Boston, Massachusetts 02115

Received September 4, 2008; Revised Manuscript Received November 11, 2008; Accepted November 12, 2008

Abstract: The efficacy of drug delivery systems can be significantly enhanced by making them target-specific via the attachment of various ligands to their surface. We attempted to enhance tumor accumulation and therapeutic effect of doxorubicin-loaded long-circulating liposomes (Doxil, ALZA Corp.) by coupling to their surface the anticancer monoclonal antibody 2C5 (mAb 2C5) with nucleosome (NS)-restricted activity, that can recognize the surface of various tumor but not normal cells via the surface-bound nucleosomes released from the apoptotically dying neighboring tumor cells and specifically targets pharmaceutical carriers to tumor cells *in vitro* and *in vivo*. Antibody coupling to PEGylated doxorubicin-liposomes was performed by the “post-insertion” technique. The pharmacokinetics of plain and immuno-targeted Doxil-mimicking liposomes, as well as their accumulation in primary Lewis lung carcinoma (LLC) tumors in mice was followed by real-time gamma-scintigraphy upon liposomal membrane labeling with ¹¹¹In. Therapeutic action of various liposomal formulations was followed by registering primary tumor growth, determining tumor weight upon mice sacrifice, and by counting the number of metastases in the liver and lungs. 2C5 antibody-targeted liposomes demonstrate significantly enhanced accumulation in LLC tumors. Targeted doxorubicin-loaded PEG-liposomes were significantly more effective in inhibiting tumor growth and metastatic process in the LLC tumor models in mice. Our results clearly show the remarkable capability of 2C5-targeted Doxil to specifically deliver its cargo into various tumor manifestations (solid and metastatic) significantly increasing the efficacy of therapy.

Keywords: Tumor targeting; long-circulating liposomes; doxorubicin; anticancer antibody; γ -scintigraphy; tumor growth inhibition; metastases

Introduction

Liposomal formulations of various anticancer drugs are widely used in experimental and clinical oncology with some of them, such as Doxil (doxorubicin incorporated into long-circulating PEGylated liposomes), becoming drugs of choice under certain conditions.^{1–3} The use of the liposomal carrier

allows for decreased side effects of cancer chemotherapeutics, such as nonspecific toxicity, and for enhanced drug delivery into tumors.^{4,5} Thus, Doxil demonstrates decreased

* Corresponding author. Mailing address: Department of Pharmaceutical Sciences, Bouvé College of Health Sciences, Northeastern University, Mugar Building, Room 312, 360 Huntington Avenue, Boston, MA 02115. Tel: 617-373-3206. Fax: 617-373-8886. E-mail: v.torchilin@neu.edu.

[†] Current address: Formulation R&D, Vascular Systems, Atrium Medical Corporation, 5 Wentworth Dr., Hudson, NH 03051. Tel: 603-880-1433, x5441. Fax: 603-880-0313. E-mail: telbayoumi@atriummed.com.

- (1) Skubitz, K. M. Phase II trial of pegylated-liposomal doxorubicin (Doxil) in sarcoma. *Cancer Invest.* **2003**, 21 (2), 167–176.
- (2) Tejada-Berges, T.; Granai, C. O.; Gordinier, M.; Gajewski, W. Caelyx/Doxil for the treatment of metastatic ovarian and breast cancer. *Expert Rev. Anticancer Ther.* **2002**, 2 (2), 143–150.
- (3) Harris, L.; et al. Liposome-encapsulated doxorubicin compared with conventional doxorubicin in a randomized multicenter trial as first-line therapy of metastatic breast carcinoma. *Cancer* **2002**, 94 (1), 25–36.
- (4) Gabizon, A. A.; Lyass, O.; Berry, G. J.; Wildgust, M. Cardiac safety of pegylated liposomal doxorubicin (Doxil/Caelyx) demonstrated by endomyocardial biopsy in patients with advanced malignancies. *Cancer Invest.* **2004**, 22 (5), 663–669.

cardiotoxicity and improved tumor accumulation via the enhanced permeability and retention (EPR) mechanism.^{6,7} Clinically, EPR (“passively”)-targeted PEG-liposomes have been successfully used to deliver doxorubicin to skin, cervical or breast cancers and have been shown to enhance therapeutic activity compared with the same drug administered in the free form or entrapped in conventional liposomes.^{6,8,9} The effective delivery of doxorubicin in PEG-liposomes (Doxil/Caelyx) in the treatment of solid tumors in patients with breast-carcinoma metastases has resulted in a marked increase in survival.^{2,10} Doxil/Caelyx is also in phase II clinical trials for patients with squamous cell cancer of head and neck and ovarian cancer.² The encapsulation of doxorubicin into PEG-liposomes has not only increased the tumor concentration of the drug via the EPR effect but also decreased its volume of distribution, and consequently the therapy-associated toxicity. Still, Doxil is responsible for some nonspecific toxicity (cardiomyopathy and myelosuppression),^{11,12} and additionally provokes certain mucocutaneous reactions.^{12,13} Further decrease in its nonspecific toxicity could be achieved by active targeting of the nanoencapsulated drug into tumors, since one can hypothesize that the active targeting in addition to the EPR-mediated accumulation will accelerate drug accumulation in the tumor, which should result in lesser exposure of normal tissues to the drug, and the presence of the protein molecule (antibody or its fragment) on the preparation will enhance its clearance from the blood decreasing this exposure still further.

Active targeting of long-circulating liposomes to the tumor tissue involves the use of specific “vector” molecules that show affinity toward certain components characteristic

for target tumor tissues.¹⁴ Although the active targeting may further diminish unwanted interactions with normal tissues and cells, its main advantage over the passive targeting is the ability to deliver a larger drug payload directly into the tumor. The use of a targeting moiety could be especially important for tumors with immature vasculature, such as tumors in the early stages of their development, and for delocalized tumors. Vector molecules capable of recognizing tumors include antibodies,¹⁵ peptides,¹⁶ saccharides,¹⁷ hormones, transferrin,¹⁸ and some low molecular weight compounds like folate and some vitamins.¹⁹

Several attempts have been made to further improve the anticancer efficiency of the Doxil by targeting it with vector molecules specific to receptors typical for cancer cells. Thus, an attachment of folate residues to the liposomal surface improved the ability of liposomes to selectively recognize cancer cells that overexpress folate receptors.²⁰ These folate-targeted liposomes loaded with doxorubicin demonstrated a substantial increase in cytotoxicity toward target cells *in vitro*²¹ and *in vivo*.²⁰ Modification with folate promoted the internalization and nuclear delivery of liposomes and overcame multiple drug resistance.¹⁹ Similar notable results were obtained with doxorubicin-loaded long-circulating liposomes modified with RGD-peptide motif targeting the neovasculature of the angiogenic tumors.²² Using a small cell lung cancer cell line, it was shown that targeted liposomes were internalized much faster, delivered doxorubicin to the cell nuclei more efficiently, and were more cytotoxic compared to nontargeted liposomes.²³ Furthermore, doxorubicin-loaded liposomes additionally modified with Fab’ fragments of

- (5) Safra, T.; et al. Pegylated liposomal doxorubicin (doxil): reduced clinical cardiotoxicity in patients reaching or exceeding cumulative doses of 500 mg/m². *Ann. Oncol.* **2000**, *11* (8), 1029–1033.
- (6) Gabizon, A.; Shmeeda, H.; Barenholz, Y. Pharmacokinetics of pegylated liposomal Doxorubicin: review of animal and human studies. *Clin. Pharmacokinet.* **2003**, *42* (5), 419–436.
- (7) Gabizon, A.; et al. Prolonged circulation time and enhanced accumulation in malignant exudates of doxorubicin encapsulated in polyethylene-glycol coated liposomes. *Cancer Res.* **1994**, *54* (4), 987–992.
- (8) Gabizon, A.; et al. Dose dependency of pharmacokinetics and therapeutic efficacy of pegylated liposomal doxorubicin (DOXIL) in murine models. *J. Drug Targeting* **2002**, *10* (7), 539–548.
- (9) Alberts, D. S.; et al. Efficacy and safety of liposomal anthracyclines in phase I/II clinical trials. *Semin. Oncol.* **2004**, *31* (6 Suppl. 13), 53–90.
- (10) Ranson, M. R.; et al. Treatment of advanced breast cancer with sterically stabilized liposomal doxorubicin: results of a multicenter phase II trial. *J. Clin. Oncol.* **1997**, *15* (10), 3185–3191.
- (11) Lyass, O.; et al. Correlation of toxicity with pharmacokinetics of pegylated liposomal doxorubicin (Doxil) in metastatic breast carcinoma. *Cancer* **2000**, *89* (5), 1037–1047.
- (12) Danesi, R.; et al. Pharmacokinetic-pharmacodynamic relationships of the anthracycline anticancer drugs. *Clin. Pharmacokinet.* **2002**, *41* (6), 431–444.
- (13) Charrois, G. J.; Allen, T. M. Drug release rate influences the pharmacokinetics, biodistribution, therapeutic activity, and toxicity of pegylated liposomal doxorubicin formulations in murine breast cancer. *Biochim. Biophys. Acta* **2004**, *1663* (1–2), 167–177.

- (14) Torchilin, V. P. Recent advances with liposomes as pharmaceutical carriers. *Nat. Rev. Drug Discovery* **2005**, *4* (2), 145–160.
- (15) Park, J. W.; et al. Tumor targeting using anti-her2 immunoliposomes. *J. Controlled Release* **2001**, *74* (1–3), 95–113.
- (16) Schifferers, R. M.; et al. Anti-tumor efficacy of tumor vasculature-targeted liposomal doxorubicin. *J. Controlled Release* **2003**, *91* (1–2), 115–122.
- (17) Eliaz, R. E.; Szoka, F. C., Jr. Liposome-encapsulated doxorubicin targeted to CD44: a strategy to kill CD44-overexpressing tumor cells. *Cancer Res.* **2001**, *61* (6), 2592–2601.
- (18) Kobayashi, T.; et al. Effect of transferrin receptor-targeted liposomal doxorubicin in P-glycoprotein-mediated drug resistant tumor cells. *Int. J. Pharm.* **2007**, *329* (1–2), 94–102.
- (19) Goren, D.; et al. Nuclear delivery of doxorubicin via folate-targeted liposomes with bypass of multidrug-resistance efflux pump. *Clin. Cancer Res.* **2000**, *6* (5), 1949–1957.
- (20) Shmeeda, H.; et al. Intracellular uptake and intracavitary targeting of folate-conjugated liposomes in a mouse lymphoma model with up-regulated folate receptors. *Mol. Cancer Ther.* **2006**, *5* (4), 818–824.
- (21) Lee, R. J.; Low, P. S. Folate-mediated tumor cell targeting of liposome-entrapped doxorubicin *in vitro*. *Biochim. Biophys. Acta* **1995**, *1233* (2), 134–144.
- (22) Xiong, X. B.; et al. Enhanced intracellular uptake of sterically stabilized liposomal Doxorubicin *in vitro* resulting in improved antitumor activity *in vivo*. *Pharm. Res.* **2005**, *22* (6), 933–939.
- (23) Xiong, X. B.; et al. Intracellular delivery of doxorubicin with RGD-modified sterically stabilized liposomes for an improved antitumor efficacy: *in vitro* and *in vivo*. *J. Pharm. Sci.* **2005**, *94* (8), 1782–1793.

antidisialoganglioside antibodies selectively and almost completely inhibited the metastatic growth of human neuroblastoma in nude mouse model.²⁴ The modification of Doxil with anti-HER2 monoclonal antibody fragments resulted in a formulation that demonstrated marked anticancer efficiency against tumor lines overexpressing HER2 far superior to that of control nontargeted liposomes both in cell-culture and in *in vivo* models.¹⁵

Earlier, we have identified a family of natural antibodies with nucleosome-restricted specificity capable of effective recognition and binding of a broad variety of cancer cells (but not normal ones) via the cancer cell surface-bound nucleosomes released from apoptotically dying neighboring cancer cells.^{25,26} Extracellular and tumor cell-bound NSs were found in tumor cell cultures,²⁷ and in patients with tumors,²⁸ where they arise from apoptotic cells present in every *in vivo* developing tumor.²⁹ Some antibodies belonging to the group of antinuclear autoantibodies (ANA) demonstrate a NS-restricted specificity being evidently produced as a result of NS uptake by B cells, which, upon the antigen processing display NS histone peptides recognized by appropriate receptors on T helper cells in turn, inducing the presenting B cells to produce ANA with NS-restricted specificity.³⁰

The ability to recognize the surface of tumor but not normal cells was observed for some monoclonal ANAs demonstrating,²⁶ and tumor cell surface-bound NSs were proposed to be their target on tumor cell surface.²⁵ The binding of extracellular NSs to tumor cell surface was hypothesized to be mediated by specific NS receptors reported by several investigators to be present on the surface of tumor cells. Thus, a 94 kDa protein was recognized as a NS receptor in human B-lymphoblastoid Raji cell line, monkey CVI cells, and rat pancreas islet tumoral cell line

RINm,³¹ while another 50 kDa protein domain, calreticulin, was identified as a NS-binding site in a different study.³² For antibody-mediated tumor targeting purposes, it is especially important that NSs are exposed on the cell surface of tumor cells.³³

Functionally, extracellular chromatin fragments have been shown to inhibit the tumor cell killing by NK cells *in vitro*.^{28,34} These findings in fact suggest considering the NS release by dying tumor cells as a tumor self-defense mechanism that protects the surviving tumor cells from host immune attack. In this case, the increased production of NS-specific cytotoxic autoantibodies by a tumor-bearing organism may be considered a response that counteracts tumor self-defense.³³

In addition to their own anticancer activity,^{30,33} these antibodies and their typical representative, the monoclonal antibody 2C5 (mAb 2C5), when used in subtherapeutic quantities, can serve as effective targeting moieties for the tumor-specific delivery of various drug-loaded pharmaceutical nanocarriers.^{35,36} Earlier, we have obtained encouraging data on the increased *in vitro* cytotoxicity of Doxil modified with mAb 2C5.^{37,38} To attach the mAb 2C5 to Doxil liposomes on top of the protective layer of PEG, we have used earlier developed protocol of antibody premodification with *p*-nitrophenyl-carbonyl-PEG-phosphatidyl ethanolamine

- (24) Pastorino, F.; et al. Doxorubicin-loaded Fab' fragments of anti-disialoganglioside immunoliposomes selectively inhibit the growth and dissemination of human neuroblastoma in nude mice. *Cancer Res.* **2003**, 63 (1), 86–92.
- (25) Iakubov, L. Z.; Torchilin, V. P. A novel class of antitumor antibodies: nucleosome-restricted antinuclear autoantibodies (ANA) from healthy aged nonautoimmune mice. *Oncol. Res.* **1997**, 9 (8), 439–446.
- (26) Iakubov, L. Z.; Rokhlin, O.; Torchilin, V. P. Anti-nuclear autoantibodies of the aged reactive against the surface of tumor but not normal cells. *Immunol. Lett.* **1995**, 47 (1–2), 147–149.
- (27) Bell, D. A.; Morrison, B. The spontaneous apoptotic cell death of normal human lymphocytes in vitro: the release of, and immunoproliferative response to, nucleosomes in vitro. *Clin. Immunol. Immunopathol.* **1991**, 60 (1), 13–26.
- (28) Le Lann, A. D.; et al. In vitro inhibition of natural-killer-mediated lysis by chromatin fragments. *Cancer Immunol. Immunother.* **1994**, 39 (3), 185–192.
- (29) Wyllie, A. H. Apoptosis (the 1992 Frank Rose Memorial Lecture). *Br. J. Cancer* **1993**, 67 (2), 205–208.
- (30) Torchilin, V. P.; Iakubov, L. Z.; Estrov, Z. Therapeutic potential of antinuclear autoantibodies in cancer. *Cancer Ther.* **2003**, 1, 179–190.

- (31) Jacob, L.; et al. A monoclonal anti-double-stranded DNA autoantibody binds to a 94-kDa cell-surface protein on various cell types via nucleosomes or a DNA-histone complex. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, 86 (12), 4669–4673.
- (32) Seddiki, N.; et al. Calreticulin, a potential cell surface receptor involved in cell penetration of anti-DNA antibodies. *J. Immunol.* **2001**, 166 (10), 6423–6429.
- (33) Chakilam, A. R.; Pabba, S.; Mongayt, D.; Iakubov, L. Z.; Torchilin, V. P. A single monoclonal antinuclear autoantibody with nucleosome-restricted specificity inhibits the growth of diverse human tumors in nude mice. *Cancer Ther.* **2004**, 2 (4), 353–364.
- (34) Le Lann-Terrisse, A. D.; Fournie, G. J.; Benoist, H. Nucleosome-dependent escape of tumor cells from natural-killer-mediated lysis: nucleosomes are taken up by target cells and act at a postconjugation level. *Cancer Immunol. Immunother.* **1997**, 43 (6), 337–344.
- (35) Elbayoumi, T. A.; Pabba, S.; Roby, A.; Torchilin, V. P. Antinucleosome antibody-modified liposomes and lipid-core micelles for tumor-targeted delivery of therapeutic and diagnostic agents. *J. Liposome Res.* **2007**, 17 (1), 1–14.
- (36) Torchilin, V. P.; Lukyanov, A. N.; Gao, Z.; Papahadjopoulos-Sternberg, B. Immunomicelles: Targeted pharmaceutical carriers for poorly soluble drugs. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, 100 (10), 6039–6044.
- (37) Lukyanov, A. N.; Elbayoumi, T. A.; Chakilam, A. R.; Torchilin, V. P. Tumor-targeted liposomes: doxorubicin-loaded long-circulating liposomes modified with anti-cancer antibody. *J. Controlled Release* **2004**, 100 (1), 135–144.
- (38) Elbayoumi, T. A.; Torchilin, V. P. Enhanced cytotoxicity of monoclonal anticancer antibody 2C5-modified doxorubicin-loaded PEGylated liposomes against various tumor cell lines. *Eur. J. Pharm. Sci.* **2007**, 32 (3), 159–168.

(pNP-PEG-PE) conjugate³⁹ with the subsequent spontaneous “micelle transfer” incorporation of the modified antibody into the membrane of PEGylated liposomes via the hydrophobic PE moiety.⁴⁰ As a result, mAb 2C5-modified Doxil demonstrated significantly higher cytotoxicity toward various cancer cells, including those resistant to doxorubicin, than all control preparations. Our previous data indicated that the specific internalization of the mAb 2C5-Doxil into cytosol, along with the nuclear localization of their drug load, inside the target cancer cells were mainly responsible for the superior anticancer activity.³⁸

We present here the results of our extended *in vivo* studies on the specific tumor-targeting capacity of mAb 2C5-modified doxorubicin-loaded PEGylated liposomes, and their significantly enhanced therapeutic efficacy against both primary and metastatic tumor growths.

Materials and Methods

Materials. Cholesterol (Chol), fully hydrogenated soy phosphatidylcholine (HSPC), *N*-(carbonyl-methoxy-poly(ethylene glycol 2000)-1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine sodium salt (MPEG2000-DSPE), and phosphatidylethanolamine (PE) were from Avanti Polar Lipids, Inc. (Alabaster, AL) and used without further purification. Doxil was purchased from Pharmacia Inc. (West Roxbury, MA). Doxorubicin was purchased from Sigma Chem., Inc. (St. Louis, MO). Triethylamine (TEA), octyl glucoside (OG) and diethylenetriaminepentaacetic acid anhydride (DTPA) were products of Sigma. Poly oxyethylene3400-bis(*p*-nitrophenyl carbonate) (PEG(pNP)₂) was purchased from SunBio (Orinda, CA). Cell culture media, DMEM, fetal bovine serum (FBS) and concentrated solutions of Na-pyruvate, nonessential amino acids, L-glutamine and penicillin/streptomycin stock solutions were from CellGro (Kansas City, MO). Female BALB/c and C57/BL mice were purchased from Charles River Laboratories (Cambridge, MA). All solvents and other chemicals were analytical grade preparations. The mAb 2C5 was produced in ascites via the I.P. injection of 1.5×10^6 hybridoma cells/mL into pristine primed BALB/c 4 week old male mice. The production and the purification of the mAb 2C5 were carried out by Harlan Bioproducts (Indiannapolis, IL) using the cell line from our laboratory. Control IgG2a isotype-matching antibody clone UPC-10 was from Sigma.

Methods. (a) Attachment of Antibody to Liposomes. To prepare antibody (mAb 2C5 or nonspecific IgG) conjugates with PEG3400-PE, an excess of pNP-PEG-PE dis-

persed in 10 mg/mL micellar solution of octyl glucoside in 5 mM Na-citrate, 150 mM NaCl, pH 5.0, was added to an equal volume of 1 mg/mL solution of a protein (mAb 2C5 or IgG) in Tris-buffered saline (TBS), pH 8.5. The mixtures were incubated for 24 h at pH 8.5 at 4 °C.³⁹ To obtain doxorubicin-loaded liposomes modified with mAb 2C5 or nonspecific IgG, the reaction mixtures from the conjugation reactions (see above) were mixed in equal volumes with Doxil and incubated for 5 h at 4 °C. Then, the remaining octyl glucoside and free, nonincorporated mAb 2C5 were removed by dialysis using cellulose ester dialysis tubes with a cutoff size of 300,000 Da. The complete removal of the nonincorporated mAb 2C5 (if such fraction did exist at all) was confirmed by the presence of only one peak of liposomes on the gel-chromatogram of the final sample (no peaks of free mAb 2C5-PEG-PE or mAb 2C5-PEG-PE micelles).³⁷ The absence of the protein band after dialysis in the control sample consisting of a simple mixture of free mAb 2C5 and Doxil liposomes (without the presence of the cross-linker pNP-PEG-PE), indicates that the extensive dialysis process was efficient in removing virtually all antibodies, which could be nonspecifically adsorbed on the liposomal surface, and the modification of the antibodies with the pNP-PEG-PE cross-linker is essential for stable incorporation of the antibody-conjugates onto the liposomal membrane surface.^{37,41} Earlier, it was also shown that the procedure used does not cause any significant loss of doxorubicin by liposomes.^{37,38,42}

(b) Preparation of Control Liposomes. Control PEGylated liposomes mimicking Doxil composition but containing no doxorubicin were prepared using the same lipid components and in the same concentrations as in Doxil. A lipid film was obtained from *N*-(carbonyl-methoxypolyethylene glycol 2000)-1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine sodium salt (3.19 mg/mL), fully hydrogenated soy phosphatidylcholine (HSPC, 9.58 mg/mL), and cholesterol (3.19 mg/mL). The lipid film was suspended in HEPES-buffered saline (HBS), pH 7.4, and sonicated with a probe-type sonicator at 10 W power for 15 min, followed by several passages through mini-extruder with 100 nm pore size polycarbonate filter, until approximately 100 nm liposomes with narrow size distribution were obtained.^{37,38}

All liposomal preparation were controlled for their size (via the dynamic light scattering) and for zeta potential using a Zeta PLUS particle size analyzer with ZETA PALS System, Brookhaven Corp. (Holtsville, NY) at 25 °C.

(c) Cell Cultures. Murine Lewis lung carcinoma (LLC) were purchased from the American type Culture Collection (Manassas, VA), and were maintained in DMEM cell culture medium supplemented with FBS to 10%, Na pyruvate to 1

(39) Torchilin, V. P.; et al. *p*-Nitrophenylcarbonyl-PEG-PE-liposomes: fast and simple attachment of specific ligands, including monoclonal antibodies, to distal ends of PEG chains via *p*-nitrophenylcarbonyl groups. *Biochim. Biophys. Acta* **2001**, *1511* (2), 397–411.

(40) Elbayoumi, T. A.; Torchilin, V. P. Enhanced accumulation of long-circulating liposomes modified with the nucleosome-specific monoclonal antibody 2C5 in various tumours in mice: gamma-imaging studies. *Eur. J. Nucl. Med. Mol. Imaging* **2006**, *33* (10), 1196–1205.

(41) Elbayoumi, T. A.; Torchilin, V. T. Tumor-targeted immunoliposomes for delivery of chemotherapeutics and diagnostics. *J. Pharm. Innovation* **2008**, *3* (1), 51–58.

(42) Elbayoumi, T. A.; Torchilin, V. P. Tumor-specific antibody-mediated targeted delivery of Doxil® reduces the manifestation of auricular erythema side effect in mice. *Int. J. Pharm.* **2008**, *357* (1–2), 272–279.

mM, and penicillin and streptomycin to 50 units/mL and 50 $\mu\text{g/mL}$, respectively.

(d) ^{111}In Radiolabeling of Liposomes. Doxil-mimicking liposomes containing the amphiphilic chelate DTPA-PE (HSPC:CHOL:MPEG2000-DSPE:DTPA-PE in 3:2:0.3:0.3 molar ratios, respectively) were prepared along with the IgG- and 2C5-immuno-analogues. The loading of the liposome-incorporated DTPA-PE with ^{111}In was performed via the transchelation mechanism. DTPA-PE-containing liposomes were incubated for 1 h with ^{111}In chloride in 1 M citrate buffer at RT, and then dialyzed overnight against HBS at 4 $^{\circ}\text{C}$ to remove free label.⁴⁰

(e) Tumor Growth in Mice. Using subcutaneous injection into fat pads in the lower abdominal region, LLC tumors were implanted in 8 week old C57/BL mice (5×10^4 cells/mouse) and in 6–8 week old BALB/c mice (3×10^5 cells/mouse), for therapeutic and gamma-imaging studies, respectively. The time for the appearance of the tumor varies from one cell line to another and usually takes 10–14 days. Mice were constantly monitored and allowed for free access to food and water (following the animal care protocol no. R01210 approved by Northeastern University Institutional Animal Care and Use Committee, in accordance with the “Principles of laboratory animal care”, NIH publication no. 85-23, revised in 1985) and tumor volumes were calculated using the formula $0.5(\text{length} \times \text{width}^2)$ by measuring the dimensions of the tumor at regular time intervals.

(f) Tumor Accumulation of ^{111}In -Labeled Liposomes in Mice. When the tumor diameter reached 5–8 mm, mice were injected with 0.1 mL of 4 mg/mL ^{111}In -radiolabeled Doxil-mimicking liposomal formulations via the lateral tail vein. The accumulation of ^{111}In -radiolabeled Doxil-mimicking liposomal formulations in the developed tumor was also visualized using an Ohio Nuclear 400 radio-isotope camera (Ohio-Nuclear Inc., Solon, OH) equipped with a high energy collimator and NU Mac computer (NC systems, Boulder, CO) at 2, 4 and 6 h postinjection after anesthetizing the mice by injecting a mixture of xylazine and ketamine intraperitoneally. Digital pictures of the tumor-bearing mouse were taken using a Kodak digital camera (Eastman Kodak Company, Rochester, NY).³⁵ Images were analyzed using ImageJ software (U.S. National Institutes of Health, Bethesda, MD) for calculation of mean signal intensity per equivalent area of selected tumor location.

(g) Single-Dose Pharmacokinetics of ^{111}In -Labeled Doxil-Mimicking Liposomes. *In vivo* biodistribution studies of ^{111}In -radiolabeled Doxil-mimicking liposomal formulation and their IgG and mAb 2C5 analogues were performed in healthy 8 week old female BALB/c mice in two separate experiments. In each experiment, 72 BALB/c mice were injected with 0.1 mL of 4 mg/mL ^{111}In radiolabeled Doxil-mimicking liposomal formulations via the lateral tail vein. At time points of 15, 30, 120, 360, 720, 1440 min postinjection, blood was collected using a Pasteur pipet from the retro-orbital plexus of the eye, and then, the mice were euthanized with carbon dioxide followed by organ collection. Each time point had at least 4 mice and the organs collected

were liver, kidney, spleen, lung, muscle and tail. The amount of the radioactivity was quantified as CPM using a Beckman 5500B gamma-counter (Beckman Instruments, Columbia, MD) and the amount of the accumulated radioactivity per gram of tissue was calculated followed by calculation of the pharmacokinetic parameters like elimination rate constant (K_{el}), area under the curve (AUMC), and half-life (MRT) using noncompartmental statistical analysis of the blood data, using PK Solutions 2.0 (Summit research Services, Montrose, CO).^{35,40}

(h) *In Vivo* Therapeutic Efficacy of 2C5-Modified Doxil in Murine Tumor Models. Therapeutic treatment of LLC tumor-bearing mice started 5 days after tumor implantation with the dose intensity of 2 mg doxorubicin/kg/every 5 days. Each treatment group consisting of 8–10 mice received the corresponding doxorubicin liposomal formulation for 4 times. Tumor diameters were measured and approximate tumor volume was calculated as $0.5(\text{length} \times \text{width}^2)$ ²⁴ every 3 days during the treatment period to monitor tumor growth. 35 days after the first injection, mice were sacrificed and tumors were excised and weighed.

(i) Histological Evaluation. After 35 days of initiation of treatment, lungs and livers were surgically removed and weighed, then fixed for 24 h in Bouins solution (15:5:1 saturated picric acid:neutral buffer formalin:acetic acid). Afterward, samples were submerged in 95% alcohol for 24 h. Finally, metastatic nodules/foci per mouse were visually counted and the mean value was estimated for each sample group. Then, metastasis percentage, determined by percent of mice showing any metastasis in each group, was calculated.^{43,44}

(j) Statistics. Differences in the apparent tumor volumes during treatment and postmortem tumor weights were compared using Student's *t* test for two independent samples and Kruskal–Wallis analysis with Tukey's HSD Post-Hoc test for all pair mean comparisons, for three or more independent samples. These tests were analyzed using kaledagraph software, ver. 3.6 (Synergy Software, Reading, PA). For easy representation of the difference in tumor volumes during the treatment between the control and treated groups, the average tumor volume (mm^3) of the group vs days was plotted. Tumor weights were plotted as box plots, which represent the median, quartiles and extremes in a group.

Results

The method of antibody attachment to liposomes by transferring PEG-PE-modified antibodies from their loose micelles onto the liposome surface used in this study results

(43) Lu, D. Y.; Xu, B.; Ding, J. Antitumor effects of two bisdioxopiperazines against two experimental lung cancer models *in vivo*. *BMC Pharmacol.* **2004**, *4* (1), 32.

(44) Nakamura, K.; et al. Inhibitory effect of Cordyceps sinensis on spontaneous liver metastasis of Lewis lung carcinoma and B16 melanoma cells in syngeneic mice. *Jpn. J. Pharmacol.* **1999**, *79* (3), 335–341.




Figure 1. Whole body imaging of LLC-tumor-bearing mice 4 h after injection with ^{111}In -labeled liposomes. Panel I: calculated radiologic signal intensity. Panel II: Gamma-scintigraphic images of mice injected with (A) control Doxil-mimicking liposomes; (B) nonspecific IgG-modified Doxil-mimicking liposomes; (C) mAb 2C5-modified Doxil-mimicking liposomes. Circles indicate tumor locations.

in attaching approximately 70–80 antibody molecules per single 100 nm liposome.^{37,38} The antibody incorporation was not accompanied by any significant loss of the liposomal doxorubicin, and virtually no difference in drug release was registered between the original Doxil and the immuno-Doxil formulations (modified with either the IgG-PEG-PE or the mAb 2C5-PEG-PE) in 48 h *in vitro* release experiment.^{37,38,42} Immuno-modification of Doxil did not noticeably change the liposome size (remained in the range of approximately 90 to 120 nm) or the net charge at the liposome surface (zeta potential of -25 to -23).^{37,38,40} The physicochemical characterization of immunoliposomes similar to those used in this study were reported by us earlier, and it could be useful to mention here that the number of liposome-bound antibodies was estimated as approximately 80 molecules per single liposome.^{37,38,40}

It was also repeatedly shown that liposomes, including doxorubicin-containing PEGylated liposomes, modified with mAb 2C5-PEG-PE demonstrate strong and specific binding with mAb 2C5 antigen, nucleosomes, while nonspecific IgG-modified liposomes and nonmodified liposomes do not bind with the nucleosome monolayer.^{37,38}

In Vivo Tumor Accumulation of mAb 2C5-Modified Liposomes. The investigation of tumor accumulation of ^{111}In -labeled Doxil-mimicking liposomes and their IgG and mAb 2C5 immuno-analogues in LLC murine tumor model clearly demonstrated an enhanced tumor accumulation of 2C5-modified liposomes. The intensity of the radiologic signal accumulated in primary subcutaneous (sc) tumor corresponding to mAb 2C5-modified liposomes, was scaled almost 2–3 times that of the nonspecific IgG-bearing or plain liposomes formulations, measured at the same time (see Figure 1). The whole body direct gamma-imaging of tumor-bearing mice, starting only 4 h postinjection time-point, clearly confirmed the enhanced accumulation of mAb 2C5-targeted preparations in both tumors.

Single Dose Pharmacokinetics of ^{111}In -Labeled Doxil-Mimicking Liposomes. The pharmacokinetic behavior of various liposomal preparations was studied in healthy




Figure 2. Single dose pharmacokinetics of various ^{111}In -labeled Doxil-mimicking liposomal preparations in healthy BALB/C mice ($n = 5$, results indicated \pm SD).

BALB/c mice (Figure 2). Similar to the kinetic profile of analogous preparation in 4T1-tumor bearing mice,⁴⁰ blood clearance data show that both IgG and mAb 2C5 immuno-analogues of the ^{111}In -labeled Doxil-mimicking liposomes expectedly cleared somewhat faster from the body compared to plain liposomes, yet both formulations retained the ability to long circulate in the blood, with half-lives (MRT values) of about 16.2 ($K_{el} = 0.065 \text{ h}^{-1}$) and 15.8 ($K_{el} = 0.067 \text{ h}^{-1}$) hours, respectively, compared to 21.6 h ($K_{el} = 0.045 \text{ h}^{-1}$) in case of the nonmodified formulation. This corresponds well to results reported earlier with other targeted liposomes.²⁴ Furthermore, the decrease noticed in the calculated area under the curves (AUC) for the Doxil-mimicking liposomes ($\text{AUC} = 13630\% \text{ cpm} \cdot \text{h}^2/\text{g}$) was less than 2-fold in case of immuno-modification with either IgG ($\text{AUC} = 6930\% \text{ cpm} \cdot \text{h}^2/\text{g}$) or mAb 2C5 ($\text{AUC} = 6380\% \text{ cpm} \cdot \text{h}^2/\text{g}$).

The observed slight increase in the clearance of the antibody-modified liposomes from the blood could be also explained by the presence of whole antibodies on the liposomal surface, which makes them more liable to be taken up by RES via the Fc-mediated uptake. The use of smaller

mAb 2C5 fragments, such as Fab', to prepare cancer-targeted immunoliposomes, could effectively prevent this Fc receptor-mediated clearance by RES, although this was not specifically studied in our set of experiments.

Previous work with LLC-tumor bearing mice for the biodistribution of the radiolabeled liposomes, in various organs over 48 h, indicated that while immuno(IgG- or mAb 2C5-modified)-liposomes are mainly excreted by liver and spleen; they do have negligible accumulation in nontarget tissues.⁴⁰

In Vivo Therapeutic and Antimetastatic Activity of 2C5-Modified Doxil in Murine Lewis Lung Carcinoma Model. Based on previous therapeutic studies of Doxil in murine tumor models,⁴⁵ the dose of 2.2 mg of doxorubicin/kg/every 5 days (corresponds to about 6 half-lives for the original Doxil and approximately 7 half-lives of the antibody-modified Doxil formulations, which means almost complete elimination of drug by the time for the following dose) for four consecutive administrations was chosen (i.e., total dose given is just about 9 mg/kg). Some previous studies with Doxil in similar models used similar dosing intensity of 9 mg/kg/every two weeks for two doses or 4.5 mg/kg/every 3 days for four doses (i.e., total dose equals 18 mg/kg).⁴⁵ Using the selected treatment regimen, the therapeutic efficacy of mAb 2C5-modified Doxil was compared to that of both the nonspecific IgG-modified Doxil and the original Doxil formulations, in addition to the negative control PBS injections. The effect was assessed in terms of the average tumor volume observed during treatment period of 36 days and in terms of the average postmortem tumor weight at the end of the treatment period.

In this study, murine Lewis lung carcinoma (LLC) tumor model, which produces lung and liver metastasis when implanted subcutaneously, was used, and the assessment for the therapeutic efficacy against the primary tumor growth followed the criteria as mentioned above. Considering the primary tumors, the results show that compared to all controls, the mAb 2C5-modified Doxil, as with other tumors, demonstrated a significantly enhanced therapeutic efficacy in mice expressed as smaller tumor volumes starting at 20 days after tumor implantation ($p \leq 0.05$). The final average tumor weight in animals treated with this preparation was only ca. 0.52 g compared to ca. 1.6 g in animals treated with IgG-modified Doxil or original Doxil and ca. 2.3 g in animals receiving only the buffer injections (Figure 3A,B).

As follows from Tables 1 and 2, the metastatic growth in lungs and livers of host animals was also most significantly inhibited in the case of mAb 2C5-modified Doxil. There was a significant difference in the number of lung and liver metastases between the targeted preparation-treated group and control groups – untreated and treated with unmodified Doxil. Although, the unmodified Doxil significantly inhibits the metastatic process compared to untreated controls, its




Figure 3. Therapeutic activity, expressed as tumor volumes (A) and postmortem tumor weight (B) of 2C5-modified Doxil against control preparations in LLC-implanted mice. Arrows indicate treatment schedule, 2 mg/kg/q 5 days ($n = 8-10$), * ($P \leq 0.05$, vs corresponding IgG-Doxil), nonparametric Kruskal–Wallis with Tukey's HSD Post-Hoc test. (Results presented as mean \pm SD).

Table 1. Lung Metastases in LLC Tumor-Bearing Mice^a

treatment	no. of lung metastatic foci	% of metastases (no. of mice with metastases/gp)
PBS control	43 \pm 11 a,b	80 (8/10)
Doxil	24 \pm 8 a,c	45 (4/9)
mAb 2C5-modified Doxil	11 \pm 1.5 b,c	22 (2/9)

^a At a, b, c; $P \leq 0.05$.

specific targeting to tumors with mAb 2C5 significantly enhanced this inhibition.

One has to note here that although the high accumulation of immunoliposomes in liver and spleen as the ultimate consequence of the RES-mediated clearance is a well-known phenomenon, it is not very likely that liposomes residing in the liver for metabolism can exert a significant therapeutic effect systemically serving as a secondary drug depot. One observation supporting the absence of the depot effect is the rapid therapeutic effect of the treatment on the implanted tumors, where within about a week of therapy and right after

(45) Charrois, G. J.; Allen, T. M. Multiple injections of pegylated liposomal Doxorubicin: pharmacokinetics and therapeutic activity. *J. Pharmacol. Exp. Ther.* **2003**, 306 (3), 1058–1067.

Table 2. Liver Metastases in LLC Tumor-Bearing Mice^a

treatment	rel liver wt (%)	no. of liver metastatic foci	% of metastases (no. of mice with metastases/gp)
PBS control	5.1 ± 0.12	52 ± 17 a,b	60 (6/10)
Doxil	4.94 ± 0.07	33 ± 10.5 a	22 (2/9)
mAb 2C5-modified Doxil	4.76 ± 0.05	18 b	11 (1/9)

^a At a, b; *P* ≤ 0.05.

the second dose, the tumor volumes of Doxil-treated animals starts to markedly decrease, with clearly superior tumor size reduction in case of targeted Doxil. This indeed corresponds well with the earlier observation of the fast high tumor-specific accumulation of the immuno-targeted liposomes rather than suggesting any organ-based depot effect. At the same time, one cannot reject the possibility of a certain effect of the drug escaping cellular metabolism to affect neighboring metastatic foci in close vicinity within liver tissues, especially if these cancerous metastases were transformed/derived from the original Kupffer cells.

Discussion

The toxic side-effects of anticancer drugs, resulting from the lack of specificity of conventional therapies, usually limit the increase in dose intensity often required to eradicate the cancerous growth. One of the most evolved strategies that have been developed in the last two decades was the enhancement of the tumor specific delivery of the chemotherapeutic drug, mainly anthracyclins, through actively targeting them via monoclonal antibodies capable of selectively binding antigens that are overexpressed on cancer cells. At the same time, the clinical support received for several liposomal formulations of doxorubicin (Doxil/Caelyx) and daunorubicin (Daunoxome) was entirely based on their unique ability to improve the pharmacokinetics of the liposome-laden drugs. Yet, despite the advances with prolonged circulation and liposome accumulation in the targeted tumor interstitium, the current clinically approved Doxil formulation has resulted in only a moderate increased antitumor efficacy. Moreover, while this formulation has shown a substantial reduction of the toxicity profile of the free doxorubicin, these long-circulating liposomes suffered from the introduction of new side effects, mainly skin toxicity manifested as hand-foot syndrome and mucositis.^{46,47} These adverse effects are virtually inherent to the Stealth liposomal carriers and can be overcome by modification of the liposome composition. The active targeting of these liposomes via monoclonal antibodies and their fragments has received extensive research attention in the past decade, in order to

render them more tumor-avid, hence facilitating the delivery of these liposomes with their drug cargoes into the cellular site of drug action and evading undesired side effects. This line of research stimulated our attempts to utilize the mAb 2C5 with its unique ability to specifically recognize a broad variety of tumors, as a targeting ligand to actively target Doxil/Caelyx, toward aggressive tumors *in vivo*.⁴²

From our earlier *in vitro* experiments,^{35,37,38} the remarkable anticancer activity of mAb 2C5-Doxil against the various tumor cell lines of diverse origins was evident, mainly due to mAb 2C5 proven superior specific surface cell binding and subsequent specific uptake inside the cancer cells. These experiments naturally led us to a comprehensive set of *in vivo* studies.

Using the LLC murine *in vivo* model, the biodistribution data demonstrated that tumor accumulation ratios of ¹¹¹In radiolabeled 2C5-modified Doxil-mimicking liposomes (prepared by adding 0.5 mol % of amphiphilic chelate, DTPA-PE to lipid composition), compared to the neighboring muscle were almost double that of the nontargeted formulations after 24 h, and even after 48 h, despite their relatively rapid clearance.⁴⁰

Moreover, the whole body gamma-imaging of the LLC tumor-bearing mice models, at 4 h postadministration, confirms the tumor-preferential distribution of the 2C5-modified Doxil-mimicking liposomes compared to control formulations. Based on the pharmacokinetics and the biodistribution profile of the immunoliposomes, it is warranted to assume that this mAb 2C5-modification of the Doxil-mimicking liposomes has made them significantly less EPR-dependent, and a much fewer number of passages of the 2C5-liposomes in a relatively short period of time (approximately 4–6 h) has resulted in a great increase in the tumor accumulation of these liposomes, and consequently a markedly stronger tumor-specific signal compared to that of the EPR-dependent, plain Doxil-mimicking liposomes.

The pharmacokinetic analysis indicated that both IgG and mAb 2C5 immuno-analogues of the ¹¹¹In-labeled Doxil-mimicking liposomes suffered from faster clearance from the body, mainly through the uptake by RES organs: liver and spleen. In spite of that, both immuno-formulations retained sufficient ability to circulate long in the blood for about 16 h, compared to 21 h in the case of the nonmodified formulation. This fairly hassled clearance of the antibody-modified liposomes from the body was expected, owing to the presence of whole antibodies on the liposomal surface, which makes them susceptible to the uptake by the RES. The use of smaller mAb 2C5 fragments such as Fab' to prepare cancer-targeted immuno-liposomes should also be possible and will probably improve the blood residence time of the mAb 2C5-modified liposomes to be close to that of the plain liposomes, as reported with other groups.⁴⁸ The

(46) Uziely, B.; et al. Liposomal doxorubicin: antitumor activity and unique toxicities during two complementary phase I studies. *J. Clin. Oncol.* **1995**, *13* (7), 1777–1785.

(47) Lotem, M.; et al. Skin toxic effects of polyethylene glycol-coated liposomal doxorubicin. *Arch. Dermatol.* **2000**, *136* (12), 1475–1480.

(48) Brignole, C.; et al. Development of Fab' fragments of anti-GD(2) immunoliposomes entrapping doxorubicin for experimental therapy of human neuroblastoma. *Cancer Lett.* **2003**, *197* (1–2), 199–204.

gamma-tracking of the radiolabeled liposomal formulations along with the biodistribution data in fact allows for considering the decrease in circulation half-life of the radiolabeled liposomes due to the immuno-modification with mAb 2C5 as having an only a minor effect on the development of a sharper and faster tumor-specific signal this preparation actually produces.

The selection of the suitable dose regimen, 2.2 mg/kg/q 5 days for 4 doses with total dose of 9 mg/kg, was based upon both the pharmacokinetic data of the liposomal formulations and the effective Doxil therapeutic regimen used in similar tumor models (total therapeutic dose of 18 mg/kg). It was chosen as a subtherapeutic dose of Doxil in order to be able to investigate the statistical significance of the therapeutic difference between the various Doxil formulations. Considering the primary sc solid tumor growth in the tested LLC lung tumor model, compared to all controls, the mAb 2C5-modified Doxil demonstrated a significantly improved therapeutic efficacy in mice, resulting in markedly smaller tumor volumes starting less than three weeks after tumor implantation, and significantly lower final average tumor weights, compared to other treatments ($p \leq 0.05$). These impressive therapeutic results confirm the important role of the tumor-specific accumulation of the mAb 2C5-modified Doxil in achieving high drug concentration in the tumor. Although the diffusion and deep tumor penetration are not expected for particles of the size of liposomes, especially in case of large tumors with high interstitial pressure, making our approach more applicable for smaller tumor, however it is quite likely that the encapsulated drug, being much more diffusible than the carrier itself, can reach distant tumor cells surrounding the area of extravasated liposomes, and exert such a significant antineoplastic effect.

One can assume that a significantly enhanced effect of the mAb 2C5-targeted formulation on the developments of metastases in the case of LLC liver and lung metastases compared to the original Doxil ($p \leq 0.05$) is in a sense limited by the fact that most metastatic nodules are considerably smaller in size than primary tumors, and with no or very small necro-apoptotic cores, compared to solid primary LLC tumors. Hence, they may have limited amounts of NSs bound onto their surfaces and available as targets for the recognition by the mAb 2C5-modified Doxil. Hypothetically,




Figure 4. Schematics of tumor targeting and uptake of nucleosome-specific monoclonal antibody 2C5-modified Doxil liposomes, for both primary site and metastatic cells.

the pretreatment of a cancerous condition with pro-apoptotic drugs can increase the apoptotic death of cancer cells in the body, nucleosome release, and nucleosome attachment to cancer cells in metastatic sites making them better targets for mAb 2C5-modified drugs and drug delivery systems.⁴⁹ The general scheme of targeting primary tumors and metastatic cells with mAb 2C5-modified drug-loaded long-circulating liposomes is presented in Figure 4.

Conclusions

The present study provides clear evidence that the simple immuno-modification of Doxil with the cancer-specific antinucleosome mAb 2C5 resulted in the successful development of a strong tumor-targeted doxorubicin delivery system, demonstrating both prolonged circulation in the body and fast and specific accumulation in solid LLC tumors. This resulted in a significant inhibition of both primary and metastatic growths of this aggressive tumor in mice. The mAb 2C5-modified Doxil provides a promising opportunity to further enhance the therapeutic index of the original formulation and broaden the spectrum of its antitumor activity.

Acknowledgment. This work was supported by NIH Grant 2R01 HL55519 to V.P.T.

MP8001528

- (49) Iakoubov, L. Z.; Torchilin, V. P. Nucleosome-releasing treatment makes surviving tumor cells better targets for nucleosome-specific anticancer antibodies. *Cancer Detect. Prev.* **1998**, 22 (5), 470–475.