

Biorecognition and Subcellular Trafficking of HPMA Copolymer–Anti-PSMA Antibody Conjugates by Prostate Cancer Cells

Jihua Liu,[†] Pavla Kopečková,^{†,‡} Patrick Bühler,[§] Philipp Wolf,[§] Huaizhong Pan,[†]
Hillevi Bauer,[†] Ursula Elsässer-Beile,[§] and Jindřich Kopeček^{*,†,‡}

Department of Pharmaceutics and Pharmaceutical Chemistry and Department of Bioengineering, University of Utah, Salt Lake City, Utah, and Department of Urology, Experimental Urology, University of Freiburg, Freiburg, Germany

Received December 29, 2008; Revised Manuscript Received March 31, 2009; Accepted April 3, 2009

Abstract: A new generation of antibodies against the prostate specific membrane antigen (PSMA) has been proven to bind specifically to PSMA molecules on the surface of living prostate cancer cells. To explore the potential of anti-PSMA antibodies as targeting moieties for macromolecular therapeutics for prostate cancer, fluorescently labeled HPMA (*N*-(2-hydroxypropyl)methacrylamide) copolymer–anti-PSMA antibody conjugates (P-anti-PSMA) were synthesized and the mechanisms of their endocytosis and subcellular trafficking in C4-2 prostate cancer cells were studied. Radioimmunoassays showed the dissociation constants of P-anti-PSMA for C4-2 prostate cancer cells in the low nanomolar range, close to values for free anti-PSMA. It indicated that conjugation of anti-PSMA to HPMA copolymers did not compromise their binding affinity. The rate of endocytosis of P-anti-PSMA was much faster than that of control HPMA copolymer conjugates containing nonspecific IgG. Selective pathway inhibitors of clathrin-mediated endocytosis and of macropinocytosis inhibited the internalization of P-anti-PSMA. Inhibition of clathrin-mediated endocytosis was further evidenced by down-regulation of clathrin heavy chain expression by siRNA. Using a dominant-negative mutant of dynamin (Dyn K44A) to abolish the clathrin-, caveolae-independent endocytic pathway, we found that some of P-anti-PSMA adopted this pathway to be endocytosed into C4-2 cells. Thus multiple receptor-mediated endocytic pathways, including clathrin-mediated endocytosis, macropinocytosis, and clathrin-, caveolae-independent endocytosis, were involved in the internalization of P-anti-PSMA. The extent of the participation of each pathway in P-anti-PSMA endocytosis was estimated. Membrane vesicles containing P-anti-PSMA rapidly colocalized with membrane vesicles overexpressing Rab7, a late endosome localized protein, demonstrating that a part of P-anti-PSMA was transported to late endosomes.

Keywords: HPMA copolymer; drug delivery; antibody targeting; endocytosis; clathrin-mediated endocytosis

Introduction

Polymer therapeutics including polymer–protein conjugates, drug–polymer conjugates, supramolecular, and other

nanosized drug delivery systems represent a compensatory and promising approach on the improvement of cancer treatment, due to lack of tumor selectivity of most low-molecular-weight anticancer chemotherapeutic agents. Conjugating low-molecular-weight anticancer drugs to polymers establishes (passive) tumor selectivity due to the enhanced permeability and retention (EPR) effect.¹ However, one way

* Corresponding author. Mailing address: Department of Pharmaceutics and Pharmaceutical Chemistry, University of Utah, 30 S. 2000 E. Rm. 201, Salt Lake City, UT 84112-5820. Phone: (801) 581-7211. Fax: (801) 581-7848. E-mail: Jindrich.Kopecek@utah.edu.

[†] Department of Pharmaceutics and Pharmaceutical Chemistry, University of Utah.

[‡] Department of Bioengineering, University of Utah.

[§] University of Freiburg.

(1) Shiah, J. G.; Dvořák, M.; Kopečková, P.; Sun, Y.; Peterson, C. M.; Kopeček, J. Biodistribution and antitumor efficacy of long-circulating *N*-(2-hydroxypropyl)methacrylamide copolymer-doxorubicin conjugates in nude mice. *Eur. J. Cancer* **2001**, *37*, 131–139.

to achieve high local concentration of polymer therapeutics in tumor tissues is incorporation of a targeting moiety able to actively guide polymer therapeutics to the tumor sites.² Clinical success of monoclonal antibodies bodes well for their use as targeting moieties³ in drug delivery systems. Indeed, targeted polymer therapeutics have improved the therapeutic index with minimal side effects in both preclinical and clinical settings.^{4–7}

The incorporation of OV-TL16 antibody, recognizing the CD47 (OA-3) antigen expressed on most of human ovarian carcinomas, into *N*-(2-hydroxypropyl)methacrylamide (HPMA) copolymer based polymer therapeutics resulted in their enhanced accumulation in human ovarian OVCAR-3 carcinoma xenografts with concomitant decrease in tumor proliferation.⁴ More efficient tumor targeting and higher antitumor efficacy were demonstrated in targetable HPMA copolymer–drug conjugates mediated by Fab' antibody fragment from OV-TL16 antibody than the nontargeted HPMA copolymer drug conjugate.⁸

Identification of tumor cell surface specific molecules enables development of targeted anticancer polymer therapeutics. Prostate specific membrane antigen (PSMA) is highly expressed in prostate cells with a minimal expression in tissues of the brain, proximal small intestine, salivary glands, and kidney.^{9–11} The expression of PSMA is upregulated in malignant disease, with the greatest level detected in metastatic androgen independent prostate cancer.^{12,13} It has been found also abundantly expressed in tumor-associ-

ated neovasculature in a variety of solid tumors.^{10,14} PSMA is a dimeric type II integral membrane glycoprotein with an intracellular segment, a transmembrane domain, and an extracellular domain.¹¹

Recently, Elsässer-Beile and co-workers developed a new generation of monoclonal antibodies and recombinant antibody fragments against cell-adherent PSMA.¹⁵ In their studies, mAbs and scFv from mice immunized with the native form of PSMA demonstrated binding activity and internalization into LNCap cells.¹⁵ In addition, the antibodies possessed great specificities, verified by lack of binding activity toward a variety of PSMA-negative cells. A recombinant immunotoxin based on the anti-PSMA antibody possessed selective toxicity to prostate cancer cells.^{16,17} Moreover, a bispecific diabody against PSMA and CD3 was active in T-cell mediated lysis of prostate cancer cells.¹⁸

In addition to antibodies, aptamers, synthetic oligonucleotides, were identified as capable of binding to PSMA.¹⁹ Highly selective and effective therapeutic modalities based on aptamers have been developed to deliver chemotherapeutics, functional siRNA and toxin for the treatment of

- (2) Allen, T. M. Ligand-targeted therapeutics in anticancer therapy. *Nat. Rev. Cancer* **2002**, 2, 750–763.
- (3) Reichert, J. M.; Rosensweig, C. J.; Faden, L. B.; Dewitz, M. C. Monoclonal antibody successes in the clinic. *Nat. Biotechnol.* **2005**, 23, 1073–1078.
- (4) Shiah, J. G.; Sun, Y.; Kopečková, P.; Peterson, C. M.; Straight, R. C.; Kopeček, J. Combination chemotherapy and photodynamic therapy of targetable *N*-(2-hydroxypropyl)methacrylamide copolymer-doxorubicin/mesochlorin *e*₆-OV-TL 16 antibody immunconjugates. *J. Controlled Release* **2001**, 74, 249–253.
- (5) Lu, Z. R.; Kopečková, P.; Kopeček, J. Polymerizable Fab' antibody fragments for targeting of anticancer drugs. *Nat. Biotechnol.* **1999**, 17, 1101–1104.
- (6) Peer, D.; Karp, J. M.; Hong, S.; Farokhzad, O. C.; Margalit, R.; Langer, R. Nanocarriers as an emerging platform for cancer therapy. *Nat. Nanotechnol.* **2007**, 2, 751–760.
- (7) Seymour, L. W.; Ferry, D. R.; Anderson, D.; Hesslewood, S.; Julyan, P. J.; Poyner, R.; Doran, J.; Young, A. M.; Burtles, S.; Kerr, D. J. Hepatic drug targeting: Phase I evaluation of polymer-bound doxorubicin. *J. Clin. Oncol.* **2002**, 20, 1668–1676.
- (8) Lu, Z. R.; Shiah, J. G.; Kopečková, P.; Kopeček, J. Polymerizable Fab' antibody fragment targeted photodynamic cancer therapy in nude mice. *STP Pharma Sci.* **2003**, 13, 69–75.
- (9) Fair, W. R.; Israeli, R. S.; Heston, W. D. Prostate-specific membrane antigen. *Prostate* **1997**, 32, 140–148.
- (10) Silver, D. A.; Pellicer, I.; Fair, W. R.; Heston, W. D.; Cordon-Cardo, C. Prostate-specific membrane antigen expression in normal and malignant human tissues. *Clin. Cancer Res.* **1997**, 3, 81–85.
- (11) Israeli, R. S.; Powell, C. T.; Corr, J. G.; Fair, W. R.; Heston, W. D. Expression of the prostate-specific membrane antigen. *Cancer Res.* **1994**, 54, 1807–1811.

- (12) Kawakami, M.; Nakayama, J. Enhanced expression of prostate-specific membrane antigen gene in prostate cancer as revealed by in situ hybridization. *Cancer Res.* **1997**, 57, 2321–2324.
- (13) Wright, G. L., Jr.; Grob, B. M.; Haley, C.; Grossman, K.; Newhall, K.; Petrylak, D.; Troyer, J.; Konchuba, A.; Schellhammer, P. F.; Moriarty, R. Upregulation of prostate-specific membrane antigen after androgen-deprivation therapy. *Urology* **1996**, 48, 326–334.
- (14) Liu, H.; Moy, P.; Kim, S.; Xia, Y.; Rajasekaran, A.; Navarro, V.; Knudsen, B.; Bander, N. H. Monoclonal antibodies to the extracellular domain of prostate-specific membrane antigen also react with tumor vascular endothelium. *Cancer Res.* **1997**, 57, 3629–3634.
- (15) Elsässer-Beile, U.; Wolf, P.; Gierschner, D.; Bühler, P.; Schultze-Seemann, W.; Wetterauer, U. A new generation of monoclonal and recombinant antibodies against cell-adherent prostate specific membrane antigen for diagnostic and therapeutic targeting of prostate cancer. *Prostate* **2006**, 66, 1359–1370.
- (16) Wolf, P.; Gierschner, D.; Bühler, P.; Wetterauer, U.; Elsässer-Beile, U. A recombinant PSMA-specific single-chain immunotoxin has potent and selective toxicity against prostate cancer cells. *Cancer Immunol. Immunother.* **2006**, 55, 1367–1373.
- (17) Wolf, P.; Alt, K.; Bühler, P.; Katzenwadel, A.; Wetterauer, U.; Tacke, M.; Elsässer-Beile, U. Anti-PSMA immunotoxin as novel treatment for prostate cancer? High and specific antitumor activity on human prostate xenograft tumors in SCID mice. *Prostate* **2008**, 68, 129–138.
- (18) Bühler, P.; Wolf, P.; Gierschner, D.; Schaber, I.; Katzenwadel, A.; Schultze-Seemann, W.; Wetterauer, U.; Tacke, M.; Swamy, M.; Schamel, W. W.; Elsässer-Beile, U. A bispecific diabody directed against prostate-specific membrane antigen and CD3 induces T-cell mediated lysis of prostate cancer cells. *Cancer Immunol. Immunother.* **2008**, 57, 43–52.
- (19) Lupold, S. E.; Hicke, B. J.; Lin, Y.; Coffey, D. S. Identification and characterization of nuclease-stabilized RNA molecules that bind human prostate cancer cells via the prostate-specific membrane antigen. *Cancer Res.* **2002**, 62, 4029–4033.

prostate cancer.^{20–25} Aptamer-based macromolecules as cellular imaging and cellular labeling agents also have been investigated.^{26,27}

PSMA, presumably like other cell surface receptors, undergoes internalization constitutively upon binding to its putative ligand. This process is referred to as receptor-mediated endocytosis, in which more and more elaborate mechanisms have been recognized and characterized recently.²⁸ Nevertheless, how the PSMA directs the conjugates to be endocytosed into the tumor cells once it is bound to polymer carriers as a targeting moiety has not been investigated.

Pinocytosis, the fluid-phase endocytosis, has been recently recognized to occur in all cells by at least four basic mechanisms: clathrin-mediated endocytosis (CME),²⁹ caveolae-mediated endocytosis,³⁰ clathrin- and caveolae-independent endocytosis,^{28,31} and macropinocytosis.³² These pathways differ with regard to the nature of the cargo, the mechanism of vesicle formation, the regulation of cargo

entry, and especially, the ultimate intracellular destinations of cargo.^{28,33}

CME occurs constitutively in all mammalian cells²⁸ and continuously takes up essential nutrients, such as transferrin and low-density lipoprotein. Cargoes internalized by cells through CME are subsequently transported to early endosomes, from which some cargo is recycled back to the plasma membrane, whereas other consignments are delivered to late endosomes, and ultimately, lysosomes for degradation.²⁹ The caveolae-mediated endocytic pathway is less defined than CME. Caveolae is a noncoated flask shaped membrane invagination, which is formed by an assembly of glycosphingolipids and cholesterol (lipid raft) and the integrated transmembrane protein caveolin. Two well-studied ligands of this pathway are cholera toxin subunit B (CTX) and SV40 simian virus; CTX eventually targets to Golgi and SV40 to the endoplasmic reticulum.^{30,34} Macropinocytosis is traditionally considered to involve the internalization of large size macromolecules or significant amounts of fluid together with large areas of plasma membrane. However, it is actin-dependent and regulated by phosphoinositide (PI) 3-kinases.³⁵ The mechanisms that govern caveolae- and clathrin-independent endocytosis remain largely unknown. One of distinction that can be used to define this pathway is the independence of dynamin, a GTPase protein surrounding the neck of vesicle pits to mediate the scission of a wide range of vesicles such as clathrin-coated vesicles and caveolae.^{31,36}

The importance of understanding endocytic pathways involved in the internalization and subcellular trafficking of polymer–drug conjugates has been greatly appreciated recently.^{37,38} For example, one can design efficient endosomal escape or lysosomal enzymatic cleavage based on the trafficking and localization of polymer–drug conjugates. One also can optimize the physicochemical characteristics of polymer–drug conjugates to guide them to the proper membrane limited subcellular organelle. In the present study, the endocytic mechanisms and subcellular trafficking of

- (20) McNamara, J. O.; Andrechek, E. R.; Wang, Y.; Viles, K. D.; Rempel, R. E.; Gilboa, E.; Sullenger, B. A.; Giangrande, P. H. Cell type-specific delivery of siRNAs with aptamer-siRNA chimeras. *Nat. Biotechnol.* **2006**, *24*, 1005–1015.
- (21) Farokhzad, O. C.; Cheng, J.; Teply, B. A.; Sherifi, I.; Jon, S.; Kantoff, P. W.; Richie, J. P.; Langer, R. Targeted nanoparticle-aptamer bioconjugates for cancer chemotherapy in vivo. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 6315–6320.
- (22) Cheng, J.; Teply, B. A.; Sherifi, I.; Sung, J.; Luther, G.; Gu, F. X.; Levy-Nissenbaum, E.; Radovic-Moreno, A. F.; Langer, R.; Farokhzad, O. C. Formulation of functionalized PLGA-PEG nanoparticles for in vivo targeted drug delivery. *Biomaterials* **2007**, *28*, 869–876.
- (23) Dhar, S.; Gu, F. X.; Langer, R.; Farokhzad, O. C.; Lippard, S. J. Targeted delivery of cisplatin to prostate cancer cells by aptamer functionalized Pt(IV) prodrug-PLGA-PEG nanoparticles. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 17356–17361.
- (24) Wullner, U.; Neef, I.; Eller, A.; Kleines, M.; Tur, M. K.; Barth, S. Cell-specific induction of apoptosis by rationally designed bivalent aptamer-siRNA transcripts silencing eukaryotic elongation factor 2. *Curr. Cancer Drug Targets* **2008**, *8*, 554–565.
- (25) Chu, T. C.; Marks, J. W., 3rd; Lavery, L. A.; Faulkner, S.; Rosenblum, M. G.; Ellington, A. D.; Levy, M. Aptamer:toxin conjugates that specifically target prostate tumor cells. *Cancer Res.* **2006**, *66*, 5989–5992.
- (26) Farokhzad, O. C.; Khademhosseini, A.; Jon, S.; Hermmann, A.; Cheng, J.; Chin, C.; Kiselyuk, A.; Teply, B.; Eng, G.; Langer, R. Microfluidic system for studying the interaction of nanoparticles and microparticles with cells. *Anal. Chem.* **2005**, *77*, 5453–5459.
- (27) Bagalkot, V.; Zhang, L.; Levy-Nissenbaum, E.; Jon, S.; Kantoff, P. W.; Langer, R.; Farokhzad, O. C. Quantum dot-aptamer conjugates for synchronous cancer imaging, therapy, and sensing of drug delivery based on bi-fluorescence resonance energy transfer. *Nano Lett.* **2007**, *7*, 3065–3070.
- (28) Conner, S. D.; Schmid, S. L. Regulated portals of entry into the cell. *Nature (London)* **2003**, *422*, 37–44.
- (29) Lakadamyali, M.; Rust, M. J.; Zhuang, X. Ligands for clathrin-mediated endocytosis are differentially sorted into distinct populations of early endosomes. *Cell* **2006**, *124*, 997–1009.
- (30) Pelkmans, L. Secrets of caveolae- and lipid raft-mediated endocytosis revealed by mammalian viruses. *Biochim. Biophys. Acta* **2005**, *1746*, 295–304.

- (31) Kirkham, M.; Parton, R. G. Clathrin-independent endocytosis: new insights into caveolae and non-caveolar lipid raft carriers. *Biochim. Biophys. Acta* **2005**, *1746*, 349–363.
- (32) Swanson, J. A. Shaping cups into phagosomes and macropinosomes. *Nat. Rev. Mol. Cell Biol.* **2008**, *9*, 639–649.
- (33) Haucke, V. Cargo takes control of endocytosis. *Cell* **2006**, *127*, 35–37.
- (34) Perret, E.; Lakkaraju, A.; Deborde, S.; Schreiner, R.; Rodriguez-Boulan, E. Evolving endosomes: how many varieties and why. *Curr. Opin. Cell Biol.* **2005**, *17*, 423–434.
- (35) Lindmo, K.; Stenmark, H. Regulation of membrane traffic by phosphoinositide 3-kinases. *J. Cell Sci.* **2006**, *119*, 605–614.
- (36) Praefcke, G. J.; McMahon, H. T. The dynamin superfamily: universal membrane tubulation and fission molecules. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 133–147.
- (37) Bareford, L. M.; Swaan, P. W. Endocytic mechanisms for targeted drug delivery. *Adv. Drug Delivery Rev.* **2007**, *59*, 748–758.
- (38) Richardson, S. C.; Wallom, K. L.; Ferguson, E. L.; Deacon, S. P.; Davies, M. W.; Powell, A. J.; Piper, R. C.; Duncan, R. The use of fluorescence microscopy to define polymer localisation to the late endocytic compartments in cells that are targets for drug delivery. *J. Controlled Release* **2008**, *127*, 1–11.




Figure 1. Scheme of synthesis of HPMA copolymer–anti-PSMA antibody conjugates (P-anti-PSMA).

HPMA copolymer–anti-PSMA antibody conjugates were investigated in depth in live cells.

Experimental Section

Materials. Alexa Fluor 633 labeled transferrin, Alexa Fluor 647-labeled dextran 10 kDa and Hoechst 33342 were purchased from Molecular Probes (Carlsbad, CA). Chlorpromazine, filipin complex, mevinolin, 5-(*N*-ethyl-*N*-isopropyl)-amiloride, LY 294002, and wortmannin were from Sigma-Aldrich (St. Louis, MO). Clathrin HC siRNA and control siRNA were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Lipofectamine 2000 was from Invitrogen (Carlsbad, CA). The plasmid encoding EGFP-dynamin wild type (EGFP-dynamin^{WT}) and dynamin deficient mutant plasmid (EGFP-dynamin^{K44A}) were gifts from Dr. Diane McVey Ward (University of Utah). The plasmids encoding EYFP-Rab5 and EYFP-Rab7 were kindly provided by Dr. Xiaowei Zhuang (Harvard University). LNCap derived androgen independent C4-2 cells and the prostate cancer cell line PC-3 were purchased from American type Culture Collection (ATCC) (Manassas, VA). Fetal bovine serum was from Hyclone (Logan, UT). C4-2 and PC-3 cells were maintained in RPMI 1640 medium supplemented with 10% fetal bovine serum at 37 °C in an incubator with 5% CO₂. The sterile 35 mm Petri dish with 14 mm glass bottom microwell for confocal microscopy was from MatTek Corporation (Ashland, MA). Three monoclonal antibodies against different epitopes of PSMA (3A/12, 3F/11, 3E/7)

were generated as previously described.¹⁵ In the present study, all the data were generated with PSMA (3A/12) and its polymeric counterpart (P-anti-PSMA) except the determination of the antigen binding affinity.

Methods. UV–vis spectra were measured on a Varian Cary 400 Bio UV–visible spectrophotometer. Mass spectra of all synthesized compounds were obtained using a mass spectrometer Voyager-DE (STR Biospectrometry Workstation, PerSeptive Biosystems, Framingham, MA). The molecular weight and molecular weight distribution of polymers were measured on the ÅKTA FPLC system (GE Healthcare, formerly Amersham) equipped with UV and RI detectors using a Superose 6 HR10/30 column with PBS (pH 7.4) as the mobile phase. The average molecular weights were calculated using a calibration with polyHPMA fractions.

Synthesis of HPMA Copolymer–Antibody Conjugates. The conjugates were prepared in several steps (Figure 1). Polymeric precursors containing reactive thiazolidine-2-thione (TT) groups for binding antibody were synthesized by solution radical copolymerization of monomers in DMSO. The comonomers, HPMA (key monomer unit), MA-Gly-Gly-TT (*N*-methacryloylglycylglycine thiazolidine-2-thione; monomer unit bearing reactive TT group at the end of diglycine spacer³⁹) and MA-FITC ((5-[3-(methacryloylamino)propyl]thioureidyl]fluorescein; monomer unit containing

(39) Šubr, V.; Ulbrich, K. Synthesis and properties of new *N*-(2-hydroxypropyl)methacrylamide copolymers containing thiazolidine-2-thione. *React. Funct. Polym.* **2006**, *66*, 1525–1538.

FITC⁴⁰), were prepared as previously described. In the last step, the antibodies were coupled to the polymer precursors by aminolysis of TT groups in PBS.

Synthesis of Reactive Polymer Precursors. Polymer precursor containing TT and FITC (P-FITC-TT) was prepared by polymerization of HPMA (270 mg, 94.5 mol %), MA-GG-TT (31 mg, 5 mol %), MA-FITC (5.3 mg, 0.5 mol %) and 20 mg AIBN in 1.8 mL of DMSO containing 10 μ L of CH₃COOH. After purification by reprecipitation into acetone the polymer contained 0.32 mmol of TT/g (4.9 mol % TT; determined by UV spectrophotometry using extinction coefficient of monomer 10800 M⁻¹ cm⁻¹ in MeOH, 305 nm) and 0.027 mmol of FITC/g (0.4 mol %); yield 0.25 g; M_w 42 kDa, M_w/M_n 1.5.

Polymer precursor containing Texas Red (P-TxR-TT) was prepared from precursor containing only TT groups (prepared exactly as the P-FITC-TT omitting the MA-FITC in the polymerization mixture) by modifying a small amount of reactive TT groups by Texas Red cadaverin (Molecular Probes Texas RedC₅). Briefly, 100 mg of polymer precursor containing 0.038 mmol of TT groups was dissolved in 0.5 mL of DMSO, and 1.7 mg (0.0025 mmol) of Texas Red cadaverin was added and stirred for 30 min at room temperature (rt). The polymer was precipitated into acetone and dried. The content of TxR groups was 0.022 mmol/g (0.35 mol %; extinction coefficient 86000 M⁻¹ cm⁻¹ in PBS, 593 nm).

Binding of Anti-PSMA Ab to Polymer Precursors. A typical procedure was as follows: The polymer precursor (2 mg) was dissolved in 50 μ L of deionized H₂O and mixed with a cold solution of 2 mg of antibody in 0.5 mL of PBS, pH 7.4 while stirring. The mixture was stirred at 4 °C for 30 min. Then the pH was gradually raised up to 8.6 within 4 h at rt. The reaction mixture was left overnight in refrigerator. Next day the pH of reaction mixture was increased to 9 and the released TT was removed using a Sephadex G-25 (PD-10 column). The conjugate was fractionated on a Superose 6 (16/60) column (Amersham GE Healthcare) to separate a small amount (<5%) of high-molecular-weight fraction (eluted on the exclusion limit of the column, M_w > 500 kDa), the nonbound polymer and free antibody (Figure 2). The conjugate fraction was concentrated by ultrafiltration to 1.5–2.5 mL (Amicon membrane; cut off 30 kDa), and the final molecular weight profile was measured using Superose 6 (HR 10/30) column in PBS. The content of FITC (or TxR) containing polymer in concentrated conjugate fraction was measured spectrophotometrically, and the content of Ab was determined by the Lowry method (after subtraction of the background corresponding to polymer). The molecular weight of the conjugates was approximately 300 kDa as determined by online light-scattering detector (MiniDawn, Wyatt Technology Corp.).




Figure 2. Typical example of the SEC elution profile of P-anti-PSMA using Superose 6 HR16/60 column (AKTA/FPLC, buffer PBS).

Radioiodination of Antibody and Conjugates. Free antibody and copolymer antibody conjugates (P-anti-PSMA) were radioiodinated by the Iodogen method.^{41,42} Briefly, 300 μ L of antibody or copolymer antibody conjugates (0.5 mg/mL in PBS, pH 7.4) was added into an Iodogen precoated tube followed by addition of 10 μ L (0.5 mCi) of Na¹²⁵I. After 5 min the labeled antibody or P-anti-PSMA was purified using a PD-10 column (Amersham GE Healthcare) pre-equilibrated with PBS containing 0.5% BSA. The specific radioactivity was 1.5 μ Ci/ μ g.

Radioimmunoassay. This was performed to determine the antigen binding affinity of the free antibody and P-anti-PSMA. Briefly, cells grown in 96-wells at 80–90% confluence were incubated with Hanks balanced salt solution containing HEPES (20 mM), NaN₃ (10 mM) and BSA (0.5%) 1 h before experiments on ice. Subsequently, cells were incubated with the above buffer containing a serial concentration of labeled and unlabeled free antibody or P-anti-PSMA on ice for 4 h. After incubation, the buffer was discarded and cells were extensively washed to remove unbound antibody or P-anti-PSMA conjugates. Cells were solubilized with 1 M NaOH for 1 h and counted for radioactivity. The values of radioactivity were analyzed using the SigmaPlot software.

Transfection. One day before transfection, 1×10^5 cells were seeded in a 35 mm Petri dish with 14 mm glass bottom microwell in the growth media. For each transfection sample, 2 μ g of EGFP-dynamin^{WT}, EGFP-dynamin^{K44A}, EYFP-Rab5 or EYFP-Rab7 expression plasmids and 50 pmol of clathrin heavy chain siRNA or scrambled sequence (control) siRNA were diluted in 100 μ L of Opti-MEM I and mixed well; 5 μ L of Lipofectamine 2000 was diluted with 100 μ L of Opti-MEM I and kept for 5 min at room temperature. Then the two mixtures were combined and let to stand for 20 min at room temperature. Prior to exposure to transfection complex, cells were washed with Opti-MEM I twice and incubated with 800 μ L of Opti-MEM I. The 200 μ L mixtures were

(40) Omelyanenko, V.; Kopečková, P.; Gentry, C.; Kopeček, J. Targetable HPMA copolymer-adriamycin conjugates. Recognition, internalization, and subcellular fate. *J. Controlled Release* **1998**, 53, 25–37.

(41) Fraker, P. J.; Speck, J. C., Jr. Protein and cell membrane iodinations with a sparingly soluble chloroamide, 1,3,4,6-tetrachloro-3a,6a-diphenylglycoluril. *Biochem. Biophys. Res. Commun.* **1978**, 80, 849–857.

(42) Omelyanenko, V.; Kopečková, P.; Gentry, C.; Shiah, J. G.; Kopeček, J. HPMA copolymer-anticancer drug-OV-TL16 antibody conjugates. 1. Influence of the method of synthesis on the binding affinity to OVCAR-3 ovarian carcinoma cells in vitro. *J. Drug Targeting* **1996**, 3, 357–373.

then applied into dishes and incubated with cells for 5 h. After that, the transfection complex was discarded and cells were further incubated in growth media. Experiments were performed at 24–48 h post-transfection.

Internalization Experiments. Cells were plated into sterile 35 mm Petri dishes with 14 mm glass bottom microwells at a density of 1×10^5 cells per well. Cells were incubated for 2 days before internalization experiments were performed. P-anti-PSMA and Alexa Fluor 633-labeled transferrin were added to the culture medium and incubated with C4-2 cells at 37 °C for the indicated time periods. Alexa Fluor 647-labeled dextran 10 kDa was added to culture medium at a concentration of 50 $\mu\text{g/mL}$ and incubated with cells overnight followed by adding P-anti-PSMA. Ten minutes before microscopy, the DNA dye, Hoechst 33342, was added into dishes and incubated with cells. Then cells were washed with PBS twice and live-cell fluorescence imaging was performed immediately.

Inhibition of Internalization via Endocytic Pathways by Selective Inhibitors. Cells were incubated in serum free medium containing chlorpromazine 10 μM , wortmannin 1 μM , amiloride 10 μM , or LY294002 10 μM . Thirty minutes later, P-anti-PSMA was added into the medium (in the presence of these inhibitors) and cells were incubated for another 4 h. Experiments were performed immediately after incubation.

Cholesterol Depletion. To deplete cells of cholesterol, the cells were incubated in serum free medium for 1 h and then filipin 2.5 $\mu\text{g/mL}$ or mevinolin 10 μM was added and the cells were incubated for another 30 min. Experiments were carried out at the end of incubation.

Quantitative Study of Uptake by Flow Cytometry. The cells were seeded in 12-well plates at a density of 2.5×10^5 cells per well and incubated in growth medium for 2 days (37 °C, 5% CO_2). Then the cells were incubated with conjugates at 37 °C for indicated time periods. After incubation, medium containing conjugates was discarded. Cells were harvested and washed with cold PBS three times followed by flow cytometry analysis immediately. The amount of 1.0×10^4 cells was collected and analyzed for each sample.

Statistical Analysis. Statistical analysis was performed using Student's *t* test with $0.01 < *p < 0.05$ or $**p < 0.01$ as significant difference. The experiments were performed in triplicate. Cells treated with each individual inhibitor were compared with cells without exposure to inhibitors.

Results

Synthesis and Characterization of P-anti-PSMA Conjugates. The synthesis of polymer precursors and of HPMA copolymer–anti-PSMA antibody conjugates is shown in Figure 1. The polymer precursor P(FITC)-(GG-TT) contained 4.9 mol % of TT, 0.4 mol % of FITC. The M_w was 42 kDa and M_w/M_n 1.5. The polymer precursor P(TxR)-(GG-TT) contained 4.8 mol % of TT, 0.35 mol % of TxR. The M_w was 50 kDa, and M_w/M_n was 1.5.

Table 1. Characterization of P-anti-PSMA

conjugate	polymer precursor	antibody	conjugate composition ^a (wt %)	
			Ab	polymer
P(FITC)-3F11	P(FITC)-(GG-TT)	3F11	63.9	36.1
P(FITC)-3E7	P(FITC)-(GG-TT)	3E7	61.8	38.2
P(TxR)-3A12	P(TxR)-(GG-TT)	3A12	62.0	38.0
P(FITC)-IgG ^b	P(FITC)-(GG-TT)	IgG	66.5	33.5

^a Molecular ratio of Ab: polymer was calculated for all conjugates as ~1:3. ^b Human IgG.

The antibodies were covalently bound to HPMA copolymer precursors (P(FITC)-(GG-TT) and P(TxR)-(GG-TT)) via amide bonds formed by aminolysis of reactive thiazolidine-2-thione groups on the HPMA copolymer. This method involves the reaction of amino groups on the surface of antibody (mostly ϵ -amino groups of lysine). The intention was to modify the antibody only moderately to avoid conformation changes of the antibody molecule and prevent the decrease of its affinity to the target. Data from our previous study⁴² showed that the amino groups in the vicinity of binding site might be less reactive than in the other part of the antibody molecule. The K_d of the conjugate prepared by aminolysis was of the same order as the original antibody.⁴² The reaction conditions in this study were optimized to attach approximately three polymer chains per molecule of Ab. The weight ratio of Ab to polymer precursor was 1:1, and the concentration of Ab in the reaction mixture was 0.4 wt %. Such conditions generated only a small amount of high-molecular-weight (branched or cross-linked) fraction; it was removed by SEC fractionation. The molecular weight of the conjugates calculated from the chemical composition, approximately 300 kDa, was confirmed by SEC equipped with online laser light scattering detector; the estimated size was 10–12 nm.

The characteristics of conjugates are summarized in Table 1. A typical example of the size exclusion chromatography elution profile from fractionation of conjugates using Superose 6 HR16/60 column (AKTA/FPLC, Pharmacia column, buffer PBS) is shown in Figure 2.

Determination of the Antigen Binding Affinity of the Free Antibodies and Copolymer Antibody Conjugates. The PSMA molecule binding affinity of the anti-PSMA antibodies and P-anti-PSMA conjugates was determined by radioimmunoassay in C4-2 cells highly expressing PSMA molecules. The nonspecific binding of the antibody and copolymer antibody conjugates to cells was estimated in PC-3 cells that do not express PSMA. Three monoclonal antibodies against different epitopes of PSMA and their corresponding copolymer conjugates were examined, and the averages of dissociation constants (affinity) from three experiments are listed in Table 2. The binding affinities of all three antibodies were not compromised by conjugation to copolymer drug carriers. As expected, the affinity of antibodies attached to HPMA copolymer were moderately lower, but still in the same order of magnitude as the native Ab.

Table 2. The Dissociation Affinities of Free Anti-PSMA Antibodies and P-anti-PSMA^a

sample	3A/12	3F/11	3E/7	P-anti-PSMA(3A/12)	P-anti-PSMA(3F/11)	P-anti-PSMA(3E/7)
<i>K_d</i> (nM)	7.5	6.8	12	24	16	42

^a The dissociation affinities of three different anti-PSMA antibodies and their corresponding copolymer conjugates in C4-2 cells were measured, and the averages of dissociation affinities from three experiments were listed. The error of determination was approximately $\pm 25\%$.

The HPMA Copolymer Anti-PSMA Conjugates Are Endocytosed into PSMA Positive Cells through Receptor Mediated Endocytosis. It has been shown previously that the mAbs 3/A12, 3/F11 and 3/E7 bind to cell adherent PSMA and are internalized.¹⁵ To prove that P-anti-PSMA is also internalized upon the binding to PSMA, the P-anti-PSMA was incubated with C4-2 cells and PC-3 cells (Figure 3a). P-anti-PSMA was internalized into C4-2 cells whereas uptake of P-anti-PSMA was not observed in PC-3 cells, indicating that P-anti-PSMA was internalized into C4-2 cells through PSMA-mediated endocytosis. Dextran 10 kDa, a fluid phase endocytosis marker, was coincubated and endocytosed into PC-3 cells, demonstrating PC-3 cells' ability to execute fluid phase endocytosis, which subsequently confirmed the binding specificity of P-anti-PSMA to PSMA molecules.

The Rate of Uptake of P-anti-PSMA Is Faster Than That of HPMA Copolymer Conjugate with Nonspecific IgG. As a control, an HPMA copolymer conjugate with nonspecific IgG (P-IgG) was synthesized and the kinetics of P-anti-PSMA uptake was evaluated in C4-2 cells using flow cytometry (Figure 3b). The uptake of P-anti-PSMA could be detected after 5 min of incubation of P-anti-PSMA with cells whereas the uptake of P-IgG was seen only after 1 h of incubation. The uptake of P-anti-PSMA increased more rapidly with time of incubation than that of P-IgG. It implied that uptake of P-anti-PSMA is facilitated by receptor-mediated endocytosis.

Both CME and Macropinocytosis Participate in the Internalization of P-anti-PSMA by C4-2 Cells. Mechanisms of endocytosis that might be involved in the internalization of P-anti-PSMA were investigated utilizing a variety of commonly used pathway selective inhibitors, including chlorpromazine, an inhibitor of clathrin mediated endocytosis, filipin and mevinolin, cholesterol disruption agents, inhibitors of caveolae-mediated endocytosis and amiloride, wortmannin, and LY 294002, macropinocytosis inhibitors. Endocytosis of P-anti-PSMA was diminished in cells exposed to chlorpromazine, amiloride, wortmannin and LY 294002, observed through the confocal microscopic study (Figure 4a). The quantitative estimation of uptake of P-anti-PSMA was carried out by flow cytometry analysis (Figure 4b). The uptake of P-anti-PSMA was inhibited by nearly 40% in cells exposed to chlorpromazine at the indicated concentration. Suppressed uptake was displayed in cells treated with three inhibitors of macropinocytosis. In particular, wortmannin, an inhibitor of phosphoinositide (PI) 3-kinases that plays a role on the regulation of macropinocytosis, reduced uptake of P-anti-PSMA nearly up to 50% at the indicated concentration. Hence, the result revealed that more than one mechanism




Figure 3. (a) Internalization of P-anti-PSMA in PSMA positive C4-2 and PSMA negative PC-3 cells. Cells were preincubated with Alexa 647 labeled dextran 10 kDa, a fluid phase pinocytosis marker, for 16 h at 37 °C and then exposed to P-anti-PSMA for 2 h. [The concentration of P-anti-PSMA used in the whole study was 12×10^{-3} mg/mL of anti-PSMA and 7.4×10^{-3} mg/mL of copolymer.] Medium containing the conjugate was washed out followed by confocal microscopy study. [Images were taken in live cells in the whole study.] The pictures were from one confocal Z slice. [All pictures shown were from one confocal Z slice taken using an Olympus confocal microscope (FV 1000).] (b) Kinetic study of uptake of P-anti-PSMA by C4-2 cells. The cells were incubated with either P-anti-PSMA or HPMA copolymer–nonspecific IgG conjugate (P-IgG) using comparable amounts of anti-PSMA and IgG, as well as FITC, for the indicated period of time, followed by flow cytometry analysis.

are contributing to the internalization of P-anti-PSMA by C4-2 cells.

Down-Regulation of Clathrin Expression Diminishes the Internalization of P-anti-PSMA in C4-2 Cells. To further confirm that internalization of P-anti-PSMA was




Figure 4. Inhibition of P-anti-PSMA uptake using pathway selective inhibitors. Cells were pretreated with chlorpromazine (10 μ M), an inhibitor of clathrin-mediated endocytosis; filipin (2.5 μ g/mL) and mevinolin (10 μ M), inhibitors of caveolae-mediated endocytosis; amiloride (10 μ M), wortmannin (10 μ M), and LY 294002 (10 μ M), macropinocytosis inhibitors, for 0.5 h. The cells were incubated for another 2 h with P-anti-PSMA in the presence of inhibitors. (a) Confocal microscopy images of cells demonstrate diminished endocytosis. (b) Quantitative analysis of uptake by flow cytometry was carried out after incubation. The percentages of uptake of each sample in comparison to control are shown. The experiments were performed in triplicate, and asterisks (*) and (**) indicate statistically significant differences ($0.01 < p < 0.05$) and ($p < 0.01$) compared to control, respectively.

mediated by CME, the most well studied mechanism of receptor-mediated endocytosis, the influence of down-regulating the expression of clathrin protein, the main assembly unit of clathrin coated pits, on uptake of P-anti-PSMA was assessed. C4-2 cells were transfected with siRNA specific against clathrin heavy chain and protein expression was nearly completely abolished as detected by Western blotting (data not shown). Internalization of transferrin was

abolished in cells transfected with clathrin specific siRNA, as demonstrated by the disappearance of intracellular transferrin and retention of transferrin on plasma membrane. Similarly, the intracellular fluorescence of P-anti-PSMA decreased in cells transfected with clathrin specific siRNA, compared to control cells, indicating that internalization of P-anti-PSMA was regulated by clathrin (Figure 5a). In addition, C4-2 cells were incubated with P-anti-PSMA and




Figure 5. (a) Inhibition of the endocytosis of transferrin and P-anti-PSMA using clathrin-specific siRNA. C4-2 cells were transfected with clathrin heavy chain siRNA and incubated for 48 h, followed by incubation with transferrin and P-anti-PSMA for 1 h. (b) Colocalization of P-anti-PSMA with transferrin. P-anti-PSMA and Alexa Fluor 633 labeled transferrin were coincubated for 1 h. (c) Endocytosis of P-anti-PSMA was not completely abolished by dynamin deficiency. Cells were transfected with EGFP-dynamin^{WT} or EGFP-dynamin^{K44A}, a dynamin-deficient mutant that has been shown to block both clathrin- and caveolae-mediated endocytosis. After transfection, cells were incubated for 2 days and then exposed to P-anti-PSMA and transferrin.

Alexa Fluor 633 labeled transferrin and colocalization was analyzed. P-anti-PSMA was colocalized with transferrin (Figure 5b). Therefore, CME is the mechanism through which P-anti-PSMA is endocytosed into C4-2 cells.

Clathrin- and Caveolae-Independent Endocytosis Also Takes Part in P-anti-PSMA Internalization. The clathrin- and caveolae-independent pathway was also evaluated in the study. Because the clathrin- and caveolae-independent pathway is dynamin independent, a dominant negative (deficient) mutant of human dynamin (dynamin^{K44A}) has been widely used to characterize this pathway. In the present study, the wild type of dynamin (dynamin^{WT}) and dynamin^{K44A} were overexpressed as EGFP fusion protein in C4-2 cells to

determine the participation of clathrin- and caveolae-independent pathway in the internalization of P-anti-PSMA. Consequently, a P-anti-PSMA conjugate labeled with TxR was synthesized and used in this study. The internalization of P-anti-PSMA was examined in overexpression-positive cells. In cells overexpressing EGFP-dynamin^{WT} (highlighted with single arrow in Figure 5c), all the P-anti-PSMA was internalized and accumulated in the perinuclear area. However, in cells overexpressing EGFP-dynamin^{K44A} (highlighted with double arrow in Figure 5c), retention of P-anti-PSMA at the plasma membrane coexisted with perinuclear P-anti-PSMA, indicating that endocytosis of P-anti-PSMA was not completely blocked. The intracellular P-anti-PSMA bypassed dynamin and was endocytosed. Association of P-anti-PSMA with plasma membrane indicated that the endocytosis of some of P-anti-PSMA is regulated by dynamin.

P-anti-PSMA Antibody Conjugates Are Transported to Late Endosomes via Endosomes. Subcellular trafficking of P-anti-PSMA containing membrane vesicles was subsequently inspected. Rab small GTPases, members of the Ras superfamily, are localized at distinct membrane vesicles and are responsible for membrane vesicle formation, development and trafficking.^{43–46} It is well accepted that Rab5 and Rab7 are distinct domains of early endosomes and late endosomes, respectively.^{43–47} We expressed Rab5 and Rab7, fused to yellow fluorescent proteins (YFP). C4-2 cells were transfected with DNA constructs of YFP fused Rab5 and Rab7, and subsequently the trafficking of P-anti-PSMA in live cells was tracked. The P-anti-PSMA labeled with TxR was used in this study. In Figure 6, green signals represent either Rab5 expressing early endosomes or Rab7 expressing late endosomes, while the red signal signifies membrane vesicles entrapping TxR labeled P-anti-PSMA conjugates. Membrane limited vesicles containing P-anti-PSMA were observed to colocalize with Rab5 expressing vesicles as quickly as within 15 min, demonstrating that membrane vesicles containing P-anti-PSMA rapidly mature into early endosomes. However, Rab5 gradually disappeared from vesicles containing P-anti-PSMA conjugates within 1 h. The membrane limited vesicles containing P-anti-PSMA began to mature into late endosomes after incubation for 22 min. The maturation was seen to consistently increase up to 240 min.

Discussion

Understanding the mechanism of internalization and subcellular trafficking of targeted polymer drug conjugates is

- (43) Pfeffer, S. A model for Rab GTPase localization. *Biochem. Soc. Trans.* **2005**, *33*, 627–630.
- (44) Zerial, M.; McBride, H. Rab proteins as membrane organizers. *Nat. Rev. Mol. Cell Biol.* **2001**, *2*, 107–117.
- (45) Pfeffer, S.; Aivazian, D. Targeting Rab GTPases to distinct membrane compartments. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 886–896.
- (46) Pfeffer, S. Filling the Rab GAP. *Nat. Cell Biol.* **2005**, *7*, 856–857.
- (47) Rink, J.; Ghigo, E.; Kalaidzidis, Y.; Zerial, M. Rab conversion as a mechanism of progression from early to late endosomes. *Cell* **2005**, *122*, 735–749.




Figure 6. Subcellular trafficking of P-anti-PSMA conjugates. C4-2 cells were transfected with DNA constructs of YFP fused Rab5 and Rab7 by using lipofectamine 2000 according to manufacturer's recommendation. After 24 h, cells were incubated with P-anti-PSMA for indicated period of time. Membrane vesicles containing P-anti-PSMA were observed quickly to fuse to endosomes but rapidly disappeared from early endosomes. P-anti-PSMA containing vesicles began to fuse to late endosomes after incubation for 22 min. The fusion was seen to consistently increase up to 240 min.

of great importance for the explanation of their treatment effects and further optimization of design and synthesis. Recently, rapid advancements have been made and provided new insights into the complexity of physiological endocytic pathways. Different pathways can share a common basic pathway to which additional levels of sophistication are added.^{48,49} In addition, the same cargo can be internalized through distinct endocytic mechanisms. For instance, transforming growth factor- β receptor (TGF β R) is internalized through clathrin-mediated and caveolae-mediated endocytosis.⁵⁰

Not surprisingly, studies have shown that multiple endocytic pathways can concurrently take action when a

macromolecular drug conjugate is exposed to a single type of cell. It is because of the fact that macromolecular drug conjugates hijack the complicated physiological endocytic machinery. Rejman et al. demonstrated, using pathway inhibitors, that polyethyleneimine (PEI)/DNA polyplexes are internalized into cells through both CME and caveolae-mediated endocytosis.^{51–53} It was confirmed also by using colocalization experiments.^{54,55} The cellular uptake of pul-lulan-spermine, a cationic polysaccharide, occurred via CME

- (48) Bonifacino, J. S.; Rojas, R. Retrograde transport from endosomes to the trans-Golgi network. *Nat. Rev. Mol. Cell Biol.* **2006**, *7*, 568–579.
- (49) van der Goot, F. G.; Gruenberg, J. Intra-endosomal membrane traffic. *Trends Cell Biol.* **2006**, *16*, 514–521.
- (50) Le Roy, C.; Wrana, J. L. Clathrin- and non-clathrin-mediated endocytic regulation of cell signalling. *Nat. Rev. Mol. Cell Biol.* **2005**, *6*, 112–126.

- (51) Rejman, J.; Bragonzi, A.; Conese, M. Role of clathrin- and caveolae-mediated endocytosis in gene transfer mediated by lipo- and polyplexes. *Mol. Ther.* **2005**, *12*, 468–474.
- (52) Rejman, J.; Conese, M.; Hoekstra, D. Gene transfer by means of lipo- and polyplexes: role of clathrin and caveolae-mediated endocytosis. *J. Liposome Res.* **2006**, *16*, 237–247.
- (53) Hoekstra, D.; Rejman, J.; Wasungu, L.; Shi, F.; Zuhorn, I. Gene delivery by cationic lipids: in and out of an endosome. *Biochem. Soc. Trans.* **2007**, *35*, 68–71.
- (54) von Gersdorff, K.; Sanders, N. N.; Vandenbroucke, R.; De Smedt, S. C.; Wagner, E.; Ogris, M. The internalization route resulting in successful gene expression depends on both cell line and polyethyleneimine polyplex type. *Mol. Ther.* **2006**, *14*, 745–753.

and caveolae-mediated endocytosis, as proved by using appropriate inhibitors.⁵⁶ Most recently, a comprehensive model suggested that the Tat peptide simultaneously used three endocytic pathways: macropinocytosis, CME, and caveolae-mediated endocytosis.⁵⁷ In addition, a novel arginine-grafted dendritic block copolymer, R-PAMAM-PEG-PAMAM-R G5 (PPP5-R), was shown to simultaneously employ the same three pathways.⁵⁸

This study revealed that P-anti-PSMA is internalized by C4-2 cells through multiple endocytic pathways. CME is a major pathway of C4-2 cells internalizing P-anti-PSMA. Clathrin-coated pits are formed by the assembly of cytosolic coat proteins, the main assembly unit being clathrin. Clathrin is a three-legged structure formed by three heavy chains, each with a tightly associated clathrin light chain. We disrupted the formation of clathrin-coated pits by siRNA-mediated down-regulation of clathrin heavy chain protein expression. This genetic approach together with colocalization of P-anti-PSMA with transferrin and internalization in the presence of chlorpromazine inhibitor strongly indicated that CME is one of the mechanisms by which P-anti-PSMA molecules are internalized. This conclusion is indirectly supported by the biochemical characteristics of PSMA. Transferrin is the best-characterized ligand of CME. The entire sequence of PSMA is related to the transferrin receptor (TfR) with 20% homology. The overall structure and domain organization of TfR and PSMA are similar, as revealed by the crystal structure of PSMA.⁵⁹ It was proposed that small cytoplasmic motifs of TfR1 and PSMA, YXRF and MXXXL, respectively, solely regulate their uptake by endocytosis.⁶⁰

In addition to CME, a fraction of P-anti-PSMA was internalized into C4-2 cells through macropinocytosis.

Distinct from CME, macropinocytosis is initiated by plasma membrane protrusion to the external milieu, a process of extensive plasma membrane reorganization. Macropinocytosis occurs spontaneously or upon the stimulation of growth-factor-receptor.^{32,61,62} Presumably, a large volume of extracellular fluid or a macromolecule of large size triggers macropinocytosis. However, no regulation has been found to control the size or morphology of the enclosed macropinosome. Recently, involvement of actin filaments in macropinocytosis and regulation of macropinocytosis by PI3-kinases have been studied extensively.^{32,35} Inhibition of PI3K abolishes macropinocytosis of macromolecules (mostly in closing of cup invagination) and fusion of macropinosomes. It also interrupts the membrane protrusion in some cells. The physicochemical characteristics of cargo that is endocytosed by macropinocytosis remain unknown. However, as a common mechanism of fluid phase endocytosis, macropinocytosis has recently been discovered to actively participate in the internalization of a variety of macromolecules, such as intact protein, cell penetrating peptides, synthetic polymers and nanoparticles.^{62–66} Using three specific inhibitors of macropinocytosis, we verified that macropinocytosis takes part in internalization of P-anti-PSMA.

Clathrin- and caveolae-independent endocytosis was another mechanism found participating in the internalization of P-anti-PSMA. This endocytic pathway is least well-defined. Due to the independency of clathrin and caveolin, this pathway was defined by the absence of dynamin recruitment. In fact, the evidence of the existence of this pathway originated from the analysis of cells expressing a temperature-sensitive dynamin mutant. Dynamin is a multidomain GTPase that assembles into a “collar” around the neck of vesicle invagination to mediate the release of vesicles

- (55) van der Aa, M. A.; Huth, U. S.; Häfele, S. Y.; Schubert, R.; Oosting, R. S.; Mastrobattista, E.; Hennink, W. E.; Peschka-Süss, R.; Koning, G. A.; Crommelin, D. J. Cellular uptake of cationic polymer-DNA complexes via caveolae plays a pivotal role in gene transfection in COS-7 cells. *Pharm. Res.* **2007**, *24*, 1590–1598.
- (56) Kanatani, I.; Ikai, T.; Okazaki, A.; Jo, J.; Yamamoto, M.; Imamura, M.; Kanematsu, A.; Yamamoto, S.; Ito, N.; Ogawa, O.; Tabata, Y. Efficient gene transfer by pullulan-spermine occurs through both clathrin- and raft/caveolae-dependent mechanisms. *J. Controlled Release* **2006**, *116*, 75–82.
- (57) Duchardt, F.; Fotin-Mleczek, M.; Schwarz, H.; Fischer, R.; Brock, R. A comprehensive model for the cellular uptake of cationic cell-penetrating peptides. *Traffic* **2007**, *8*, 848–866.
- (58) Kim, T. I.; Baek, J. U.; Yoon, J. K.; Choi, J. S.; Kim, K.; Park, J. S. Synthesis and characterization of a novel arginine-grafted dendritic block copolymer for gene delivery and study of its cellular uptake pathway leading to transfection. *Bioconjugate Chem.* **2007**, *18*, 309–317.
- (59) Davis, M. I.; Bennett, M. J.; Thomas, L. M.; Bjorkman, P. J. Crystal structure of prostate-specific membrane antigen, a tumor marker and peptidase. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 5981–5986.
- (60) Rajasekaran, S. A.; Anilkumar, G.; Oshima, E.; Bowie, J. U.; Liu, H.; Heston, W.; Bander, N. H.; Rajasekaran, A. K. A novel cytoplasmic tail MXXXL motif mediates the internalization of prostate-specific membrane antigen. *Mol. Biol. Cell* **2003**, *14*, 4835–4845.

- (61) Amyere, M.; Mettlen, M.; Van Der Smissen, P.; Platek, A.; Payastre, B.; Veithen, A.; Courtoy, P. J. Origin, originality, functions, subversions and molecular signalling of macropinocytosis. *Int. J. Med. Microbiol.* **2002**, *291*, 487–494.
- (62) Jones, A. T. Macropinocytosis: searching for an endocytic identity and role in the uptake of cell penetrating peptides. *J. Cell Mol. Med.* **2007**, *11*, 670–684.
- (63) Mennesson, E.; Erbacher, P.; Piller, V.; Kieda, C.; Midoux, P.; Pichon, C. Transfection efficiency and uptake process of polyplexes in human lung endothelial cells: a comparative study in non-polarized and polarized cells. *J. Gene Med.* **2005**, *7*, 729–738.
- (64) Gratton, S. E.; Napier, M. E.; Ropp, P. A.; Tian, S.; DeSimone, J. M. Microfabricated particles for engineered drug therapies: elucidation into the mechanisms of cellular internalization of PRINT particles. *Pharm. Res.* **2008**, *25*, 2845–2852.
- (65) Kaplan, I. M.; Wadia, J. S.; Dowdy, S. F. Cationic TAT peptide transduction domain enters cells by macropinocytosis. *J. Controlled Release* **2005**, *102*, 247–253.
- (66) Magzoub, M.; Sandgren, S.; Lundberg, P.; Oglecka, K.; Lilja, J.; Witttrup, A.; Goran Eriksson, L. E.; Langel, U.; Belting, M.; Graslund, A. N-terminal peptides from unprocessed prion proteins enter cells by macropinocytosis. *Biochem. Biophys. Res. Commun.* **2006**, *348*, 379–385.

into the cytoplasm. It is involved in the scission of a wide range of vesicles such as clathrin-coated vesicles and caveolae.³⁶

Most recently, there was a report that dextran 10 kDa also adopts clathrin- and caveolae-independent pathway to be endocytosed into cells.³¹ We thus inspected the similarity between dextran and P-anti-PSMA internalization. In EGFP-dynamin^{K44A} positive cells, the internalization of dextran 10 kDa did not remarkably decline, similarly to P-anti-PSMA (data not shown). This supports the conclusion that a portion of P-anti-PSMA internalization is dynamin-independent, similarly to dextran 10 kDa, and occurs via the clathrin- and caveolae-independent pathway.

The clathrin-coated-pit formation and regulation, as well as trafficking of membrane vesicles derived from CME, are relatively well understood. Clathrin-coated pits enclosing cargoes pinching off from plasma membranes are subsequently transported to early endosomes. Sorting occurs at this stage. Some lipid and protein cargoes that possess housekeeping features are recycled back to the plasma membrane through recycling endosomes or the trans-Golgi network. Other molecules, destined to be degraded, congregate rapidly within multivesicular endosomes, which move along the microtubule network to late endosomes. Late endosomes act as another level sorting station from which most of the molecules presumably are transported to lysosomes for degradation.^{48,49}

The route of macropinosome transport in the endocytic pathway remains unclear. There is a tendency to agree that macropinosomes rapidly develop into early endosomes.^{67–69} Nevertheless, there is a discrepancy whether macropinosomes or derived early endosomes continue to progress into late endosomes and lysosomes.^{67–69} Similarly, the membrane vesicles enclosing cargoes from clathrin-, caveolae-independent entry might fuse with endosomes after pinching off. However, the final destination of cargo remains unsolved.

Trafficking of P-anti-PSMA is anticipated to be sophisticated based on the complexity of its entry into C4-2 cells. Hence, we decided to dissect and pick up the mainframe of P-anti-PSMA trafficking that is of importance for the design and optimization of macromolecular drug conjugates. We visualized the early endosomes and late endosomes by overexpression of Rab GTPase proteins. It was able to provide more conclusive results compared to commonly used lysotracker due to the high specificity. We found that some of membrane vesicles trapping P-anti-PSMA rapidly developed or fused to endosomes after budding off. Some of membrane vesicle trapping P-anti-PSMA rapidly fused to late endosomes, and it seemed that they consistently remained in Rab7 positive membrane vesicles for a prolonged time (24 h). Presumably, these Rab7 positive membrane vesicles ultimately mature into lysosomes. Interestingly, we found that in a specific period of time P-anti-PSMA disappeared from early endosomes, while it accumulated in late endosomes. It is well in agreement with the report from Zerial and co-workers that Rab5 is converted to Rab7 during early to late endosome trafficking.³⁸ Based on these results, polymer–drug conjugates can be designed containing drugs attached via either endosome escaping spacers or lysosome enzyme cleavable spacers for the best of treatment effects.

Overall, our results demonstrated that multiple mechanisms could be involved in the endocytosis of antibody–polymer conjugates. Trafficking of membrane vesicles trapping targeted macromolecular drug conjugates can be traced, which will guide the design and optimization of targeted macromolecular drug conjugates.

Abbreviations Used

PSMA, prostate specific membrane antigen; HPMA, *N*-(2-hydroxypropyl)methacrylamide; P-anti-PSMA, HPMA (*N*-(2-hydroxypropyl)methacrylamide) copolymer–anti-PSMA antibody conjugates; P-IgG, HPMA copolymer conjugate with nonspecific IgG; CME, clathrin-mediated endocytosis; dynamin K44A, dominant-negative mutant of dynamin.

Acknowledgment. We thank Dr. Diane McVey Ward (University of Utah) and Dr. Xiaowei Zhuang (Howard Hughes Medical Institute, Harvard University) for their generous gifts of EGFP-dynamin^{WT}, EGFP-dynamin^{K44A}, EYFP-Rab5 and EYFP-Rab7. The research was supported in part by NIH Grant RO1 CA132831 from the National Cancer Institute (to J.K.).

MP8002682

- (67) Kerr, M. C.; Lindsay, M. R.; Luetterforst, R.; Hamilton, N.; Simpson, F.; Parton, R. G.; Gleeson, P. A.; Teasdale, R. D. Visualisation of macropinosome maturation by the recruitment of sorting nexins. *J. Cell Sci.* **2006**, *119*, 3967–3980.
- (68) Schnatwinkel, C.; Christoforidis, S.; Lindsay, M. R.; Uttenweiler-Joseph, S.; Wilm, M.; Parton, R. G.; Zerial, M. The Rab5 effector Rabankyrin-5 regulates and coordinates different endocytic mechanisms. *PLoS Biol.* **2004**, *2*, E261.
- (69) Hamasaki, M.; Araki, N.; Hatae, T. Association of early endosomal autoantigen 1 with macropinocytosis in EGF-stimulated A431 cells. *Anat. Rec.* **2004**, *277*, 298–306.