

Confocal Images of Circulating Tumor Cells Obtained Using a Methodology and Technology That Removes Normal Cells

Priya Balasubramanian,[†] Liying Yang,[†] James C. Lang,[‡] Kris R. Jatana,[‡]
David Schuller,[‡] Amit Agrawal,[‡] Maciej Zborowski,[§] and Jeffrey J. Chalmers^{*,†,||}

William G. Lowrie Department of Chemical and Biomolecular Engineering, The Ohio State University, 125 Koffolt Laboratories, 140 West 19th Avenue, Columbus, Ohio 43210, College of Medicine and Public Health, Department of Otolaryngology—Head and Neck Surgery, The Ohio State University, Columbus Ohio 43210, and Department of Biomedical Engineering, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, Ohio

Received February 23, 2009; Revised Manuscript Received May 8, 2009; Accepted May 13, 2009

Abstract: A completely negative enrichment technology was used to detect circulating tumor cells, CTCs, in the peripheral blood of head and neck cancer patients. Of 32 blood samples, 63% contained CTCs and the number of CTCs identified per mL of blood collected ranged from 0 to 214. The final purity ranged from 1 CTC in 9 total cells to 1 CTC in 20,000 total cells, the final purity being both a function of the number of CTCs and the performance of the specific enrichment. Consistent with previous reports, CTC were positively identified if: (1) they contained a nucleus based on DAPI stain, (2) stained positive for cytokeratins, and (3) have a high nuclei to cytoplasmic ratio. In addition, for a blood sample to be considered positive for CTCs, the enriched sample must be positive for epithelial growth factor receptor, EGFR, as measured by RT-PCR. While most of the blood samples were obtained during surgery, a number were taken prior to and during surgery. In all of the pre- and postsurgery paired samples, significant numbers of CTCs were detected. A number of these enriched samples were observed under confocal microscope in addition to the microscopic observations under traditional wide-field fluorescent microscope. As expected, the FITC stained cytokeratins appeared in the cytoplasm and the average size of these positively stained cells, on the cytospin, was in the range of 8–12 μm . Future studies will involve the investigation if cancer stem cell and mesenchymal markers are present on these CTCs and correlations of patient outcome to the number and type of CTC present.

Keywords: Circulating tumor cells; immunomagnetic cell separation; immunocytochemistry; RT-PCR; head and neck squamous cell carcinoma (HNSCC)

Introduction

The genesis of overt metastases in a number of different types of cancer, including head and neck squamous cell

carcinoma, HNSCC, is based on the concept that tumor cells that dissociate from the primary cancer obtain access to circulation either directly through blood vessels or after transit in lymphatic channels. According to the classical view of metastatic spread, the dissemination of cancer cells into the circulation occurs during the later stages of progression toward malignancy. However, recent research in breast cancer has demonstrated that this process in fact occurs early during tumor development. Thus, detection of such cells in patients with newly diagnosed solid tumors has been considered an appealing strategy to provide evidence of future metastasis.^{1–3} Overall, the existence of circulating tumor cells (CTCs) and the settlement of these cells in secondary organs, such as liver, bone and lungs, as dis-

* Corresponding author. Mailing address: The Ohio State University, Dept of Chemistry & Biomolecular Engineering, 140 W 19th Ave, Columbus, Ohio 43210. Tel: (216) 292-2727. Fax: (216) 292-3769. E-mail: chalmers.1@osu.edu.

[†] William G. Lowrie Department of Chemical and Biomolecular Engineering, The Ohio State University.

[‡] Department of Otolaryngology—Head and Neck Surgery, The Ohio State University.

[§] Cleveland Clinic. Tel: (216) 445-9330. Fax: (216) 444-9198. E-mail: zborow@bme.ri.ccf.org.

^{||} Director, University Cell Analysis and Sorting Core.

seminated metastatic tumor cells (DTCs) is generally accepted.² Genome and transcriptome analyses of single disseminated tumor cells demonstrated that the majority of DTCs are cells with genetic aberrations compatible with malignancy. These cells are most likely direct descendants of the primary tumor, although the genetic changes can be incongruent with the dominant genotype of the corresponding primary tumor.^{4–6}

The detection of the presence of tumor cells outside the primary tumor would serve three purposes that could be, potentially, clinically useful: (1) as unambiguous evidence for an early occult spread of tumor cells; (2) as a relevant risk factor for subsequent metastasis and, thus, poor prognosis; and (3) as a marker for monitoring treatment susceptibility. From a research perspective, genotyping and phenotyping of CTCs can provide a detailed insight into the metastatic process and permit direct exploration of targeted treatment strategies.^{2,3}

The available literature regarding the prognostic relevance of the presence of DTC in bone marrow was inconsistent for breast cancer until a recent pooled analysis.⁷ The results of this large study with over 4000 patients showed that the presence of DTCs in bone marrow of patients with stage I–III breast cancer is a strong, adverse prognostic factor for all the clinically relevant end-points. The patient sample included disease-free, distant disease-free, breast cancer specific and overall survival in the entire cohort. It also included subsets of patients treated with chemotherapy only, endocrine therapy only, and those patients with small primary tumors and no lymph node metastases who received no systemic adjuvant therapy. A different study with metastatic breast cancer patients indicated that the detection of greater than 5 CTCs per 7.5 mL of peripheral blood adversely affects progression-free and overall survival and is a predictive marker for lack of response to treatment.⁸ This correlation was developed through studies performed in the first FDA approved diagnostic device to detect CTC: the CellSearch

system, currently sold by Veridex, LLC. This system is a semiautomated immunomagnetic enrichment system that involves positive selection of CTCs using anti-EpCAM antibodies coated with ferrofluids. The enriched CTCs are then labeled for immunofluorescent imaging with DAPI, anti-CD45 antibodies to label blood cells and anticytokeratin antibodies to label CTCs. The CellSearch system has been used in a number of other studies, including a much larger study where 70% of over 1400 metastatic breast cancer patients had at least 5 CTCs per 7.5 mL of blood.⁹

Several studies suggest that CTCs are descendants of the corresponding primary tumor.^{10–13} In contrast, it is thought that DTCs do not share such broad molecular similarities; therefore, it remains unanswered if a clinically desirable blood test for CTC detection would be able to provide the same information as the bone marrow test for DTC detection.

From a clinical point of view it will be necessary to determine what percentage of CTCs and DTCs are detectable, but nonviable, and which of them are actually capable of forming solid metastases. Several studies show the ability to profile DTCs and to differentiate their biological properties using anticytokeratin antibodies in combination with antibodies against tumor-associated markers.^{14–19} Differential gene or protein expression studies show that CTCs have malignant potential.^{10,13} In addition to tumor cell profiles, information on markers such as epidermal growth factor receptor

- (1) Fehm, T. Micrometastatic spread in breast cancer: detection, molecular characterization and clinical relevance. *Breast Cancer Res* **2008**, *10* (Suppl. 1), S1.
- (2) Pantel, K.; Brakenhoff, R. H.; Brandt, B. Detection, clinical relevance and specific biological properties of disseminating tumour cells. *Nat. Rev. Cancer* **2008**, *8* (5), 329–40.
- (3) Pantel, K.; Brakenhoff, R. H. Dissecting the metastatic cascade. *Nat. Rev. Cancer* **2004**, *4* (6), 448–56.
- (4) Schmidt-Kittler, O. From latent disseminated cells to overt metastasis: genetic analysis of systemic breast cancer progression. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100* (13), 7737–42.
- (5) Klein, C. A. Genetic heterogeneity of single disseminated tumour cells in minimal residual cancer. *Lancet* **2002**, *360* (9334), 683–9.
- (6) Klein, C. A. Combined transcriptome and genome analysis of single micrometastatic cells. *Nat. Biotechnol.* **2002**, *20* (4), 387–92.
- (7) Braun, S. A pooled analysis of bone marrow micrometastasis in breast cancer. *N. Engl. J. Med.* **2005**, *353* (8), 793–802.
- (8) Cristofanilli, M. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N. Engl. J. Med.* **2004**, *351* (8), 781–91.

- (9) Riethdorf, S. Detection of circulating tumor cells in peripheral blood of patients with metastatic breast cancer: a validation study of the CellSearch system. *Clin. Cancer Res.* **2007**, *13* (3), 920–8.
- (10) Fehm, T. Cytogenetic evidence that circulating epithelial cells in patients with carcinoma are malignant. *Clin. Cancer Res.* **2002**, *8* (7), 2073–84.
- (11) Hayes, D. F. Monitoring expression of HER-2 on circulating epithelial cells in patients with advanced breast cancer. *Int. J. Oncol.* **2002**, *21* (5), 1111–7.
- (12) Racila, E. Detection and characterization of carcinoma cells in the blood. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95* (8), 4589–94.
- (13) Meng, S. HER-2 gene amplification can be acquired as breast cancer progresses. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101* (25), 9393–8.
- (14) Reimers, N. Expression of extracellular matrix metalloproteases inducer on micrometastatic and primary mammary carcinoma cells. *Clin. Cancer Res.* **2004**, *10* (10), 3422–8.
- (15) Hensen, A. Comparative evaluation of urokinase-type plasminogen activator receptor expression in primary breast carcinomas and on metastatic tumor cells. *Int. J. Cancer* **2003**, *107* (6), 903–9.
- (16) Braun, S. ErbB2 overexpression on occult metastatic cells in bone marrow predicts poor clinical outcome of stage I–III breast cancer patients. *Cancer Res.* **2001**, *61* (5), 1890–5.
- (17) Braun, S. Tumor-antigen heterogeneity of disseminated breast cancer cells: implications for immunotherapy of minimal residual disease. *Int. J. Cancer* **1999**, *84* (1), 1–5.
- (18) Pantel, K. Differential expression of proliferation-associated molecules in individual micrometastatic carcinoma cells. *J. Natl. Cancer Inst.* **1993**, *85* (17), 1419–24.
- (19) Pantel, K. Frequent down-regulation of major histocompatibility class I antigen expression on individual micrometastatic carcinoma cells. *Cancer Res.* **1991**, *51* (17), 4712–5.

(EGFR),²⁰ epithelial cell adhesion molecule (EpCAM),¹⁷ uPA receptor (uPAR),¹⁵ and extracellular matrix metallo-proteinase inducer (EMMPRIN)¹⁴ may have therapeutic implications for targeted therapy strategies. Consequently, more efficient and cost-effective methods to enrich CTCs and DTCs could potentially expedite and support further important studies on tumor cell characterization and by doing so, potentially improve the prognostic and predictive value of CTCs and DTCs.

In contrast to breast cancer, few studies on CTCs in HNSCC have been conducted. One of these studies, by Partridge et al. (2003),²¹ reported that the presence of CTC in samples of bone marrow and central venous blood, collected preoperatively or postoperatively from head and neck cancer patients, indicated a high risk of local and distant recurrence and reduced survival.

CTC Detection Technology. Several methodologies have been developed to detect and characterize DTCs in bone marrow or CTCs in peripheral blood of cancer patients. These methodologies fall under two main categories: Immunological based assays and molecular based assays. Although these approaches can be used directly, an enrichment step prior to the detection is preferred. A number of methodologies exist for enriching rare cancer cells, including density gradient separation to enrich for nucleated cells (i.e., Ficoll), and magnetic cell separation either targeting the cancer cell, positive immunomagnetic cell separation, PIMS, or negative immunomagnetic cell separation, NIMS, where normal blood cells are targeted. Unfortunately, most of the reported studies using some form of magnetic cell separation to separate or enrich rare cancer cells do not provide data, or complete data, on the performance of the magnetic separation step. Consequently, it makes comparison of the performance of technology, as well as the overall study, difficult or impossible.

Recently, two new approaches to enrich for CTCs have been reported. In one of these approaches, a microfluidic device containing anti-EpCAM antibodies conjugated to “microposts” was developed to facilitate interactions between CTCs and the posts, when whole blood flows through the device under controlled laminar flow conditions. It is reported to have high specificity and sensitivity; however, since it is another version of positive selection, it suffers from the bias of assuming that all CTCs express the epithelial surface marker EpCAM.²² The other method involves enrichment of CTCs by filtration before genomic analysis. Basically, this approach assumes that the CTCs

are bigger than typical blood cells; therefore, the normal cells will flow through the filtration media while the CTCs are retained.²³ As with other positive selection approaches, this has an intrinsic bias that CTCs are different in size than normal blood cells.

In contrast, the Chalmers and Zborowski laboratories have developed a purely negative enrichment technology in which only normal blood cells are targeted and removed, thereby allowing rare, non-hematopoietic cells to be significantly enriched (Yang et al. 2009).²⁴ They reported that this system is able to reduce the number of normal blood cells in a cancer patient's blood from 4.05×10^9 to 8.04×10^3 cells per mL and still recover, on average, 2.32 CTCs per mL of blood. For all of the cancer patient blood samples tested at the time of publication, and in which CTCs were detected (20 out of 26 patients) the average recovery of CTCs was 21.7 per mL of blood, with a range of 282 to 0.53 CTC.

Materials and Methods

Blood Collection. Ten to 18.5 mL of peripheral blood was obtained from patients who presented with, and underwent surgical resection for, HNSCC and have not been previously treated for this disease. Blood samples were collected in green-top BD-Vacutainer (Cat. #367874, BD Biosciences) and stored at 4 °C until processing. Samples were processed within 24 h after procurement. Operators were blinded to clinical correlative information during the cell suspension processing and analysis.

Cell Culture. The head and neck cancer cell line SCC-4 was purchased from ATCC (Manassas, VA) and maintained in Dulbecco's modified Eagle medium (DMEM) (Cat. #30-2002, ATCC, Manassas, VA) supplemented with 10% fetal bovine serum (FBS) (Cat. #30-2020, ATCC, Manassas, VA) and 1% penicillin–streptomycin (Cat. #30-2300, ATCC, Manassas, VA). Cells were harvested using Accutase (Cat. #AT104, Innovative Cell Technologies, Carlsbad, CA) as per the manufacturer's instruction.

Reagents Used. Tetrameric antibody complexes, TAC, from Stem Cell technologies (Vancouver, BC) were used to immunomagnetically label peripheral blood lymphocytes, PBLs. The specific TAC used in this study targeted the CD45 cell surface antigen and the dextran coated, magnetic nanoparticles. 0.5 μ L of the TAC complex was added per million cells.

Separation Methodology. The immunomagnetic separation was carried out as described in Yang et al. 2009²⁴ and will only be summarized here. An overall view of the separation process is shown in Figure 1. Red blood cells in the blood samples were lysed by mixing blood and lysis

(20) Schlimok, G.; Riethmuller, G. Detection, characterization and tumorigenicity of disseminated tumor cells in human bone marrow. *Semin. Cancer Biol.* **1990**, *1* (3), 207–15.

(21) Partridge, M. Detection of rare disseminated tumor cells identifies head and neck cancer patients at risk of treatment failure. *Clin. Cancer Res.* **2003**, *9* (14), 5287–94.

(22) Nagrath, S. Isolation of rare circulating tumour cells in cancer patients by microchip technology. *Nature* **2007**, *450* (7173), 1235–9.

(23) Zheng, S. Membrane microfilter device for selective capture, electrolysis and genomic analysis of human circulating tumor cells. *J. Chromatogr. A* **2007**, *1162* (2), 154–61.

(24) Yang, L. Optimization of an enrichment process for circulating tumor cells from the blood of head and neck cancer patients through depletion of normal cells. *Biotechnol. Bioeng.* **2009**, *102* (2), 521–34.




Figure 1. Flow diagram of current process to enrich for rare cancer cells in human blood.




Figure 2. Photograph of a gel of a RT-PCR product for EGFR for a positive sample, lane 1, a negative sample, lane 2, and molecular weight bands, lane 3.

buffer (154 mM NH_4Cl , 10 mM KHCO_3 , 0.1 mM EDTA) at a ratio of 1:25, incubating it for 5 min at room temperature before pelleting the remaining blood cells at 350g for 5 min. Nucleated cell concentration was estimated by diluting 20 μL with 3% acetic acid (1:25) and counting the cells using a hemocytometer. The cell suspension obtained, consisting mostly of nucleated cells, is then first labeled with FcR blocking reagent and then anti-CD45 TAC and incubated for 30 min at room temperature on a shaker. Without washing the cells, magnetic nanoparticles are then added to the cell suspension and incubated for 15 min at room temperature in a shaker. The immunomagnetically labeled cell suspension is subsequently run through the deposition quadrupole magnetic sorter (dQMS) system to obtain an enriched sample.

Confocal Images and Immunostaining. Confocal images were obtained using a Zeiss LSM 510 confocal/multi photon laser scanning microscope equipped with Argon/2 laser (458, 477, 488, 514 nm) and a Titanium Sapphire laser (750 nm). The cells were viewed with a 63 \times (NA 1.2) apochromatic water objective, and images of different fields were taken. The microscope was set up to take multitrack images and the excitation and emission filter sets configured individually so that there is no fluorescence bleed-through between the channels. The argon (488 nm) laser with appropriate emission filters was used for the visualization of FITC. The DAPI staining was viewed by the use of a tunable IR laser which excites the fluorophore using two photons at the point of focus. An aliquot of the enriched sample after immunomagnetic separation was preserved in neutral buffered formalin prior to preparing a cytospin. After washing the slides with PBS (Cat. #SH3025601, Thermo Sci Hyclone), the cells were incubated with anticytokeratin FITC CK3-6H5 (1:10) (Cat.




Figure 3. Histogram of the number of CTCs detected per milliliter of patient blood sample ($n = 32$).

#130-080-101, Miltenyi Biotech) for 30 min at 37 °C followed by a 30 min blocking with PBS pH 7.4 with 1% BSA at room temperature. The slides were then washed with PBS for 15 min, air-dried, and mounted with Vectashield mounting medium with 4',6-diamidino-2-phenylindole (DAPI) (Cat. # H-1200, Vector Laboratories).

RT-PCR Analysis. As presented previously by Tong et al. (2007)²⁵ and Yang et al. (2009),²⁴ a portion of the sample before enrichment and after enrichment was preserved in RNeasy lysis buffer for further molecular analysis. RNA was extracted from these samples using Trizol reagent (Cat. #15596-018) and/or Picopure RNA isolation kit (Cat. #KIT0204, Molecular Devices) depending upon the number of cells in the sample. Aliquots of 1 μg of RNA was reverse transcribed according to manufacturer's protocol. Following reverse transcription 2 μL of the cDNA product was amplified for EGFR generating a band at 301bp. The EGFR primer sequences are as follows: GGGAGCAGCGATGCGA and CTCCACTGTGTTGAG-GGCAAT. 35 cycles of PCR were performed, each consisting of a denaturation step (94 °C for 1 min), an annealing step (60 °C for 1 min) and an extension step (72 °C for 1.5 min). Final extension for 10 min at 72 °C was allowed, and the mixture was cooled down to 4 °C. RT-PCR products were separated by electrophoresis on a 1.5% agarose gel containing ethidium bromide and analyzed by direct visualization, and photographed, under UV light. Figure 2 presents representative photographs of a positive product from a patient, a negative product, and a lane containing molecular weight bands.

(25) Tong, X. Application of immunomagnetic cell enrichment in combination with RT-PCR for the detection of rare circulating head and neck tumor cells in human peripheral blood. *Cytometry, Part B* 2007, 72 (5), 310–23.




Figure 4. Set of progressive “z-slices” of confocal images of stained cells on a cytospin slide of an enriched peripheral blood sample from a head and neck cancer patient. Each of the four images is a computer combined digital image of separate images filtered for DAPI (nuclei) staining (blue) and FITC (cytokeratin) staining (green-yellow). A total of 14 “z-slices” were obtained, and panels A, B, C, and D correspond to slices 3, 5, 6, and 9, respectively. Note, the lower the number, the further from the objective; consequently, the largest view of nuclei, and the surrounding cytokeratin, is presented in panels B and C.

Results

As reported previously by Yang et al. (2009),²⁴ for a cell to be considered a CTC by visual observation using a fluorescent microscope, it must (1) be positive for a nucleus

based on DAPI stain, (2) be positive for cytokeratins based on FITC fluorescence, and (3) have a high nuclei to cytoplasmic ratio. In addition, for a blood sample to be considered positive for CTCs, the enriched sample must be





Figure 5. A combined (all z slices) confocal image of SCC4 cells after being processed on a cytopsin using the same labeling protocol as was used on the blood sample presented in Figure 4.

positive for EGFR, based on visual observations of gels run on RT-PCR product.


For a sample size of 32 blood samples from different patients, 63% of the samples contained CTCs, based on the identification protocol defined above. The number of CTCs detected per mL of blood collected ranged from 0 to 214. The final number of CTCs were estimated by counting the number of CTCs on the slide and multiplying it by the corresponding dilution factor. Figure 3 is a histogram indicating the distribution of these 32 samples with respect to the number of CTCs per mL of blood. Using the optimized enrichment protocol reported by Yang et al. (2009),²⁴ the final purity of the enriched sample is

a function of the number of CTCs recovered as well as the final number of non-CTCs remaining. In our study, this purity defined as the ratio of number of CTCs to number of non-CTCs ranged from 1 CTC in 9 total cells to 1 CTC in 25,000 total cells. All of these patient samples were also analyzed for EGFR expression by RT-PCR. For samples in which no cytokeratin positive cells were identified based on visual inspection, the RT-PCR analysis for EGFR was also negative. Conversely, for all but two of the cytopsin samples that were considered positive for CTCs, the samples were positive for EGFR.

To further determine the accuracy of our ability to visually identify and characterize the CTCs, we analyzed a number of the cytopsin samples of the enriched samples using confocal microscopy. Figure 4 is a set of confocal images of an enriched, peripheral blood sample from a HNSCC patient. A total of 14 “z slices” were taken, and Figures 4A through 4D are representative, progressive slices moving from the bottom of the cytopsin to the top. As can be observed, the nuclei are most pronounced in Figures 4B and 4C, while the cytokeratin is present around the nuclei, as expected. Also, it can be noted that some of the cells in the field of view do not fit the criteria to be considered a CTC. In particular, in the lower left is a cell, negative for cytokeratins and with a “kidney shaped” nucleus, suggestive of being a monocyte or granulocyte. Near the center of Figures 4A–4D, cytokeratin positive cellular debris can be observed. While not counted as a CTC, this debris most likely represents a CTC that had undergone apoptosis. For the specific patient sample shown in Figure 4, a total of 214 CTCs per mL of blood was determined, and the final purity was 1 CTC in 31 non-cytokeratin positive cells. Obviously, visual inspection



A



B

Figure 6. A confocal image of a stained cytopsin of an enriched peripheral blood sample prior to (A) and during surgery (B). Note the presence of what appears to be an embolus in panel B.

of Figure 4 indicates a different ratio of cytokeratin positive to cytokeratin negative cells. Care was taken in the inspection of the cytospin to find this specific image to maximize the number of cytokeratin positive cells in a single field of view.

As a control, a cytospin using the same staining protocol used on patient samples was made using cultured SCC-4 cells, and a confocal image of a single, z-slice of these stained cells is presented in Figure 5. As expected, the cells stained positively for the nuclei and cytokeratins which surround the nuclei.

To demonstrate the reproducibility of these confocal images and the presence of CTCs in patients' blood prior to surgery, samples were taken before and after surgery and processed using the same separation and staining protocol mentioned earlier. Figures 6A and 6B are representative images of two cytospins from a sample taken prior to and during surgery, respectively.

Discussion

It is suggested that a number of significant conclusions can be made from the results presented in this study. First, a large range in concentration of CTCs in the peripheral blood of cancer patients exists. Second, confocal images indicate that the reagents used to target cytokeratins, a FITC conjugated pan anti-cytokeratin antibody, did in fact target structures in the cell's cytoplasm. Third, cytospins can be made that contain a very high percentage of cytokeratin positive cells. Fourth, these CTCs can be present in the blood of cancer patients, at high concentrations, prior to surgery, eliminating the possibility that the presence of CTCs is the result of tumor manipulation during surgery; however, it is still possible that manipulation can increase the number of CTCs. Fifth, CTCs can be of similar size to PBLs. It has been suggested in other publications that CTCs are typically larger than normal blood cells.

Ongoing studies in our laboratory involve studying the expression of other markers which have been reported to be potentially important with respect to metastatic cancer. These include cancer stem cell (CSC) markers which have been reported, or suggested to be identified, in various solid cancers including breast, prostate, colon, brain, pancreatic, and head and neck cancer.^{26–32}

Since a purely negative enrichment technology is used, it is also possible, especially with multicolor confocal microscopy, to investigate if any of the CTCs have mesenchymal characteristics. There is a growing body of literature which suggests that epithelial cancer undergoes a process called *epithelial–mesenchymal transition* (EMT) through which the tumor cells undergo a loss of polarity, lose cell–cell junctions, and acquire a mesenchymal phenotype. It is also proposed that this mesenchymal phenotype is “motile”, “migratory” and “invasive” and it also promotes the escape of the cancer cells from the primary site and leads to the development of metastases.^{33–36}

Acknowledgment. Grant support is gratefully acknowledged: National Science Foundation (BES-0124897 to J.J.C.) the National Cancer Institute (R01 CA62349 to M.Z., R01 CA97391-01A1 to J.J.C.) and the State of Ohio Third Frontier Program (ODOD 26140000: TECH 07-001).

MP9000519

(26) Al-Hajj, M. Prospective identification of tumorigenic breast cancer cells. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100* (7), 3983–8.

- (27) Collins, A. T. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res.* **2005**, *65* (23), 10946–51.
- (28) O'Brien, C. A. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature* **2007**, *445* (7123), 106–10.
- (29) Singh, S. K. Identification of human brain tumour initiating cells. *Nature* **2004**, *432* (7015), 396–401.
- (30) Lee, C. J.; Li, C.; Simeone, D. M. Human pancreatic cancer stem cells: implications for how we treat pancreatic cancer. *Transl. Oncol.* **2008**, *1* (1), 14–8.
- (31) Prince, M. E. Identification of a subpopulation of cells with cancer stem cell properties in head and neck squamous cell carcinoma. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104* (3), 973–8.
- (32) Reya, T. Stem cells, cancer, and cancer stem cells. *Nature* **2001**, *414* (6859), 105–11.
- (33) Chaffer, C. L.; Thompson, E. W.; Williams, E. D. Mesenchymal to epithelial transition in development and disease. *Cells Tissues Organs* **2007**, *185* (1–3), 7–19.
- (34) Berx, G. Pre-EMTing metastasis? Recapitulation of morphogenetic processes in cancer. *Clin. Exp. Metastasis* **2007**, *24* (8), 587–97.
- (35) Sarrio, D. Epithelial-mesenchymal transition in breast cancer relates to the basal-like phenotype. *Cancer Res.* **2008**, *68* (4), 989–97.
- (36) Bagnato, A.; Rosano, L. Epithelial-mesenchymal transition in ovarian cancer progression: a crucial role for the endothelin axis. *Cells Tissues Organs* **2007**, *185* (1–3), 5–94.