

Synthesis of a Highly Water-Soluble Derivative of Amphotericin B with Attenuated Proinflammatory Activity

Samusi A. Adediran, Timothy P. Day, Diptesh Sil, Matthew R. Kimbrell, Hemamali J. Warshakoon, Subbalakshmi S. Malladi, and Sunil A. David*

Department of Medicinal Chemistry, University of Kansas, Lawrence, Kansas 66047

Received June 27, 2009; Revised Manuscript Received August 4, 2009; Accepted August 7, 2009

Abstract: Amphotericin B (AmB), a well-known polyene antifungal agent, displays a marked tendency to self-associate and, as a consequence, exhibits very poor solubility in water. The therapeutic index of AmB is low and is associated with significant dose-related nephrotoxicity, as well as acute, infusion-related febrile reactions. Reports in the literature indicate that the toxicity of AmB may be related to the physical state of the drug. Reaction of AmB in dimethylformamide with bis(dimethylaminopropyl)carbodiimide yielded an unexpected *N*-alkylguanidine/*N*-acylurea bis-adduct of AmB which was highly water-soluble. The absorption spectrum of the AmB derivative in water indicated excellent monomerization, and the antifungal activities of reference AmB and its water-soluble derivative against *Candida albicans* were found to be virtually identical. Furthermore, the water-soluble adduct is significantly less active in engaging TLR4, which would suggest that the adduct may be less proinflammatory.

Keywords: Amphotericin B; solubility; toxicity; aggregation; carbodiimide; antifungal; toll-like receptors

Introduction

Amphotericin B (AmB) is a polyene antifungal agent first isolated from *Streptomyces nodosus* in 1955 from Venezuelan soil samples near the Orinoco River region.¹ For more than a half-century, AmB has remained the cornerstone of the therapy of serious systemic fungal infections,^{2,3} and even with the advent of alternate antifungal agents, such as the triazoles (voriconazole^{4–6} and posaconazole^{7,8}) and the echinocandins (caspofungin⁹), AmB is likely to remain an important therapeutic modality in the management of fulminant fungal infections.¹⁰




Figure 1. The structure of amphotericin B.

Amphotericin B, as the name suggests, is amphoteric, owing to the presence of carboxylate and amine functional groups and, is also amphipathic due to the asymmetric distribution of polar hydroxyl groups on the face of the molecule, and a markedly hydrophobic, conjugated polyene hydrocarbon on the other (Figure 1). AmB displays a marked tendency to self-associate with a critical aggregation con-

* Corresponding author. Mailing address: Sunil A. David, MD, PhD, Department of Medicinal Chemistry, University of Kansas, Multidisciplinary Research Building, Room 320D, 2030 Becker Drive, Lawrence KS 66047. Tel: 785-864-1610. Fax: 785-864-1961. E-mail: sdavid@ku.edu.

- (1) Dutcher, J. D. The discovery and development of amphotericin B. *Dis. Chest.* **1968**, *54* (Suppl. 1), 296–298.
- (2) Chapman, S. W.; Sullivan, D. C.; Cleary, J. D. In search of the holy grail of antifungal therapy. *Trans. Am. Clin. Climatol. Assoc.* **2008**, *119*, 197–215; discussion, 215–216.
- (3) Ellis, D. Amphotericin B: spectrum and resistance. *J. Antimicrob. Chemother.* **2002**, *49* (Suppl. 1), 7–10.

- (4) Herbrecht, R.; Denning, D. W.; Patterson, T. F.; Bennett, J. E.; Greene, R. E.; Oestmann, J. W.; Kern, W. V.; Marr, K. A.; Ribaud, P.; Lortholary, O.; Sylvester, R.; Rubin, R. H.; Wingard, J. R.; Stark, P.; Durand, C.; Caillot, D.; Thiel, E.; Chandrasekar, P. H.; Hodges, M. R.; Schlamm, H. T.; Troke, P. F.; de Pauw, B. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. *N. Engl. J. Med.* **2002**, *347*, 408–415.


centration of ca. 0.2 $\mu\text{g/mL}$ ¹¹ and, as a consequence, exhibits very poor solubility in water (<1 $\mu\text{g/mL}$), as well as in apolar organic solvents such as cyclohexane (20 $\mu\text{g/mL}$).¹¹ The therapeutic index of AmB is low, and is associated with significant dose-related nephrotoxicity,^{12–14} as well as acute, infusion-related febrile reactions.^{15,16} The unfavorable physical properties of the drug have engendered a variety of approaches in innovative formulations which have had significant impact on the pharmacokinetics, tissue distribution, and efficacy/toxicity ratios. An excellent review has recently been published.¹¹

It is believed that the attenuated nephrotoxicity observed with the more recent liposomal formulations is at least, in part, ascribable to the rapid clearance of the colloidal particles by the reticuloendothelial system, leading to lower C_{max} , higher V_D , and a more sustained release of the drug due to the resultant depot effect;¹¹ reticuloendothelial sequestration observed with certain formulations perhaps serves as a form of passive drug-targeting and may likely be of benefit in

conditions such as visceral leishmaniasis.^{17,18} However, early reports in the literature indicate that highly dispersed, monomeric AmB enhances its selectivity while decreasing its toxicity,^{19–21} suggesting that toxicity may be related to the physical state of the drug.²² These earlier studies were performed with surfactants, some of which unfortunately showed synergistic toxicity with AmB,^{19,21} and it was desirable for purposes of testing the hypothesis that antifungal and toxic properties of AmB are dissociable with derivatives of AmB that would be intrinsically water-soluble without the need of any excipients.

Inspired by some of the AmB derivatives obtained by semisynthesis by Carreira's research group,^{23–25} and given that our laboratory has considerable expertise with polyamine chemistry, we sought to couple spermine to the carboxylic acid group of AmB using standard carbodiimide chemistry. The solubility of AmB proved to be a significant impediment, and the only solvent that was amenable was dimethylformamide (DMF), a polar, high-dielectric solvent which, coupled with the accidental omission of scavenging agents, favored a serendipitous and unexpected side-reaction to preponderate, yielding a fully water-soluble (>45 mg/mL) mixture of *N*-acylurea/*N*-alkylguanidine derivative of AmB. This derivative was found to be monomeric by spectrophotometry, exhibited equipotent antifungal activity against *Candida albicans*, and elicited toll-like receptor-2 (TLR2) and TLR4 driven NF- κ B responses in reporter gene assays to the same extent as the parent molecule.

- (5) Powers, J. H.; Dixon, C. A.; Goldberger, M. J. Voriconazole versus liposomal amphotericin B in patients with neutropenia and persistent fever. *N. Engl. J. Med.* **2002**, *346*, 289–290.
- (6) Walsh, T. J.; Pappas, P.; Winston, D. J.; Lazarus, H. M.; Petersen, F.; Raffalli, J.; Yanovich, S.; Stiff, P.; Greenberg, R.; Donowitz, G.; Schuster, M.; Rebolli, A.; Wingard, J.; Arndt, C.; Reinhardt, J.; Hadley, S.; Finberg, R.; Laverdiere, M.; Perfect, J.; Garber, G.; Fioritoni, G.; Anaissie, E.; Lee, J. Voriconazole compared with liposomal amphotericin B for empirical antifungal therapy in patients with neutropenia and persistent fever. *N. Engl. J. Med.* **2002**, *346*, 225–234.
- (7) Smith, W. J.; Drew, R. H.; Perfect, J. R. Posaconazole's impact on prophylaxis and treatment of invasive fungal infections: an update. *Expert. Rev. Anti-Infect. Ther.* **2009**, *7*, 165–181.
- (8) Morris, M. I. Posaconazole: a new oral antifungal agent with an expanded spectrum of activity. *Am. J. Health-Syst. Pharm.* **2009**, *66*, 225–236.
- (9) Garcia-Vidal, C.; Carratala, J. Echinocandins for candidemia. *N. Engl. J. Med.* **2006**, *355*, 2791–2792.
- (10) Bohme, A.; Ruhnke, M.; Buchheidt, D.; Cornely, O. A.; Einsele, H.; Enzensberger, R.; Hebart, H.; Heinz, W.; Junghanss, C.; Karthaus, M.; Kruger, W.; Krug, U.; Kubin, T.; Penack, O.; Reichert, D.; Reuter, S.; Silling, G.; Sudhoff, T.; Ullmann, A. J.; Maschmeyer, G. Treatment of invasive fungal infections in cancer patients—recommendations of the Infectious Diseases Working Party (AGIHO) of the German Society of Hematology and Oncology (DGHO). *Ann. Hematol.* **2009**, *88*, 97–110.
- (11) Torrado, J. J.; Espada, R.; Ballesteros, M. P.; Torrado-Santiago, S. Amphotericin B formulations and drug targeting. *J. Pharm. Sci.* **2008**, *97*, 2405–2425.
- (12) Fanos, V.; Cataldi, L. Amphotericin B-induced nephrotoxicity: a review. *J. Chemother.* **2000**, *12*, 463–470.
- (13) Miano-Mason, T. M. Mechanisms and management of amphotericin B-induced nephrotoxicity. *Cancer Pract.* **1997**, *5*, 176–181.
- (14) Saliba, F.; Dupont, B. Renal impairment and amphotericin B formulations in patients with invasive fungal infections. *Med. Mycol.* **2008**, *46*, 97–112.
- (15) Khoo, S. H.; Bond, J.; Denning, D. W. Administering amphotericin B—a practical approach. *J. Antimicrob. Chemother.* **1994**, *33*, 203–213.
- (16) Gallis, H. A.; Drew, R. H.; Pickard, W. W. Amphotericin B: 30 years of clinical experience. *Rev. Infect. Dis.* **1990**, *12*, 308–329.
- (17) Berman, J. ABLE: a new and improved amphotericin B for visceral leishmaniasis? *Am. J. Trop. Med. Hyg.* **2009**, *80*, 689–690.
- (18) Thornton, S. J.; Wasan, K. M. The reformulation of amphotericin B for oral administration to treat systemic fungal infections and visceral leishmaniasis. *Expert. Opin. Drug Delivery* **2009**, *6*, 271–284.
- (19) Gruda, I.; Milete, D.; Brother, M.; Kobayashi, G. S.; Medoff, G.; Brajtburg, J. Structure-activity study of inhibition of amphotericin B (Fungizone) binding to sterols, toxicity to cells, and lethality to mice by esters of sucrose. *Antimicrob. Agents Chemother.* **1991**, *35*, 24–28.
- (20) Tancrede, P.; Barwicz, J.; Jutras, S.; Gruda, I. The effect of surfactants on the aggregation state of amphotericin B. *Biochim. Biophys. Acta* **1990**, *1030*, 289–295.
- (21) Barwicz, J.; Christian, S.; Gruda, I. Effects of the aggregation state of amphotericin B on its toxicity to mice. *Antimicrob. Agents Chemother.* **1992**, *36*, 2310–2315.
- (22) Legrand, P.; Romero, E. A.; Cohen, B. E.; Bolard, J. Effects of aggregation and solvent on the toxicity of amphotericin B to human erythrocytes. *Antimicrob. Agents Chemother.* **1992**, *36*, 2518–2522.
- (23) Paquet, V.; Carreira, E. M. Significant improvement of antifungal activity of polyene macrolides by bisalkylation of the mycosamine. *Org. Lett.* **2006**, *8*, 1807–1809.
- (24) Paquet, V.; Volmer, A. A.; Carreira, E. M. Synthesis and in vitro biological properties of novel cationic derivatives of amphotericin B. *Chemistry* **2008**, *14*, 2465–2481.
- (25) Zumbuehl, A.; Stano, P.; Sohrmann, M.; Peter, M.; Walde, P.; Carreira, E. M. A novel strategy for bioconjugation: synthesis and preliminary evaluation with amphotericin B. *Org. Biomol. Chem.* **2007**, *5*, 1339–1342.

Scheme 1. Synthesis of Bis(dimethylaminopropyl)carbodiimide


Materials and Methods

All of the solvents and reagents used were obtained commercially and used as such unless noted otherwise. Moisture or air sensitive reactions were conducted with anhydrous solvents under argon atmosphere in oven-dried (120 °C) glass apparatus. Solvents were removed under reduced pressure using standard rotary evaporators. Flash column chromatography was carried out using silica gel 60 (230–400 mesh), and thin layer chromatography was performed using silica gel HLF, precoated glass plates. All yields reported refer to isolated material judged to be homogeneous by thin-layer chromatography and NMR spectroscopy. Unless noted otherwise, NMR spectra were recorded with the chemical shifts (δ) reported in ppm relative to Me₄Si (for ¹H) and CDCl₃ (for ¹³C) as internal standards respectively.

Synthesis of Bis(dimethylaminopropyl)carbodiimide [N¹-(3-(dimethylamino)propylimino) methylene)-N³,N³-dimethylpropane-1,3-diamine] (Scheme 1). 2.5 g of 1,1'-thiocarbonyldiimidazole (**2**, 14 mmol) was heated in 50 mL of toluene at 110 °C, to which was added 1.8 equiv (2.6 g, 25 mmol, 3.2 mL, $d = 0.812$) of N¹,N¹-dimethylpropane-1,3-diamine. The reaction mixture was allowed to reflux at 110 °C overnight. The thiourea **3** (Scheme 1) was isolated by using a binary DCM/MeOH system on a 40 g normal-phase RediSep Rf silica cartridge using a CombiFlash Companion (Teledyne-Isco, Lincoln, NE) flash chromatography system (R_f : 0.3, 0.927 g, 3.77 mmol, 27% yield). The thiourea **3** was redissolved in DCM and subjected to desulfuration²⁶ by stirring with 2.0 equiv of HgO for 2 h at room temperature. After filtration of the mercuric oxide through Celite, 2.0 mL of HCl in dioxane was added and the solvent evaporated to furnish the bis(dimethylaminopropyl)carbodiimide **4** (0.4038 g, 1.9 mmol, 50% yield). ¹H NMR (CDCl₃, 400 MHz): δ 1.7–1.77 (m, 4H), 2.23 (s, 12H), 2.34 (t, $J = 7.12$ Hz, 4H), 3.28 (t, $J = 6.8$ Hz, 4H). ¹³C NMR (CDCl₃, 125 MHz): δ 29.38, 44.82, 45.54, 56.93. MH⁺: 213.212.

Synthesis of AmB–Carbodiimide Adducts (Scheme 2). To a solution of 43.3 mg of AmB (**1**, 46.5 μ mol) in 20 mL of dimethylformamide (DMF) were added 2.0 equiv of the carbodiimide **4** and a catalytic amount of 4-dimethylaminopyridine (DMAP). The solution was stirred overnight. The solvent was evaporated, and the resultant solid (45.7 mg) was analyzed by LC–MS as described below and found to be a near-equal mixture of the *N*-acylurea/*N*-alkylguanidine diadduct **5** (48%) and *N*-alkylguanidine monoadduct **6** (46%), with traces of the *N*-acylurea monoadduct **7**, the

structures of which are shown in Scheme 2. Further purification after redissolving the crude solid in a minimal volume of MeOH and precipitation by adding it to a large volume of ice-cold diethyl ether resulted in a substantial enrichment of the *N*-acylurea/*N*-alkylguanidine diadduct (>85%). The derivative was freely soluble in pure water at 45 mg/mL, the highest concentration tested. This material was used in all further assays.

LC–MS Characterization. The AmB derivative was analyzed by reverse-phase LC–ESI-MS performed on a Shimadzu LC system (LC-10AD binary pumps, SCL-10A diode array detector) using a Zorbax 3.0 mm \times 150 mm 3.5 μ m stable-bond C₁₈ reverse-phase column with a forty-minute binary gradient (CH₃CN/water, 0.1% HCOOH) from 5% to 95% of CH₃CN. ESI-MS data was acquired on an Agilent LC/MSD-TOF instrument with a mass accuracy of 3 ppm and a range of 100–3500 Da. Calibration drift was minimized on a scan-by-scan basis to less than 2 ppm by using internal standards corresponding to 922.0001 and 2721.0201 marker ions infused concurrently through a second nebulizer in the ionization chamber. MS acquisition was commenced 10 min after injection.


Absorption Spectrophotometry. Absorption spectra of reference AmB in DMSO and the AmB derivative in water were measured using a SpectraMax Plus 384 instrument (Molecular Devices, Sunnyvale, CA) from which appropriate solvent-only blank spectra were subtracted.

Determination of Minimum Inhibitory Concentration (MIC). Minimum inhibitory concentrations of reference AmB (stock solution at 1 mg/mL in DMSO) and the water-soluble AmB adduct (stock solution at 1 mg/mL in H₂O) were determined by broth microdilution method²⁷ as per NCCLS guidelines. This assay has been described previously in detail for antibacterial compounds.²⁸ Compounds were 2-fold serially diluted in a 384-well microtiter plate in Sabouraud's Dextrose Broth using a Biotek Precision 2000 automated microplate pipetting system. A single colony of *C. albicans* ATCC 26278 was dispersed in Sabouraud's Dextrose Broth and was added to the wells with appropriate vehicle-only controls. All experiments were done in quadruplicate. The microtiter plates were incubated overnight at 37 °C. The plates were read at an absorbance of 600 nm.

(26) Sheehan, J. C.; Cruickshank, P. A.; Boshart, G. L. A convenient synthesis of water-soluble carbodiimides. *J. Org. Chem.* **1961**, *26*, 2525–2528.

(27) European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution *Clin. Microbiol. Infect.* **2003**, *9*, 1–7.

(28) Balakrishna, R.; Wood, S. J.; Nguyen, T. B.; Miller, K. A.; Suresh Kumar, E. V.; Datta, A.; David, S. A. Structural correlates of antibacterial and membrane-permeabilizing activities in acylpolyamines. *Antimicrob. Agents Chemother.* **2006**, *50*, 852–861.

Scheme 2. Synthesis of AmB–DMAPCI Adducts

TLR2 and TLR4-Specific Reporter Gene Assays for Assessing Agonistic Potencies. Human embryonic kidney 293 (HEK-293) cells stably transfected (individually) with either human TLR2 or TLR4 were obtained from InvivoGen, Inc. (San Diego, CA) and were additionally nucleofected with 1 μ g of purified *pnifty2* reporter plasmid (InvivoGen) using an Amaxa Nucleofector (Gaithersburg, MD). The *pnifty2* reporter plasmid expresses a secreted embryonic alkaline phosphatase (SEAP) gene under the control of a promoter inducible by NF- κ B and AP-1 transcription factors. Stable transfectants were selected, expanded, and maintained in HEK-Blue Selection Medium (InvivoGen) containing zeocin (10 μ g/mL) and normocin (200 μ g/mL). The induction of NF- κ B was quantitated in a rapid-throughput, homogeneous format as follows: cells were incubated at a density of $\sim 10^6$ cells/mL in a volume of 80 μ L/well, in 384-well, flat-bottomed, cell culture-treated microtiter plates until confluency was achieved, and subsequently stimulated with graded concentrations of TLR agonists using an automated BioTek Precision 2000 Liquid Handler. SEAP in the medium was

assayed spectrophotometrically using an alkaline phosphatase-specific chromogen (present in HEK-detection medium as supplied by the vendor) at 650 nm. These assays have been described in detail previously by us.^{29–31}

- (29) Kimbrell, M. R.; Warshakoon, H.; Cromer, J. R.; Malladi, S.; Hood, J. D.; Balakrishna, R.; Scholdberg, T. A.; David, S. A. Comparison of the immunostimulatory and proinflammatory activities of candidate Gram-positive endotoxins, lipoteichoic acid, peptidoglycan, and lipopeptides, in murine and human cells. *Immunol. Lett.* **2008**, *118*, 132–141.
- (30) Sil, D.; Shrestha, A.; Kimbrell, M. R.; Nguyen, T. B.; Adisechan, A. K.; Balakrishna, R.; Abbo, B. G.; Malladi, S.; Miller, K. A.; Short, S.; Cromer, J. R.; Arora, S.; Datta, A.; David, S. A. Bound to Shock: Protection from Lethal Endotoxemic Shock by a Novel, Nontoxic, Alkylpolyamine Lipopolysaccharide Sequestrant. *Antimicrob. Agents Chemother.* **2007**, *51*, 2811–2819.
- (31) Wu, W.; Sil, D.; Szostak, M. L.; Malladi, S. S.; Warshakoon, H. J.; Kimbrell, M. R.; Cromer, J. R.; David, S. A. Structure-activity relationships of lipopolysaccharide sequestration in guanylhydrazone-bearing lipopolyamines. *Bioorg. Med. Chem.* **2008**, *17* (2), 709–715.




Figure 2. 2D (A) and 3D (B) plots of absorbance-versus-time profiles from photodiode array detector output of a chromatographic separation of the crude AmB–DMAPCI adducts. All three major bands show absorption spectra characteristic of AmB. The corresponding mass spectra are shown in Figure 3.

Results and Discussion

Our original intent, as mentioned earlier, in seeking to attach spermine to the carboxylate of AmB (Figure 1) in order to test if the polyamine would confer more favorable solubility began earnestly, if somewhat naïvely, with *N*-Boc protection of the mycosamine amine followed by coupling of Tri-Boc spermine^{30,32} using standard coupling procedures. Having overcome successive solubility and stability issues, and after purifying in very small yields to homogeneity the final precursor, deprotection of the Boc groups with trifluoroacetic acid resulted, to our dismay, in instantaneous and complete degradation of the molecule. Chastened, we wanted

to test if AmB could be coupled at all, even non-regioselectively, to the unprotected polyamine using the more polar 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide (EDCI). Solubility issues left us with no choice but to carry out the reaction in DMF. To our surprise, aqueous workup showed the distinct yellow-orange color of AmB partitioning largely into the water layer. LC–MS analysis of the aqueous layer indicated the presence of multiple isobaric species which eluted off the column at different times, but with masses corresponding to AmB–EDCI adduct which we assigned to the more stable and commonly encountered rearranged *N*-acylurea side product (Scheme 2).^{33–35} We reasoned (incorrectly, as it turns out) that the multiplicity of isobaric

(32) Miller, K. A.; Suresh Kumar, E. V. K.; Wood, S. J.; Cromer, J. R.; Datta, A.; David, S. A. Lipopolysaccharide Sequestrants: Structural Correlates of Activity and Toxicity in Novel Acylhomospermines. *J. Med. Chem.* **2005**, *48*, 2589–2599.

(33) Nakajima, N.; Ikada, Y. Mechanism of amide formation by carbodiimide for bioconjugation in aqueous media. *Bioconjugate Chem.* **1995**, *6*, 123–130.

species could be a consequence of *N*-acylurea regioisomers (EDCI being nonsymmetrical), and hypothesized that a symmetric carbodiimide with two amines would help resolve the isobaric problem while affording even greater solubility to the adduct. Hence we synthesized the bis(dimethylaminopropyl)carbodiimide **4** (Scheme 1). Reaction of AmB with **4**, to our surprise, yielded an unexpected bis-adduct in abundance, in addition to two isobaric products with substantially different retention times (Figures 2 and 3, peak 1). Closer examination of mass chromatograms acquired under different fragmenter voltages clearly showed that only one species ionized as both proton and sodium adducts (Figure 3, peak 2), and also showed classic —OH^+ fragments, indicating the presence of a free —COOH group, and thereby implicating the amine as the alternate nucleophile. Although we had not anticipated finding an *N*-alkylguanidine adduct, reactions of amines and carbodiimides to furnish substituted guanidines have indeed been reported in the literature.^{36,37} The other monoadduct corresponding to the *N*-acylurea adduct ionized solely as singly charged proton adduct at higher fragmenter voltages (Figure 3, peak 3) and as $+2\text{H}^+$ doubly charged species at lower voltages (data not shown). The use of 2 equiv of the carbodiimide was fortuitous in retrospect, for it provided an insight into the relative reactivity of the amine and the carboxylate of AmB with the carbodiimide that we perhaps would not have gained had we used an excess of the reagent and obtained entirely the diadduct **5**.

Precipitation of the crude product in diethyl ether results in substantial enrichment (>85%) of the *N*-alkylguanidine/*N*-acylurea diadduct **5**, which dissolved freely in water to more than 45 mg/mL (the highest we could test), which is >45,000 the aqueous solubility of the native polyene macrolide. The absorption spectrum of **5** in water (Figure 4) indicates that the predominant form is the monomeric one.^{38,39} Gratifyingly, the adduction of the alkylaminocarbodiimides to both the amine and carboxylic acid groups of AmB does not perturb significantly the function of the molecule, and the antifungal activities of reference AmB and

5 against *C. albicans* are quite comparable (Figure 5). The availability of structurally well-defined and intrinsically highly water-soluble AmB derivatives such as **5** should allow for a critical reexamination of such concepts as aggregation-dependent selectivity and toxicity given that the critical micellar concentrations of **5** are expected to be orders of magnitude higher than AmB.^{40–43}

In addition to the dose-limiting and occasionally fatal nephrotoxicity of AmB, acute infusion-related febrile reactions occur frequently. These are thought to be related to proinflammatory mediator production as a consequence of the activation of toll-like receptors (TLRs), especially TLR2 and TLR4.^{44–47} Reporter gene assays for both these TLRs are in routine use in our laboratory,²⁹ and it was of interest to compare TLR2 and TLR4 agonistic potencies of **5** relative to AmB. As shown in Figure 5, **5** is more potent than AmB in activating TLR2, while the water-soluble adduct is significantly less active in engaging TLR4 (Figure 5, bottom panel), which would suggest that the adduct **5** may be less proinflammatory. This is currently being tested using *ex vivo* human blood model systems employing assays of proinflammatory cytokine release and indices of monocyte and natural killer lymphocytic activation, and will be the subject of a future publication.

- (34) Izdebski, J.; Orlowska, A.; Anulewicz, R.; Witkowska, E.; Fiertek, D. Reinvestigation of the reactions of carbodiimides with alkoxy-carbonylamino acid symmetrical anhydrides. Isolation of two *N*-acylureas. *Int. J. Pept. Protein Res* **1994**, *43*, 184–189.
- (35) Iwasawa, T.; Wash, P.; Gibson, C.; Rebek, J. Reaction of an Introverted Carboxylic Acid with Carbodiimide. *Tetrahedron* **2007**, *63*, 6506–6511.
- (36) Hopkins, T. P.; Dener, J. M.; Boldi, A. M. Solid-phase synthesis of trisubstituted guanidines. *J. Comb. Chem.* **2002**, *4*, 167–174.
- (37) Rowley, C. N.; Ong, T. G.; Priem, J.; Richeson, D. S.; Woo, T. K. Analysis of the Critical Step in Catalytic Carbodiimide Transformation: Proton Transfer from Amines, Phosphines, and Alkynes to Guanidines, Phosphoguanidines, and Propiolamidines with Li and Al Catalysts. *Inorg. Chem.* **2008**, *47*, 12024–12031.
- (38) Adams, M.; Kwon, G. S. Spectroscopic investigation of the aggregation state of amphotericin B during loading, freeze-drying, and reconstitution of polymeric micelles. *J. Pharm. Pharm. Sci.* **2004**, *7*, 1–6.

- (39) Sanchez-Brunete, J. A.; Dea, M. A.; Rama, S.; Bolas, F.; Alunda, J. M.; Torrado-Santiago, S.; Torrado, J. J. Amphotericin B molecular organization as an essential factor to improve activity/toxicity ratio in the treatment of visceral leishmaniasis. *J. Drug Targeting* **2004**, *12*, 453–460.
- (40) Bolard, J. How do the polyene macrolide antibiotics affect the cellular membrane properties. *Biochim. Biophys. Acta* **1986**, *864*, 257–304.
- (41) Bolard, J. Interaction of polyene antibiotics with membrane lipids: physicochemical studies of the molecular basis of selectivity. *Drugs Exp. Clin. Res.* **1986**, *12*, 613–618.
- (42) Bolard, J.; Legrand, P.; Heitz, F.; Cybulska, B. One-sided action of amphotericin B on cholesterol-containing membranes is determined by its self-association in the medium. *Biochemistry* **1991**, *30*, 5707–5715.
- (43) Sanchez-Brunete, J. A.; Dea, M. A.; Rama, S.; Bolas, F.; Alunda, J. M.; Torrado-Santiago, S.; Torrado, J. J. Amphotericin B molecular organization as an essential factor to improve activity/toxicity ratio in the treatment of visceral leishmaniasis. *J. Drug Targeting* **2004**, *12*, 453–460.
- (44) Meletiadis, J.; Chanock, S.; Walsh, T. J. Defining targets for investigating the pharmacogenomics of adverse drug reactions to antifungal agents. *Pharmacogenomics* **2008**, *9*, 561–584.
- (45) Sau, K.; Mambula, S. S.; Latz, E.; Henneke, P.; Golenbock, D. T.; Levitz, S. M. The antifungal drug amphotericin B promotes inflammatory cytokine release by a Toll-like receptor- and CD14-dependent mechanism. *J. Biol. Chem.* **2003**, *278*, 37561–37568.
- (46) Razonable, R. R.; Henault, M.; Lee, L. N.; Laethem, C.; Johnston, P. A.; Watson, H. L.; Paya, C. V. Secretion of proinflammatory cytokines and chemokines during amphotericin B exposure is mediated by coactivation of toll-like receptors 1 and 2. *Antimicrob. Agents Chemother.* **2005**, *49*, 1617–1621.
- (47) Bellocchio, S.; Gaziano, R.; Bozza, S.; Rossi, G.; Montagnoli, C.; Perruccio, K.; Calvitti, M.; Pitzurra, L.; Romani, L. Liposomal amphotericin B activates antifungal resistance with reduced toxicity by diverting Toll-like receptor signalling from TLR-2 to TLR-4. *J. Antimicrob. Chemother.* **2005**, *55*, 214–222.




Figure 3. Positive ion ESI MS spectra of the chromatographic separation of the crude AmB–DMAPCI adducts. Data were acquired on an LC/MSD-TOF with a mass accuracy of 3 ppm. Note that peaks 2 and 3 are isobaric; peak 2 has a free carboxylate that ionizes also as the sodium adduct. The 1118.6788 amu signal in peak 2 is also diagnostic of a carboxylate ($-OH^+$ fragment; $M - 17.0027$).

The assumption concerning the origin of nephrotoxicity has been one of dose-dependent direct cytotoxicity on renal

epithelium. This notion is partly supported by observations that cumulative plasma concentrations (and therefore possibly




Figure 4. Absorption spectra of reference AmB (in DMSO) and the AmB-DMAPCI derivative (in water).




Figure 5. Inhibition of *C. albicans* growth by reference AmB (in DMSO) and the AmB-DMAPCI derivative (in water) determined by microplate dilution method. Data represent means and SD obtained on quadruplicates.

renal exposure to the drug) are lower with the less toxic liposomal preparations (such as AmBiosome and Abelset), compared to Fungizone.^{11,48–51} An alternate, hitherto unexamined, and testable hypothesis is that given the substantial TLR-driven proinflammatory activity of AmB (or of its more water-soluble metabolites), mesangial inflammation^{52–55} in the renal parenchymal interstitium, rather than tubular




Figure 6. TLR2- (A) and TLR4- (B) dependent activation of NF- κ B by reference AmB (in DMSO) and the AmB-DMAPCI derivative (in water) determined using TLR-specific reporter gene assays. Data represent means and SD obtained on quadruplicates.

damage is the primary cause of nephrotoxicity. If this hypothesis is correct, then a bolus dose of **5** should result in

- (48) Adedoyin, A.; Bernardo, J. F.; Swenson, C. E.; Bolsack, L. E.; Horwith, G.; DeWit, S.; Kelly, E.; Klasterksy, J.; Sculier, J. P.; DeValeriola, D.; Anaissie, E.; Lopez-Berestein, G.; Llanos-Cuentas, A.; Boyle, A.; Branch, R. A. Pharmacokinetic profile of ABELCET (amphotericin B lipid complex injection): combined experience from phase I and phase II studies. *Antimicrob. Agents Chemother.* **1997**, *41*, 2201–2208.
- (49) Adedoyin, A.; Swenson, C. E.; Bolsack, L. E.; Hellmann, A.; Radowska, D.; Horwith, G.; Janoff, A. S.; Branch, R. A. A pharmacokinetic study of amphotericin B lipid complex injection (Abelcet) in patients with definite or probable systemic fungal infections. *Antimicrob. Agents Chemother.* **2000**, *44*, 2900–2902.

- (50) Bekersky, I.; Fielding, R. M.; Dressler, D. E.; Lee, J. W.; Buell, D. N.; Walsh, T. J. Pharmacokinetics, excretion, and mass balance of liposomal amphotericin B (AmBisome) and amphotericin B deoxycholate in humans. *Antimicrob. Agents Chemother.* **2002**, *46*, 828–833.
- (51) Bekersky, I.; Fielding, R. M.; Dressler, D. E.; Lee, J. W.; Buell, D. N.; Walsh, T. J. Plasma protein binding of amphotericin B and pharmacokinetics of bound versus unbound amphotericin B after administration of intravenous liposomal amphotericin B (AmBisome) and amphotericin B deoxycholate. *Antimicrob. Agents Chemother.* **2002**, *46*, 834–840.
- (52) Sterzel, R. B.; Schulze-Lohoff, E.; Marx, M. Cytokines and mesangial cells. *Kidney Int. Suppl.* **1993**, *39*, S26–S31.
- (53) Radeke, H. H.; Resch, K. The inflammatory function of renal glomerular mesangial cells and their interaction with the cellular immune system. *Clin. Invest.* **1992**, *70*, 825–842.
- (54) Akira, S.; Hirano, T.; Taga, T.; Kishimoto, T. Biology of multifunctional cytokines: IL 6 and related molecules (IL 1 and TNF). *FASEB J.* **1990**, *4*, 2860–2867.
- (55) Gomez-Guerrero, C.; Hernandez-Vargas, P.; Lopez-Franco, O.; Ortiz-Munoz, G.; Egido, J. Mesangial cells and glomerular inflammation: from the pathogenesis to novel therapeutic approaches. *Curr. Drug Targets: Inflammation Allergy* **2005**, *4*, 341–351.

augmented nephrotoxicity, in which case the rational approach to limiting AmB toxicity would be in abrogating its TLR stimulatory activity. All known TLR2 and TLR4 agonists are amphipathic molecules as is AmB,²⁹ and TLRs, unlike classic receptors, are pattern-recognition molecules. Reducing amphipathicity by masking the polyhydroxyl domain may be one approach. These aspects are being examined.

Acknowledgment. This work was supported in part by University of Kansas Startup funds. S.A.A. and T.P.D. are grateful for support by the KUPREP (NIH 5R25GM078441) and IMSD (NIH 5R25GM062232) programs, respectively. The David lab wishes S.A.A. the very best as he embarks on his graduate studies.

MP9001602