

Cloning and Functional Expression of a Novel G_i Protein-Coupled Receptor for Adenine from Mouse Brain

Ivar von Kügelgen, Anke C. Schiedel, Kristina Hoffmann, Bernt B. A. Alsdorf, Aliaa Abdelrahman, and Christa E. Müller

Department of Pharmacology and Toxicology (I.v.K., K.H.) and Pharmaceutical Institute, Pharmaceutical Chemistry I (A.C.S., B.B.A.A., A.A., C.E.M.), University of Bonn, Bonn, Germany

Received April 12, 2007; accepted October 31, 2007

ABSTRACT

An orphan G protein-coupled receptor from the rat has recently been demonstrated to act as a transmembrane receptor for the nucleobase adenine. The receptor is possibly involved in nociception. Here we report the cloning and functional expression of an additional G_i-coupled receptor for adenine (Genbank accession code DQ386867). mRNA for this receptor was obtained from mouse brain and the mouse neuroblastoma x rat glioma hybrid cell line NG108-15. The new mouse protein sequence shares only 76% identity with that of the rat adenine receptor, suggesting that the receptors are not species homologs but distinct receptor subtypes. In human 1321N1 astrocytoma cells stably expressing the new mouse receptor, adenine and 2-fluoroadenine inhibited the isoproterenol-induced cAMP formation with IC₅₀ concentrations of 8 and 15

nM, respectively. The adenosine receptor antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX, 1 μM) as well as the P2 receptor antagonist suramin (300 μM) failed to change the responses to adenine. In contrast, pretreatment of cells with pertussis toxin abolished the effect of adenine. When the novel adenine receptor was expressed in Sf21 insect cells, a specific binding site for [³H]adenine was detected. In competition assays, the rank order of potency of selected ligands was identical to that obtained in membranes from NG108-15 cells and rat brain cortex (adenine > 2-fluoroadenine > 7-methyladenine > 1-methyladenine ≫ N⁶-dimethyladenine). In summary, our data show that a second mammalian DNA sequence encodes for a G_i-coupled GPCR activated by low, nanomolar concentrations of adenine.

Adenine was recently identified as the endogenous ligand of an orphan rat G protein-coupled receptor (GPCR) (Bender et al., 2002). The expression of mRNA for this receptor in dorsal root ganglia neurons (Bender et al., 2002), as well as reports on neurotrophic effects of adenine (Watanabe et al., 2003; Yoshimi et al., 2003), points to a role of adenine as a neuronal signaling molecule. Spinally administered adenine has been shown to facilitate electrically evoked neuronal responses in the rat dorsal horn, indicating an enhancement in nociception by the action of adenine (Matthews and Dickenson, 2004). Recent studies demonstrating the specific binding of [³H]adenine to membrane preparations provided further evidence for the expression of the adenine receptor in rat neuronal tissues as well as a number of cell lines of neuronal

origin (Gorzalka et al., 2005; Watanabe et al., 2005). The study by Gorzalka et al. (2005) also showed that an adenine receptor endogenously expressed in the mouse neuroblastoma x rat glioma hybrid cell line NG108-15 (Hamprecht, 1977) mediated an inhibition of adenylate cyclase activity. RT-PCR experiments indicated that NG108-15 cells express a mouse rather than a rat mRNA sequence encoding for an adenine receptor (Gorzalka et al., 2005). It is noteworthy that there is evidence for the occurrence of an adenine receptor in human cells (Gorzalka et al., 2005). In healthy humans, plasma concentrations of adenine amount to values of approximately 70 nM; in patients with chronic renal failure, the plasma concentration of adenine even increases to values above 1 μM (Slominska et al., 2002). Therefore, adenine may play important roles in the pathophysiology of chronic renal failure (for a recent report on a GPCR-mediated effect of adenine on the Na⁺-ATPase activity in the proximal tubule, see Wengert et al., 2007). The human genome, however, seems to encode no direct ortholog of the rat adenine receptor (see also Bender et al., 2002), indicating that one has to

I.v.K. and K.H. were supported by grants from the University of Bonn (Germany) and the Doktor Robert Pflieger Stiftung (Bamberg, Germany) and A.C.S., B.B.A.A., A.A. and C.E.M. were supported by the Deutsche Forschungsgemeinschaft (GRK-804/2).

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.107.037069.

ABBREVIATIONS: GPCR, G protein-coupled receptor; RT-PCR, reverse transcriptase-polymerase chain reaction; ANOVA, analysis of variance; CHO, Chinese hamster ovary; DPCPX, 8-cyclopentyl-1,3-dipropylxanthine; NECA, 5'-(N-ethylcarboxamido)adenosine; NG108-15 cells, mouse neuroblastoma x rat glioma hybrid cell line; PTX, pertussis toxin; Sf21 cells, *Spodoptera frugiperda* 21 cells.

search for a distinct sequence to identify human adenine receptors. Here, we report the cloning of a novel sequence encoding a GPCR for adenine. The sequence, which was found in mouse brain, mouse spleen, and NG108-15 cells, is clearly distinct from the mouse ortholog (mMrgA10) of the rat adenine receptor. Whether the mouse ortholog mMrgA10 or other members of the Mas-related gene family found in the mouse genome (Choi and Lahn, 2003; Zylka et al., 2003) operate as receptors for adenine is not known. We also demonstrate the functional expression of this second member of the family of GPCRs for adenine, which couples via a pertussis toxin-sensitive G protein to inhibition of adenylate cyclase, and characterize the pharmacological properties of the receptor. The presented results may facilitate the identification of a human GPCR for adenine. Some of these results have been presented at a meeting of the German Society of Pharmacology and Toxicology (von Kügelgen et al., 2007).

Materials and Methods

Molecular Biological Experiments. Poly A⁺ mRNA was isolated from homogenized brain tissue, the liver, the kidney and the spleen of adult male MNRI mice (Charles River WIGA, Sulzfeld, Germany) as well as from NG108-15 cells cultured as described previously (Kaulich et al., 2003) using the Oligotex Direct mRNA Mini Kit (QIAGEN, Hilden, Germany). Reverse transcriptase-polymerase chain reaction (RT-PCR) was performed using either the primer pair (sense, CACCATGGGGGAAAGCAGCACCGGTGCAG; antisense, TGGCTCTGCTTTGCTTTTTGACATCTCC) or, for expression analysis, primers for a gene-specific sequence of the new mouse gene (240 bp of Genbank accession code Q386867; sense, CTCCTTTCCTTTACACCTTCAGG; antisense, ACACCCATAGT-CATTTACATATTTGG) and Superscript One Step RT-PCR mix (Invitrogen, Karlsruhe, Germany). Annealing temperature for the PCR reaction was 56.0°C (35–40 cycles). For control experiments, the enzyme reverse transcriptase was omitted [*Taq* Platinum Polymerase (Invitrogen) was used instead of the RT-PCR enzyme mix]. PCR products were analyzed by ethidium bromide staining after agarose (0.9%) gel electrophoresis. After amplifying the product of one RT-PCR reaction using Platinum Pfx polymerase (Invitrogen), the total coding sequence was cloned into the expression vector pcDNA3.1D/V5-His-TOPO (Invitrogen). The sequence was then identified by cycle sequencing (SequiTherm Exel II DNA sequencing kit; Epicenter Technologies, Madison WI) using a LICOR Gene READIR 4200 sequencer (MWG-Biotech, Ebersberg, Germany). The sequencing was repeated at GATC (Konstanz, Germany) with identical results.

Expression of Recombinant Adenine Receptors in a Mammalian Cell Line. Human 1321N1 astrocytoma cells (European Cell Culture Collection, Salisbury, UK) were cultured at 5% CO₂ and 36.5°C in Dulbecco's modified Eagle medium (Invitrogen) containing GlutaMAX-I (Invitrogen) and 10% fetal bovine serum (Invitrogen). The cells were split once a week by treating with trypsin-EDTA (0.5 g/l; Invitrogen). 1321N1 astrocytoma cells were transfected using the pcDNA3.1 expression vector and Lipofectamine 2000 (Invitrogen) as described by the manufacturer. Cells stably expressing the receptor constructs were selected two days after transfection by culturing in the presence of 800 µg/ml G-418 (Geneticin; Invitrogen). Cells from passages 4 to 20 of the transfected 1321N1 astrocytoma cells were used for further experiments. For some experiments, 1321N1 astrocytoma cells expressing the mouse adenine receptor were pretreated with pertussis toxin 200 ng/ml for approximately 20 h (for control experiments, see *Chemicals*).

Generation of Recombinant Baculoviruses. The sequence encoding for the mouse adenine receptor (without vector-encoded epitopes) was cloned into the pVL1393 baculovirus transfer vector (BD Pharmingen, San Diego, CA) between the BamHI and BglII

restriction sites using standard techniques. Recombinant baculoviruses were generated using the Baculovirus expression vector system from BD Pharmingen, which is based on the method of Guarino and Summers (1986). In brief, 2 × 10⁶ cells were cotransfected with 0.5 µg BaculoGold DNA (Pharmingen, San Diego, CA) and 2 µg of transfer vector DNA containing the mouse adenine receptor gene using the calcium phosphate transfection method. Six days after transfection, the supernatant was harvested. The transfection supernatant was amplified to produce high titer virus stocks. The titer was determined using a modified endpoint dilution method (Reed and Muench, 1938). In brief, in a 24-well plate, 2 × 10⁵ cells/well were infected with virus dilutions from 10⁻² to 10⁻¹¹. After 3 to 5 days, cells were inspected for signs of infection, and the highest effective virus dilution was used for further transfection experiments.

Expression of Recombinant Adenine Receptors in Insect Cells. *Spodoptera frugiperda* (Sf21) cells (gift from J. Höhfeld, Cell Biology, Bonn, Germany) were grown in Insect express protein-free medium (Lonza Verviers SPRL, Verviers, Belgium). Cells were grown at 27°C and subcultured twice a week. Sf21 cells (1.8 × 10⁷) were infected with recombinant baculoviruses at a multiplicity of infection of 1. They were harvested 72 h after infection. Cells were homogenized with 20 strokes in a tight-fitting Dounce homogenizer in 50 mM Tris-HCl buffer, pH 7.4, containing 1 mM phenylmethylsulfonyl fluoride. The cell suspension was centrifuged for 10 min at 1000g, 4°C. The supernatant was then centrifuged at 48,000g, 60 min, 4°C, and the resulting pellet was resuspended in 50 mM Tris-HCl buffer, pH 7.4. The protein concentration was determined by the method of Bradford (1976). Membranes were kept frozen at -80°C until use.

Expression Analysis. 1321N1 astrocytoma cells stably expressing the mouse adenine receptor were cultured on coverslips in Dulbecco's modified Eagle's medium supplemented with 800 µg/ml G-418 for 2 days. Expression of the recombinant adenine receptor was assessed by direct immunofluorescence staining using a FITC-coupled monoclonal antibody against the V5 receptor epitope encoded by the expression vector (1:500; anti V5-FITC; Invitrogen). Expression was then verified by fluorescence microscopy on a Zeiss Axiovert 100 microscope equipped with an oil immersion 100× objective (Zeiss, Jena, Germany), a charge-coupled device camera, and a Polychrome II monochromator (excitation wavelength, 475 nm; exposure time, 1 sec; TILL Photonics, Planeg, Germany).

Analysis of Cellular Cyclic AMP Accumulation. Receptor function was assessed by analyzing the activity of cellular adenylate cyclase activity. For this purpose, nontransfected 1321N1 astrocytoma cells and 1321N1 astrocytoma cells stably expressing the recombinant mouse adenine receptor or the human P2Y₁₂ receptor were cultured on 24-well plates for 2 days. After removal of the culture medium, cells were washed with Hanks' balanced salt solution (containing 20 mM HEPES, pH 7.3) and then incubated with Hanks' balanced salt solution for 2 h at 36.5°C. Cellular cAMP production was then stimulated by the addition of 3 nM isoproterenol at 36.5°C. Solvent (control), adenine, 2-fluoro-adenine, or 2-methylthio-ADP was added together with isoproterenol. The reaction was stopped after 10 min by removal of the reaction buffer followed by the addition of a hot lysis solution (500 µl; 90°C; 4 mM Na₂EDTA, 0.01% Triton X-100; Sigma, Munich, Germany). In some experiments, the receptor antagonists DPCPX and suramin were given 10 min before adenine. The multiwell plates were shaken on ice for 1 h. cAMP levels in the supernatant were then quantified by incubation of an aliquot with cAMP binding protein and [³H]cAMP (cAMP assay; GE Healthcare, Freiburg, Germany) and liquid scintillation counting after removal of the unbound [³H]cAMP by charcoal. cAMP levels per well were calculated by linear regression from a standard curve determined for each experiment. The isoproterenol-induced cAMP production in the presence of agonists was expressed as percentage of the isoproterenol-induced cAMP production in the absence of agonists (percentage of control).

Radioligand Binding Studies. Saturation and competition assays with [^3H]adenine (27 Ci/mmol; GE Healthcare) were carried out essentially as described previously (Gorzalka et al., 2005). In brief, 50 μg of protein (NG108-15 membrane preparations) or 100 μg (Sf21 membrane preparations) was incubated with 10 nM [^3H]adenine (competition assay) in 50 mM Tris-HCl, pH 7.4, in a total volume of 200 μl . Inhibition curves were determined using six to nine different concentrations of adenine or adenine derivative spanning 3 orders of magnitude. For saturation assays, increasing amounts (0.75–150 nM) of [^3H]adenine were incubated with the Sf21 membrane preparations (100 μg) in 50 mM Tris-HCl buffer, pH 7.4, in a total volume of 200 μl . Three separate experiments were performed each in duplicate or triplicate as noted. Nonspecific binding was determined in the presence of 100 μM adenine. Incubations were carried out for 1 h at room temperature and terminated by rapid filtration through GF/B glass fiber filters (Whatman, Dassel, Germany). Filters were washed three times with ice-cold 50 mM Tris-HCl buffer, pH 7.4, 2 ml each. Filter-bound radioactivity was measured by liquid scintillation counting.

Data Analysis and Statistics. Results are presented as means \pm S.E. from n observations. Data were analyzed using Prism 4.03 (Graph Pad Software, San Diego, CA). Differences between means were tested for significance by the t test or (for multiple comparisons with the same control) by an analysis of variance followed by the Bonferroni post test (Prism). $p < 0.05$ was the significance criterion. IC_{50} values (concentrations causing half-maximal inhibition) were determined by fitting data to sigmoidal curves.

Chemicals. The following drugs were used: 7-methyladenine (Acros Organics, Geel, Belgium); adenine, ADP sodium salt, ATP sodium salt, N^6 -benzyladenine, caffeine, 8-cyclopentyl-1,3-dipropylxanthine (DPCPX), N^6 -dimethyladenine, 2-fluoroadenine, 1-methyladenine, 2-methylthio-ADP, 5'-(N -ethylcarboxamido)adenosine (NECA), and pertussis toxin (Sigma); and suramin hexasodium salt (Bayer, Leverkusen, Germany). Stock solutions of drugs were prepared with either distilled water or with dimethyl sulfoxide (N^6 -benzyladenine, N^6 -dimethyladenine, DPCPX, 2-fluoroadenine, 7-methyladenine, 1-methyladenine, NECA). The solvents were added to the buffer used for control experiments. For pertussis toxin control experiments, the solvents present in the purchased solution were added to the medium; final concentrations: 0.05% glycerol (Serva, Heidelberg, Germany), 0.5 mM NaCl, 0.05 mM Trizma base (Sigma), and 0.38 mM glycine (Merck, Darmstadt, Germany).

Results

Cloning of a Novel Receptor for Adenine. Poly A⁺ mRNA was isolated from mouse brain homogenates as well as from the mouse neuroblastoma x rat glioma hybrid cell line NG108-15. RT-PCR with both mRNA preparations and with primers for the coding sequence of the putative mouse ortholog MrgA10 of the rat adenine receptor revealed products of the expected length of approximately 1000 base pairs (for NG108-15 cells, see also Gorzalka et al., 2005). Control reactions without the enzyme reverse transcriptase showed no products confirming that the mRNA preparations contained no genomic DNA. The RT-PCR products were subcloned in the pcDNA3.1 TOPO expression vector. We were surprised to find that sequencing of the subcloned DNA sequence did not reveal the expected sequence of the mouse ortholog of MrgA10 but a distinct sequence encoding for a protein with 332 amino acids, with only 81.6% identity compared with the mouse ortholog MrgA10 and only 76.3% identity compared with the rat adenine receptor (see Fig. 1A). A high degree of variability was found in the predicted transmembrane region 3, and the predicted third extracellular

loop (bold residues in Fig. 1A). The new mouse nucleotide sequence was submitted to the Genbank under accession number DQ386867. This novel gene is located on mouse chromosome 7B3, whereas MrgA10 is localized in the region 7B4 to 7B5 (NCBI database). Additional RT-PCR experiments using specific primers for the new mouse gene (discriminating it from the mouse MrgA10 sequence) revealed products of the expected length of 240 base pairs from poly A⁺ mRNA isolated from mouse brain (B+; marked signal) and mouse spleen (S+, weak signal), but not from mRNA isolated from mouse liver L, or mouse kidney (K; Fig. 1B). This indicates the expression of mRNA for the mouse adenine receptor in neuronal tissues as well as in the spleen. Again, there were no products in reactions without the enzyme reverse transcriptase (“–” in Fig. 1B) in agreement with the absence of genomic DNA in our mRNA preparations.

Expression in Astrocytoma Cells and Functional Analysis. Next, experiments were performed to analyze the pharmacological properties of the novel construct when expressed in mammalian cells. In nontransfected human 1321N1 astrocytoma cells, adenine (0.1 and 1 μM) caused no inhibition in cellular cAMP production in the absence and presence of isoproterenol used to stimulate the cellular cAMP accumulation (data not shown). Moreover, in 1321N1 astrocytoma cells stably expressing the recombinant human P2Y₁₂ receptor, the P2Y₁₂ receptor agonist 2-methylthio-ADP (1 μM) caused the expected inhibition of the isoproterenol (10 nM)-induced increases in cAMP levels (cAMP levels in the presence of 1 μM 2-methylthio-ADP amounted to $52.0 \pm 1.7\%$ of those without 2-methylthio-ADP, control; $n = 3$; $p < 0.05$ versus control, t test), whereas 10 μM adenine again failed to cause a change ($98.9 \pm 6.0\%$ of control; $n = 3$). The lack of any effect of adenine confirms the absence of an endogenous inhibitory receptor for adenine in 1321N1 astrocytoma cells.

1321N1 astrocytoma cells were then transfected with a pcDNA3 vector containing the sequence for the novel adenine receptor and the V5-epitope. Positive clones were selected in the presence of G-418 (see *Materials and Methods*). The expression of the receptor was verified by immunofluorescence staining against the receptor epitope (see *Materials and Methods*). In contrast to nontransfected cells, in 1321N1 astrocytoma cells stably expressing the mouse receptor, adenine (1 nM–1 μM) inhibited the isoproterenol-induced cAMP formation in a concentration-dependent manner with a half-maximal concentration (IC_{50}) of 8 nM (95% confidence interval, 3–23 nM) and a maximal inhibition by approximately 40% (Fig. 2A). Likewise, the analog 2-fluoroadenine caused a concentration-dependent inhibition of cAMP production with an IC_{50} concentration of 15 nM (95% confidence interval, 7–30 nM) and a maximal inhibition by approximately 60% (Fig. 2B). The adenosine A₁ receptor antagonist DPCPX used at a concentration of 1 μM and the P2 receptor antagonist suramin (300 μM) failed to alter the inhibitory effects of adenine in cells expressing the recombinant mouse adenine receptor (Fig. 3). DPCPX and suramin also caused no change in the increases in cellular cAMP levels induced by isoproterenol in the absence of adenine (not shown). Finally, the effect of pretreatment with pertussis toxin was studied in astrocytoma cells expressing the recombinant mouse adenine receptor. Pretreatment with pertussis toxin (200 ng/ml) for 20 h did not alter the isoproterenol-induced cAMP formation (see legend to Fig. 4) but abolished the inhibitory effect of

adenine observed in cells pretreated with the solvent used for pertussis toxin (Fig. 4).

Expression of the Novel Receptor in Sf21 Cells and Analysis of Its Binding Properties. The novel mouse adenine receptor was subsequently expressed in Sf21 insect cells using the Baculovirus expression system. The adenine receptor sequence was inserted into the Baculovirus transfer vector pV11393. By cotransfecting the vector and linearized

Baculovirus DNA into Sf21 cells, recombinant Baculoviruses were generated (see *Materials and Methods*). After amplifying the viruses, Sf21 cells were infected for protein expression. Membrane preparations of noninfected Sf21 cells showed no specific binding for [³H]adenine (10 nM) (Fig. 5). In contrast, a specific binding site for [³H]adenine could be detected on membranes of Sf21 cells expressing the mouse adenine receptor (Fig. 5). Saturation experiments with [³H]-

A

Novel seq (001)	MGES STGAGF	LALNISASTM	ALTTTNPMDE	TIPG STSIKI
mMrgA10 (001)	MGESSTGAGF	LALNTSASSM	APSTTNLMDE	TIPGSIDIST
rAdenin-r (001)	MGESFTGTGF	INLNTSASTI	AVTTTNPMDK	TIPGSFNGRT
	N-terminus			
Novel seq (041)	LIPNL MIIIF	GLVGL TGNAI	VFWLL GFHLR	RNAFSVYILN
mMrgA10 (041)	LIPHLMIIIF	GLVGMTGNAI	VFWLLGFRLR	KNAFSVYILN
rAdenin-r (041)	LIPNLLIIIS	GLVGLIGNAM	VFWLLGFRLA	RNAFSVYILN
		TM1	IL1	TM2
Novel seq (081)	LALADFLFLL	CRIIASTQKL	LFSSPNITF	LLCLYTFRVI
mMrgA10 (081)	LALADFLFLL	CHIIASTLFL	LKVSYPNIIF	RKCFFSVMLV
rAdenin-r (081)	LALADFLFLL	CHIIDSTLLL	LKFSYPNIIF	LPCFNTVMMV
	TM2	EL1		TM3
Novel seq (121)	LYIAGLSMLT	AIS IER CLSV	LCPIWY RCHR	PEHTSTVMCA
mMrgA10 (121)	LYIAGLSILS	AIGTERCLSV	LCPIWYRCHC	PEHTSTVTCA
rAdenin-r (121)	<u>PYIAGLSMLS</u>	AISTERCLSV	VCPIWYRCHR	<u>PKHTSTVMCS</u>
	TM3	IL2		TM4
Novel seq (161)	AIWVLSLLIC	ILNRY FSGFL	DTKY VNDYGC	MASNFFNAAY
mMrgA10 (161)	MIWVLSLLIS	ILNKYFCVFL	DTKYVNDYGC	MASNFFTAAY
rAdenin-r (161)	AIWVLSLLIC	ILNRYFCGFL	DTKYEKDNRC	LASNFFTAAC
	TM4	EL2		TM5
Novel seq (201)	LMFLFVVLCV	SSLALL ARLF	CGTGRM KLTR	LYVTIM LTL
mMrgA10 (201)	LMFLFVVLCV	SSLALLARLF	CGAGRMKLTR	LYVTIMLTL
rAdenin-r (201)	LIFLFVVLCV	SSLALLVRSF	CGAGRMKLTR	LYATIMLTVL
	TM5	IL3		TM6
Novel seq (241)	VFLLCGL PCG	LYWFLLFWIK	NGFAVDFNF	YLASTVLSAI
mMrgA10 (241)	VFLLCGLPCG	IYWFLLSKIK	NVFIVDFDSL	FMASSVLTAL
rAdenin-r (241)	VFLLCGLPFG	IHWFLLIWIK	IDYGKFAYGL	YLAALVLTAV
	TM6	EL3		TM7
Novel seq (281)	NSSAN P IIYF	FVGSFR HRLK	HQTLKM VLQS	ALQDTP ETAE
mMrgA10 (281)	NSCANP IIYF	FVGSFRHRLQ	HQTLKMVIQS	ALQDIP ETPE
rAdenin-r (281)	NSCANP IIYF	FVGSFRH-QK	HQTLKMVLQR	ALQDTP ETAE
	TM7		C-terminus	
Novel seq (321)	NM VEMSK SKA	EP		
mMrgA10 (321)	NIVEMSK SKA	EP		
rAdenin-r (320)	NTVEMSS SKV	EP		
		C-terminus		

B



B+ B- L+ L- K+ K- S+ S-

Fig. 1. A, alignment of the predicted amino acid sequences of the novel adenine receptor (Novel seq, mouse adenine receptor, present study, Genbank nucleotide sequence accession number DQ386867; see also XM_001003331), the mouse ortholog of MrgA10 (mMrgA10; Genbank nucleotide sequence accession no. XM_195647, please note that the total sequence of XM_195647 encodes for a protein with an extended N terminus) and the rat adenine receptor (rAdenin-r; Genbank nucleotide sequence accession no. AJ311952). Residues that differ in the mouse and the rat adenine receptor are shown in bold. The seven predicted transmembrane regions (TM) are indicated by lines (IL, intracellular loop; EL, extracellular loop). **B**, RT-PCR experiment indicating the expression of mRNA for the mouse receptor for adenine in tissues using primers specific for amplification of the sequence DQ386867 and poly A⁺ mRNA isolated from total brain (B), liver (L), kidney (K), and spleen (S). The figure shows RT-PCR products (+) stained by ethidium bromide after agarose gel electrophoresis. Control experiments without reverse transcriptase (-) revealed no products. One of three independent experiments.

adenine revealed a single high-affinity binding site with a K_D value of 113 ± 17 nM, and a B_{max} value of 1.98 ± 0.39 pmol/mg protein was determined ($n = 3$) (Fig. 6).

Selected adenine derivatives, as well as compounds known to activate (NECA) or block (caffeine) adenosine receptors or to activate P2 receptor subtypes (ADP, ATP), were investigated in competition assays at the novel mouse adenine receptor expressed in Sf21 insect cell membranes. Affinity data were compared with those for the adenine receptor expressed in NG108-15 mouse neuroblastoma x rat glioma cell membranes and those expressed in rat brain cortical membrane preparations (Fig. 7 and Table 1). Unsubstituted adenine showed the highest affinity in all three test systems exhibiting K_i values in the nanomolar range ($IC_{50} = 68.5$ nM for the mouse receptor in Sf21 cells, 54.9 nM in NG108-15 cells, $K_i = 29.9$ nM in rat brain cortex). The rank order of potency for the mouse receptor expressed in Sf21 cells was adenine > 2-fluoro-adenine > 7-methyladenine > 1-methyladenine >> N^6 -benzyladenine \approx N^6 -dimethyladenine. The same rank orders of potency were observed for NG108-15 cell membranes (Fig. 7B) and rat brain cortical membranes (Table 1). All of the investigated P1 (adenosine) and P2 receptor




Fig. 2. Inhibition of isoproterenol-stimulated cAMP accumulation by adenine (A, $IC_{50} = 8$ nM; 95% confidence interval, 3–23 nM) and 2-fluoro-adenine (B, $IC_{50} = 15$ nM; 95% confidence interval, 7–30 nM) in 1321N1 astrocytoma cells stably expressing the novel (mouse) adenine receptor. Cellular cAMP production was increased by addition of isoproterenol 3 nM for 10 min at 36.5°C ; data are given as percentages of the mean increases in cellular cAMP levels in the presence of isoproterenol alone (percentage of control; average increase in cAMP levels by 15.3 ± 1.3 pmol cAMP per well; $n = 17$). Means \pm S.E. from 5 to 21 experiments. ** indicates significant differences from corresponding control ($p < 0.01$; ANOVA followed by the Bonferroni post test).

ligands showed no or only moderate affinity for adenine binding sites in all three membrane preparations (Table 1).

Discussion

An orphan rat GPCR has previously been shown to operate as a receptor for the nucleobase adenine (Bender et al., 2002) indicating that in addition to P1 receptors for adenosine (Fredholm et al., 2001; Yan et al., 2003) and P2Y receptors for extracellular nucleotides (Brunschweiler and Müller, 2006; von Kügelgen, 2006), a third group of GPCRs exists for




Fig. 3. Inhibition of isoproterenol-stimulated cAMP accumulation and interaction with $1 \mu\text{M}$ DPCPX and $300 \mu\text{M}$ suramin in 1321N1 astrocytoma cells stably expressing the novel (mouse) adenine receptor. Antagonists or solvent were added 10 min before 10 nM adenine. Data are given as percentages of the mean increases in cellular cAMP levels in the presence of isoproterenol alone (percentage of control). Means \pm S.E. from 5 to 21 experiments. * indicates significant differences from corresponding control ($p < 0.05$; ANOVA followed by the Bonferroni post test). For further details, see legend to Fig. 2.




Fig. 4. Effect of the pretreatment with pertussis toxin (PTX) on the adenine-induced inhibition of isoproterenol-stimulated cAMP accumulation in 1321N1 astrocytoma cells stably expressing the novel (mouse) adenine receptor. Cells were pretreated with 200 ng/ml PTX or its solvent for 20 h. Data are given as percentages of the mean increases in cellular cAMP levels in the presence of isoproterenol alone (percentage of control; pretreated with solvent: average increase in cAMP levels by 10.9 ± 0.4 pmol of cAMP per well; $n = 5$; pretreated with PTX: 9.8 ± 0.7 pmol cAMP per well; $n = 5$). Means \pm S.E. from five to seven experiments. **, significant difference from corresponding control (no adenine; $p < 0.01$); ##, significant difference from corresponding value obtained with cells pretreated with solvent ($p < 0.01$; ANOVA followed by the Bonferroni post test). For further details see legend to Fig. 2.

these chemically related compounds (tentatively termed P0 receptors; Brunschweiler and Müller, 2006). Human cells also seem to express receptors for adenine (Gorzalka et al., 2005). The human sequence encoding an adenine receptor, however, has yet to be identified. The present study now demonstrates that a novel sequence encodes for an additional GPCR for adenine. This receptor shares only 76% identical amino acid residues with the rat receptor, suggesting that these proteins are not species homologs but distinct members of a family of GPCRs for adenine. This view is underlined by the fact that the predicted transmembrane regions 3 of the novel receptor and the rat adenine receptor differ markedly (see TM3 in Fig. 1), whereas species homologs are known to contain conserved transmembrane regions. Moreover, the tissue distribution of the novel receptor and the rat adenine receptor differ; the latter is predominantly expressed in dorsal root ganglia (Bender et al., 2002), whereas we detected mRNA for the novel sequence in the brain and the spleen, suggesting a broader expression. As discussed below, there are also differences in pharmacological properties. A related mouse DNA sequence encodes for the mMrgA10 orphan receptor (Bender et al., 2002; see Choi and Lahn, 2003; Zylka et al., 2003). It is not known yet whether mMrgA10 also operates as a receptor for adenine. Neither for the rat adenine receptor (Bender et al., 2002) nor for the new sequence could a closely related human ortholog be identified in the genome database. A search for conserved regions of both sequences,




Fig. 5. [^3H]Adenine binding (10 nM) to membrane preparations (100 μg of protein) of noninfected (Sf21) and infected (Sf21 pVI-mAde; novel mouse receptor for adenine) Sf21 cells as well as to glass-fiber filters without protein (samples without protein). Nonspecific binding was determined in the presence of 100 μM unlabeled adenine. Data are means \pm S.E. from three (Sf21 pVI-mAde), four (samples without protein) or five (Sf21) independent experiments performed in triplicate; ns = not significant, *** indicates significant difference ($p < 0.001$; t test).




Fig. 6. Saturation curve of [^3H]adenine binding to membranes prepared from Sf21 insect cells expressing the novel (mouse) adenine receptor. A K_D value of 113 ± 17 nM and a B_{max} value of 1.98 ± 0.39 pmol/mg protein was determined ($n = 3$). Experiments were performed as described under *Materials and Methods*. The result shown represents the mean \pm S.E. of three independent experiments, each performed in duplicate.

however, may facilitate the identification of the human sequence.

For functional experiments, human 1321N1 astrocytoma cells were used. Nontransfected 1321N1 astrocytoma cells do not respond to adenine with a decrease in cAMP formation (present study). Therefore, they are suitable for a pharmacological characterization of a recombinant inhibitory receptor for adenine (despite the fact that the cells possess binding sites for adenine; Gorzalka et al., 2005). In a previous study by Bender et al. (2002), a clone of Chinese hamster ovary (CHO) cells without endogenous adenine receptors had been used for recombinant expression of the rat adenine receptor. However, the clones of CHO cells studied in our laboratory (CHO Flp-In cells and CHO K1 cells) have a high density of adenine binding sites (S. Gorzalka and C. E. Müller, unpublished observations) and possess endogenous receptors for adenine modulating intracellular cAMP levels (K. Hoffmann and I. von Kügelgen, unpublished results).

The novel recombinant mouse adenine receptor shows pharmacological properties similar but not identical to those of the adenine receptor endogenously expressed in the mouse neuroblastoma x rat glioma hybrid cell line NG108-15 [which expresses the same mouse sequence (Gorzalka et al., 2005) and, possibly, an additional subtype]. Both the native and the recombinant receptor couple to inhibition of adenylate cyclase activity via pertussis toxin-sensitive G proteins. Adenine acted as an agonist at the native (Gorzalka et al., 2005) and the recombinant receptor (present study) with half-max-




Fig. 7. Competition curves for adenine, 2-fluoroadenine, and 7-methyladenine versus 10 nM [^3H]adenine obtained with a membrane preparation of Sf21 insect cells recombinantly expressing the novel (mouse) adenine receptor (A) or membranes from NG108-15 cells, natively expressing the novel (mouse) adenine receptor (B). The IC_{50} values were calculated by nonlinear regression function for one-site competition (IC_{50} values: Sf21, adenine = 68.5 ± 9.8 nM ($n = 5$); 2-fluoroadenine = 1.47 ± 0.14 nM ($n = 2$), and 7-methyladenine = 5.76 ± 1.02 nM ($n = 2$); NG: adenine = 0.154 ± 0.056 nM ($n = 2$), 2-fluoroadenine = 0.304 ± 0.097 nM ($n = 3$), and 7-methyladenine = 1.45 ± 0.44 nM ($n = 2$)). Data are means \pm S.E. from two to five independent experiments performed in duplicate as indicated above.

TABLE 1

Comparison of affinities of adenine and selected compounds for the novel (mouse) adenine receptor and the native rat adenine receptor determined in radioligand binding studies at membrane preparations (NECA, 5'-(*N*-ethylcarboxamido)adenosine)

Results are from three independent experiments performed in triplicate, unless otherwise noted. Values in parentheses are percentage inhibition at 100 μ M; all other values are IC_{50} or K_i values.











Compound	Structure	Mouse Recombinant Receptor Expressed in Sf21 Cells ^a	NG108–15 Cells ^a	Rat Brain Cortex ^b
Adenine		0.0685 ± 0.0098^c	$0.0549 \pm 0.0422^{d,e}$ $0.154 \pm 0.056^{d,e}$	0.0299 ± 0.0034
1-Methyladenine		(62 ± 7)	(60 ± 4)	29.3 ± 9.3
2-Fluoroadenine		1.47 ± 0.14^d	0.304 ± 0.097	0.622 ± 0.144
<i>N</i> ⁶ -Benzyladenine		$(12 \pm 12)^d$	$(8 \pm 12)^d$	(30 ± 3)
<i>N</i> ⁶ -Dimethyladenine		$(0 \pm 8)^d$	$(0 \pm 6)^d$	(11 ± 2)
7-Methyladenine		5.76 ± 1.02^d	1.45 ± 0.44^d	4.13 ± 1.08
NECA		$(28 \pm 1)^d$	$(7 \pm 1)^d$	(10 ± 2)
ADP		$(28 \pm 3)^d$	$(50 \pm 5)^d$	(51 ± 3)

TABLE 1 (Continued)

Compound	Structure	Mouse Recombinant Receptor Expressed in Sf21 Cells ^a	NG108-15 Cells ^a	Rat Brain Cortex ^b
ATP		(14 ± 1) ^d	(61 ± 4) ^d	(39 ± 7)
Caffeine		(1 ± 1) ^d	(0.4 ± 1) ^d	(2 ± 1)

^a IC₅₀ values versus 10 nM [³H]adenine.

^b K_i values from Gorzalka et al. (2005).

^c Results from five independent experiments performed in duplicate.

^d Results from two independent experiments performed in duplicate.

^e IC₅₀ value from Gorzalka et al. (2005).

imal concentrations of 21 nM (NG108-15 cell membranes) and 8 nM (recombinant receptors), respectively. Adenine has been shown to activate the rat adenine receptor with a similar potency (3 nM; Bender et al., 2002). In addition to adenine, the adenine derivative 2-fluoroadenine was active at the native receptor in NG108-15 cells (Gorzalka et al., 2005) and at the recombinant receptor (Fig. 2B).

In contrast to the results obtained at the recombinant receptor expressed in astrocytoma cells (present study) and at NG108-15 membranes (Gorzalka et al., 2005), a higher IC₅₀ value for the adenine-induced inhibition of forskolin-stimulated cAMP production had been determined in intact NG108-15 cells (IC₅₀ 2.54 μM; Gorzalka et al., 2005). The difference is likely to be caused by an uptake mechanism for adenine in intact neuroblastoma NG108-15 cells, by which adenine is removed from the extracellular surface. In fact, in a recent study, only rat cortical neurons but not astrocytes were found to take up [³H]adenine, indicating that astrocytes do not express specific transport systems for adenine (Nagai et al., 2006).

An action of adenine via adenosine receptors was excluded by the lack of any interaction with the adenosine A₁ receptor antagonist DPCPX, which completely blocks adenosine A₁ and A_{2B} receptors and at least partly blocks A_{2A} and A₃ receptors at the concentration used (1 μM; Lohse et al., 1987; Müller, 1996). Moreover, adenine does not act on P2 receptors, as shown by the lack of interaction with suramin (acting as an antagonist at P2Y₁, P2Y₂, P2Y₆, P2Y₁₁, P2Y₁₂, and P2Y₁₃ receptors; Table 3 of von Kügelgen, 2006). Hence, these results confirm the idea of a group of GPCRs for the nucleobase adenine distinct from the known adenosine and P2 receptors.

In addition to functional studies, we were interested in obtaining a suitable system for radioligand binding assays. NG108-15 cell membranes natively expressing the mouse adenine receptor had previously been shown to exhibit specific binding of [³H]adenine (Gorzalka et al., 2005). However, this cell line may express more than one adenine receptor subtype or other specific binding sites for [³H]adenine. Although we tested a series of cell lines, including CHO, human

embryonic kidney, and 1321N1 astrocytoma cells, typically used for heterologous recombinant expression of GPCRs, we could not identify any mammalian cell line that did not natively express a specific binding site for [³H]adenine. Therefore, we turned our attention to nonmammalian cells and found that Sf21 insect cell membranes showed no specific binding of [³H]adenine. After infection with recombinant baculovirus containing the sequence for the novel adenine receptor, specific binding of [³H]adenine was detected in these Sf21 insect cell membranes (Fig. 5). In homologous competition assays of adenine versus [³H]adenine, concentration-dependent inhibition of [³H]adenine binding was observed (Fig. 7). Because Sf21 cell membranes express no high-affinity adenine binding proteins, they represent an ideal expression system for adenine receptor proteins for binding assays. It has been suggested that [³H]adenine may bind to glass fiber filters in a specific manner (Ye et al., 2006; but see Schiedel et al., 2007, for bacterial proteins binding adenine). However, we observed no specific binding of [³H]adenine to glass fiber filters, only nonspecific binding in the absence of Sf21 membranes, which was somewhat lower than the nonspecific binding observed in the additional presence of Sf21 membranes (Fig. 5). The specific binding of [³H]adenine to Sf21 insect cell membranes expressing the novel adenine receptor was clearly protein-dependent and increased with higher protein concentrations (data not shown).

In saturation experiments at Sf21 cells expressing the novel adenine receptor, a K_D value of 113 nM was determined for [³H]adenine. Thus, the K_D value was 4-fold higher than that determined for the rat adenine receptor in brain cortical membranes (27.2 nM; Gorzalka et al., 2005) but still in the same concentration range. A comparison of affinities of adenine and selected derivatives for the novel receptor showed that affinities were similar to but not identical with those determined at the native receptors in NG108-15 cells and the rat brain cortex (Table 1). The rank order of potency was: adenine > 2-fluoroadenine > 7-methyladenine > 1-methyladenine >> N⁶-benzyladenine = N⁶-dimethyladenine. Slight differences could be observed. For example, 2-fluoroadenine

and 7-methyladenine showed a 4- to 5-fold lower affinity for the recombinant receptor than for the native receptor (Fig. 7 and Table 1). These differences are in agreement with the expression of different receptor subtypes in NG108-15 cells. In addition, different membrane environments may contribute to the differences by changing receptor conformation (Chachisvilis et al., 2006). In radioligand binding studies, the adenosine receptor agonist NECA and the antagonist caffeine showed no affinity for the mouse adenine receptor (Table 1). The same was observed for the P2 receptor agonists ATP and ADP. IC_{50} values determined in radioligand binding studies were generally severalfold higher than IC_{50} values obtained in functional cAMP assays because of the amplification of the signal mediated by a GPCR.

In conclusion, we have identified, cloned, and pharmacologically characterized a novel receptor for the purine nucleobase adenine from mouse brain. The low sequence identity with the previously described rat adenine receptor, particularly in the predicted transmembrane region 3, as well as differences in pharmacology, suggests that the new receptor is not a species homolog but rather a distinct receptor subtype. Because human cells also possess specific binding sites for adenine, the existence of a receptor for adenine in human tissues seems very likely. However, a human sequence encoding for an adenine receptor has yet to be identified. The rat adenine receptor discovered by Bender et al. (2002) and the novel adenine receptor described in the present study seem to be members of a new, possibly larger family of receptors for purine nucleobases. The comparison of the sequences may facilitate the identification of the human sequence.

Acknowledgments

We thank Petra Spitzlei for skilful technical assistance and Prof. Dr. Jörg Höhfeld for Sf21 insect cells.

References

- Bender E, Buist A, Jurzak M, Langlois X, Baggerman G, Verhasselt P, Ercken M, Guo H-Q, Wintmolders C, Van den Wyngaert I, Van Oers I, Schoofs L, and Luyten W (2002) Characterization of an orphan G protein-coupled receptor localized in the dorsal root ganglia reveals adenine as a signaling molecule. *Proc Natl Acad Sci USA* **99**:8573–8578.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**:248–254.
- Brunschweiler A and Müller CE (2006) P2 receptors activated by uracil nucleotides—an update. *Curr Med Chem* **13**:289–312.
- Chachisvilis M, Zhang YL, Frangos JA (2006) G protein-coupled receptors sense fluid shear stress in endothelial cells. *Proc Natl Acad Sci U S A* **103**:15463–15468.
- Choi SS and Lahn BT (2003) Adaptive evolution of MRG, a neuron-specific gene family implicated in nociception. *Genome Res* **13**:2252–2259.
- Fredholm BB, IJzerman AP, Jacobson KA, Klotz KN, and Linden J (2001) International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev* **53**:527–552.
- Gorzalka S, Vittori S, Volpini R, Cristalli G, von Kügelgen I, and Müller CE (2005) Evidence for the functional expression and pharmacological characterization of adenine receptors in native cells and tissues. *Mol Pharmacol* **67**:955–964.
- Guarino LA and Summers MD (1986) Functional mapping of a trans-activating gene required for expression of a baculovirus delayed-early gene. *J Virol* **57**:563–571.
- Hamprecht B (1977) Structural, electrophysical, biochemical, and pharmacological properties of neuroblastoma-glioma cell hybrids in cell culture. *Int Rev Cytol* **49**:99–170.
- Kaulich M, Qurishi R, and Müller CE (2003) Extracellular metabolism of nucleotides in neuroblastoma x glioma NG108-15 cells determined by capillary electrophoresis. *Cell Mol Neurobiol* **23**:349–364.
- Lohse MJ, Klotz KN, Lindenborn-Fotinos J, Reddington M, Schwabe U, and Olsson RA (1987) 8-Cyclopentyl-1,3-dipropylxanthine (DPCPX)—a selective high affinity antagonist radioligand for A_1 adenosine receptors. *Naunyn Schmiedebergs Arch Pharmacol* **336**:204–210.
- Matthews EA and Dickenson AH (2004) Effects of spinally administered adenine on dorsal horn neuronal responses in a rat model of inflammation. *Neurosci Lett* **356**:211–214.
- Müller CE (1996) Adenosine receptor antagonists: structures and potential therapeutic applications. *Curr Pharm Des* **2**:501–530.
- Nagai K, Nagasawa K, Matsunaga R, Yamaji M, and Fujimoto S (2006) Novel Na^+ -independent and adenine-specific transport system for adenine in primary cultured rat cortical neurons. *Neurosci Lett* **407**:244–248.
- Reed LJ and Muench H (1938) A simple method of estimating fifty percent endpoints. *Am J Hygiene* **27**:493–497.
- Schiedel AC, Meyer H, Alsdorf B, Gorzalka S, Brüssel H, and Müller CE (2007) [3H]Adenine is a suitable radioligand for the labelling of G protein-coupled adenine receptors but shows high affinity to bacterial contaminations in buffer solutions. *Purinergic Signalling* **3**:347–358.
- Slominska EM, Szolkiewicz A, Smolenski RT, Rutkowski B, and Swierczynski (2002) High plasma adenine concentration in chronic renal failure and its relation to erythrocyte ATP. *Nephron* **91**:286–291.
- von Kügelgen I (2006) Pharmacological profiles of cloned mammalian P2Y-receptor subtypes. *Pharmacol Ther* **110**:415–432.
- von Kügelgen I, Schiedel AC, Hoffmann K, Alsdorf BBA, Abdelrahman A, and Müller CE (2007) Cloning and functional expression of a novel G_i protein-coupled receptor for adenine from mouse brain. *Naunyn Schmiedebergs Arch Pharmacol* **375** (Suppl 1):27.
- Watanabe S, Ikekita M, and Nakata H (2005) Identification of specific [3H]adenine-binding sites in rat brain membranes. *J Biochem* **137**:323–329.
- Watanabe S, Yoshimi Y, and Ikekita M (2003) Neuroprotective effect of adenine on Purkinje cell survival in rat cerebellar primary cultures. *J Neurosci Res* **74**:754–759.
- Wengert M, Adão-Novaes J, Assaife-Lopes N, Leão-Ferreira LR, and Caruso-Neves C (2007) Adenine-induced inhibition of Na^+ -ATPase activity: evidence for involvement of the G_i protein-coupled receptor in the cAMP signalling pathway. *Arch Biochem Biophys* **467**:261–267.
- Yan L, Burbiel JC, Maass A, and Müller CE (2003) Adenosine receptor agonists: from basic medicinal chemistry to clinical development. *Expert Opin Emerg Drugs* **8**:537–576.
- Ye K, Mulder-Krieger T, Beukers MW, and IJzerman AP (2006) [3H]Adenine's high filter binding precludes its use as a radioligand for adenine receptors. *Purinergic Signalling* **2**:71.
- Yoshimi Y, Watanabe S, Shinomiya T, Makino A, Toyoda M, and Ikekita M (2003) Nucleobase adenine as a trophic factor acting on Purkinje cells. *Brain Res* **991**:113–122.
- Zylka MJ, Dong X, Southwell AL, and Anderson DJ (2003) Atypical expansion in mice of the sensory neuron-specific Mrg G protein-coupled receptor family. *Proc Natl Acad Sci USA* **100**:10043–10048.

Address correspondence to: Ivar von Kügelgen, Department of Pharmacology and Toxicology, University of Bonn, D-53113 Bonn, Germany. E-mail: kugelgen@uni-bonn.de