

Activation of a Mitochondrial ATPase Gene Induces Abnormal Seed Development in *Arabidopsis*

Kon Baek^{1,3}, Pil Joon Seo^{1,3}, and Chung-Mo Park^{1,2,*}

The ATPases associated with various cellular activities (AAA) proteins are widespread in living organisms. Some of the AAA-type ATPases possess metalloprotease activities. Other members constitute the 26S proteasome complexes. In recent years, a few AAA members have been implicated in vesicle-mediated secretion, membrane fusion, cellular organelle biogenesis, and hypersensitive responses (HR) in plants. However, the physiological roles and biochemical activities of plant AAA proteins have not yet been defined at the molecular level, and regulatory mechanisms underlying their functions are largely unknown. In this study, we showed that overexpression of an *Arabidopsis* gene encoding a mitochondrial AAA protein, ATPase-in-Seed-Development (ASD), induces morphological and anatomical defects in seed maturation. The ASD gene is expressed at a high level during the seed maturation process and in mature seeds but is repressed rapidly in germinating seeds. Transgenic plants overexpressing the ASD gene are morphologically normal. However, seed formation is severely disrupted in the transgenic plants. The ASD gene is induced by abiotic stresses, such as low temperatures and high salinity, in an abscisic acid (ABA)-dependent manner. The ASD protein possesses ATPase activity and is localized into the mitochondria. Our observations suggest that ASD may play a role in seed maturation by influencing mitochondrial function under abiotic stress.

INTRODUCTION

The life cycle of higher plants is not continuous but is interrupted by developmental arrests (Goldberg et al., 1989; Kermode, 1990). A mature embryo undergoes a quiescent, desiccated stage. The discontinuous development mediated by seed maturation confers an evolutionary advantage that enables plants to cope with unfavorable environmental conditions and facilitates dispersal of the plant species (Goldberg et al., 1989; Holdsworth et al., 2008; West et al., 1994).

Seed development is divided into 3 distinct phases (Gutierrez et al., 2007; Parcy et al., 1994; West and Harada, 1993). The first developmental phase is characterized by cell division

and tissue differentiation. Embryogenesis is initiated by double fertilization. While the zygote divides asymmetrically to form the embryo proper and the suspensor, the endosperm constantly proliferates (Natesh and Rau, 1984; West and Harada, 1993). In the second developmental phase, which is also called the maturation phase, the growth of the embryo and cell cycle activities are arrested, and storage molecules, such as proteins, carbohydrates, and lipids, accumulate in the embryo, particularly in the cotyledons (Goldberg et al., 1989; Huang et al., 1992; Raz et al., 2001). The seeds finally become dehydrated in the third developmental phase, designated the late maturation phase. Mature seeds in the quiescent state are tolerant to desiccation (Crouch, 1987; Harada et al., 1988; Kermode, 1990; McCarty and Carson, 1991).

ABA is a central phytohormone mediating seed maturation and development. The endogenous content of ABA is low during early embryogenesis, but the ABA level is gradually elevated during the first half of seed maturation (Gutierrez et al., 2007). After seed maturation, the ABA content decreases. During the maturation phase, ABA inhibits precocious germination and vivipary, thereby rendering the seed dormancy (Finch-Savage and Leubner-Metzger, 2006; Nambara and Marion-Poll, 2003). In addition, ABA promotes the accumulation of seed storage compounds, such as carbohydrates, proteins, and oils, which would be required for seed germination and seedling growth prior to becoming an autotroph (Verdier and Thompson, 2008; Wobus and Weber, 1999). It also confers desiccation tolerance on mature seeds. The induction of late embryogenesis abundant (LEA) proteins by ABA is involved in the desiccation tolerance response (Finkelstein and Somerville, 1990; Keith et al., 1994; Meinke et al., 1994).

LEAFY COTYLEDON 1 (LEC1), LEC2, FUSCA 3 (FUS3), and ABA INSENSITIVE 3 (ABI3) are known to be key transcriptional regulators of seed maturation (Keith et al., 1994; Luerksen et al., 1998; Meinke et al., 1994; Parcy et al., 1997; Stone et al., 2001; West et al., 1994). LEC1, LEC2, and FUS3 are involved in the determination of embryonic cell fates during embryogenesis (Gutierrez et al., 2007; Verdier and Thompson, 2008; Wobus and Weber, 1999). These transcription factors, together with ABI3, also regulate initiation and maintenance of the maturation phase (Kagaya et al., 2005; Verdier and Thompson, 2008; Wobus and Weber, 1999). Consequently, seeds

¹Department of Chemistry, Seoul National University, Seoul 151-742, Korea, ²Plant Genomics and Breeding Institute, Seoul National University, Seoul 151-742, Korea, ³These authors contributed equally to this work.

*Correspondence: cmpark@snu.ac.kr

having mutations in these transcription factors are less resistant to desiccation, and accumulation of storage compounds is reduced in the mutant seeds (Finkelstein and Somerville, 1990; Parcy et al., 1994; West et al., 1994). However, these transcription factor genes are differentially regulated during the developmental stages, and phenotypes of the mutants are not identical, supporting that these transcription factors play both redundant and independent roles (Parcy et al., 1997; Wobus and Weber, 1999).

The ATPases associated with various cellular activities (AAA) family members play roles in diverse cellular activities (Hanson and Whiteheart, 2005). They contain a highly conserved AAA domain consisting of Walker A and B motifs and a second-region-of-homology (SRH) domain (Karata et al., 1999). The Walker A motif, which is known as the P-loop, binds to phosphates in NTP. The Walker B motif is associated with Mg^{2+} . These 2 motifs are cooperatively involved in NTP hydrolysis (Hanson and Whiteheart, 2005). The SRH domain contributes to the maintenance of ATPase activity. Mutations to the conserved amino acid residues significantly decrease ATPase activity (Karata et al., 1999). The SRH domain is characteristic of the AAA protein family. It does not exist in the Walker-type ATPases, which have only 2 consensus motifs, Walker A and B (Karata et al., 1999).

The AAA-type ATPases are widely conserved in archaeobacteria, prokaryotes, and eukaryotes, suggesting that they play a critical role in cellular activities. They are involved in various cellular activities and can thus be divided into several subfamilies according to their biochemical activities and physiological functions, including proteolytic activity, proteasome functions, vesicle-mediated secretion, membrane fusion, peroxisome biogenesis, and mitochondrial functions (Karata et al., 1999; Lupas and Martin, 2002; Patel and Latterich, 1998). More than 60 AAA-type ATPase genes have been identified in the *Arabidopsis* genome (Sugimoto et al., 2004). They regulate diverse aspects of cellular function, such as protein degradation (Lindahl et al., 1996), 26S proteasome activity (Fu et al., 1999), vesicle trafficking (Rancour et al., 2002), peroxisome biogenesis (Olsen, 1998), and hypersensitive responses in plants (Sugimoto et al., 2004).

In this study, we investigated an *Arabidopsis* AAA-type ATPase gene, *ATPase-in-Seed-Development (ASD)*, which is highly expressed in seeds. The expression pattern of the *ASD* gene correlated with the seed maturation process. Transgenic plants overexpressing the *ASD* gene (35S:*ASD*) showed abnormal seed development. The *ASD* gene was induced by abiotic stresses in an ABA-dependent manner. Subcellular localization assays and biochemical activity assays revealed that the *ASD* protein is a mitochondrial ATPase. We propose that the *ASD* protein is involved in stress regulation of seed maturation.

MATERIALS AND METHODS

Plant materials, growth conditions, and *Arabidopsis* transformation

All *Arabidopsis thaliana* lines used were in the Columbia background (Col-0). Plants were grown in a controlled culture room set at 22°C with a relative humidity of 55% under long days (16-h light/8-h dark) with white light illumination (120 μ mol photons/ m^2 s) provided by fluorescent FLR40D/A tubes (Osram, Korea). The *ASD*-deficient *asd-1* mutant (SALK-026454) was obtained from a T-DNA insertional mutant pool deposited into the *Arabidopsis* Biological Resource Center (ABRC, Ohio State University). Absence of gene expression in the mutants was verified

by reverse-transcriptase polymerase chain reaction (RT-PCR) before use.

To produce transgenic plants overexpressing the *ASD* gene, a full-size *ASD* cDNA was subcloned into the binary pH2GW7 vector under the control of the Cauliflower Mosaic Virus (CaMV) 35S promoter. *Agrobacterium*-mediated *Arabidopsis* transformation was carried out according to a modified floral dip method (Clough and Bent, 1998).

Subcellular localization

The *ASD* cDNA sequence was subcloned into the *Xba*I/*Bam*HI-digested 326-RFP expression vector (Lee et al., 2001). The expression construct was transformed into *Arabidopsis* protoplasts by a polyethylene glycol-calcium transfection method (Yoo et al., 2007). The mt-yk CD3-989 binary plasmid (Nelson et al., 2007) was used as a mitochondria-localized marker. Subcellular distribution of the *ASD* protein was visualized by differential interference contrast microscopy (DIC) and fluorescence microscopy using the Zeiss LSM510 confocal microscope (Carl Zeiss, MicroImaging GmbH, Germany).

Transcript level analysis

Quantitative real-time RT-PCR (qRT-PCR) was employed for measuring transcript levels. RNA sample preparation, reverse transcription, and quantitative PCR were carried out on the basis of the rules that have been proposed recently to ensure reproducible and accurate measurements (Udvardi et al., 2008). Extraction of total RNA from appropriate plant materials and RT-PCR conditions have been described previously (Kim et al., 2006). The RNA samples were pretreated extensively with an RNase-free DNase to eliminate any contaminating genomic DNA before use.

qRT-PCR was carried out in 96-well blocks with an Applied Biosystems 7500 Real-Time PCR System using the SYBR Green I master mix in a volume of 25 μ l. The PCR primers were designed using the software Primer Express; the primer sequences are listed in Supplementary Table 1. The 2-step thermal cycling profile used was 15 s at 94°C and 1 min at 68°C. An *elf4A* gene (At3g13920) was included in the assays as an internal control for normalizing the variations in cDNA quantities used (Gutierrez et al., 2008). The qRT-PCR reactions were carried out in biological triplicates using RNA samples extracted from 3 independent plant materials grown under identical growth conditions. The comparative $\Delta\Delta C_T$ method was used to evaluate the relative quantities of each amplified product in the samples. The threshold cycle (C_T) was automatically determined for each reaction by the System set with default parameters. The specificity of the PCR was determined by melt curve analysis of the amplified products using the standard method installed in the System.

Extraction of total RNA from seeds

Total RNA was extracted from dry seeds according to the procedure described previously (Suzuki et al., 2004). Seeds of 10–15 mg were routinely used, and total RNA was extracted using 700 μ l of extraction buffer (100 mM Tris-Cl, pH 9.5, 10 mM EDTA, 0.6 M NaCl, 0.4 M trisodium citrate, 2% (w/v) lithium dodecyl sulfate, and 5% (w/v) β -mercaptoethanol) after grinding in liquid nitrogen. The RNA samples were finally cleaned up using the Qiagen Plant Total RNA Isolation Kit (Qiagen, USA).

Histochemical staining

The primers used for subcloning of the *ASD* gene promoter were P_{ASD}:GUS-F (5'-AAAAAGCAGGCTTTCTTTATTGGGCCTTATTATAGCC) and P_{ASD}:GUS-R (5'-AGAAAGCTGGGTT

TTTTTGGATGCTCTGTTCCGG). The PCR product was subcloned into the pHGWS7 vector (Invitrogen, USA).

For histochemical analysis of β -glucuronidase (GUS) activity, plant materials were incubated in 90% acetone for 20 min on ice, washed twice with rinsing solution [50 mM sodium phosphate, pH 7.2, 0.5 mM $K_3Fe(CN)_6$, and 0.5 mM $K_4Fe(CN)_6$], and subsequently incubated at 37°C for 18–24 h in rinsing solution containing 2 mM 5-bromo-4-chloro-3-indolyl- β -D-glucuronide (X-Gluc) (Duchefa, The Netherlands). The plant materials were then incubated in a series of ethanol solutions ranging from 15% to 80% in order to remove chlorophylls from plant tissues. They were then mounted on microscope slides and visualized using a Nikon SMZ 800 microscope (Nikon, Japan).

Treatments with growth hormones and abiotic stresses

Two-week-old plants grown on 1/2 X Murashige and Skoog (MS)-agar plates (hereafter referred to as MS-agar plates) were transferred to fresh MS-agar plates supplemented with appropriate growth hormones. ABA and methyl jasmonic acid (MeJA) were used at a final concentration of 20 μ M. Salicylic acid (SA) was used at a final concentration of 100 μ M.

For the assays on the effects of drought on gene expression, 2-week-old plants grown on MS-agar plates were put on a dry 3 MM paper at room temperature for the indicated time periods. To examine the effects of high salinity on gene expression, 2-week-old plants grown on MS-agar plates were soaked in MS liquid cultures containing 200 mM NaCl and incubated under constant light for the indicated time periods. For cold treatments, 2-week-old plants grown on MS-agar plates were transferred to a cold chamber set at 4°C and incubated for the indicated time periods before harvesting plant materials. Whole plants were used for RNA extraction, unless otherwise specified.

Preparation of recombinant proteins

The wild-type *ASD* gene and a mutated *ASD* gene, in which K256 was mutated to either arginine or alanine, were fused in-frame to the 5' end of the maltose binding protein (MBP)-coding sequence in the pMBP-GW vector (Invitrogen, USA). The expression constructs were transformed into *Escherichia coli* strain BL21 cells. Cell cultures, induction, and protein purification were carried out according to the manufacturer's procedure (Novagen, Germany).

ATPase activity assay

The ATPase activity assay was carried out as previously described with minor modifications (Perlin and Spanswick, 1981). A reaction mixture containing 50 mM Tris-Cl, pH 7.5, 3 mM ATP, 3 mM $MgCl_2$, and 1 μ g of recombinant or MBP protein was incubated at 30°C for appropriate time periods. The reaction was terminated by adding a developing reagent (0.42% ammonium molybdate in 1 N H_2SO_4 :10% ascorbic acid at 5:1 [v/v] ratio). After incubation at 25°C for 30 min, the absorbance was measured at 820 nm. To determine the amount of released phosphate in the reaction mixture, potassium dihydrogen phosphate was used as the standard.

RESULTS

ASD is an AAA-type protein

The AAA proteins constitute a large superfamily (Snider et al., 2008). Among the AAA genes identified in *Arabidopsis*, the *ASD* gene is of particular interest. Gene expression analysis using the GENEVESTIGATOR database (<https://www.genevestigator.com/gv/index.jsp>) showed that the *ASD* gene is highly expressed in mature seeds and is influenced by environmental




Fig. 1. Protein structure of ASD and phylogenetic analysis of ASD homologues in plants. (A) Protein structure of ASD. The ASD protein contains a TM motif in the N-terminal region (black box) and an AAA domain in the C-terminal region (white box). The protein structure was analyzed using the software available in the ExPASy database (<http://us.expasy.org/tools/>). (B) Phylogenetic analysis of ASD and its homologues from several plant species. The accession numbers of the protein sequences analyzed are XP-002275941 (grape vine, *Vitis vinifera*), CAH10209 (wheat, *Triticum aestivum*), CAJ13559 (wheat, *Triticum turgidum*), AAV49988 (barley, *Hordeum vulgare*), ABA97668 (rice, *Oryza sativa*), and ABO99982 (potato, *Solanum tuberosum*). (C) Sequence alignment of the AAA domain from ASD and its homologues. Black boxes indicate identical residues, and gray boxes indicate biochemically conserved residues. The Walker A and B and SRH motifs are indicated above the sequences. The amino acid sequence alignment was carried out using the ClustalW server (<http://www.ebi.ac.uk/clustalw/>).

stress conditions.

Web-based bioinformatics tools were used to predict the protein structure of ASD. The analysis revealed that the ASD protein has a transmembrane motif in the N-terminal region and an AAA domain in the C-terminal region (Fig. 1A).

Protein homologues of ASD were identified in diverse plant species using the Basic Local Alignment Search Tool (BLAST) (<http://www.ncbi.nlm.nih.gov/blast/>). They included XP-002275941 from grape vine (*Vitis vinifera*), CAH10209 and CAJ13559 from wheat (*Triticum aestivum* and *Triticum turgidum*, respectively), AAV49988 from barley (*Hordeum vulgare*), ABA97668 from rice (*Oryza sativa*), and ABO99982 from potato (*Solanum tuberosum*), suggesting that the ASD protein is conserved throughout the plant kingdom (Fig. 1B). Multiple sequence alignments showed that ASD and its homologues share high levels of amino acid sequence identities, particularly within the AAA domains (Figs. 1B and 1C).




Fig. 2. Temporal and tissue-specific expression patterns of the *ASD* gene. In (A), (B), and (D), transcript levels were examined by using qRT-PCR. Biological triplicates were averaged. Bars represent standard error of the mean. (A) Temporal expression pattern. Total RNA was extracted from seeds or whole seedlings harvested at the indicated time points. d, days after germination. (B) Expression pattern in germinating seeds. Total RNA was extracted from the seeds harvested at the indicated time points. h, hours after cold imbibition. (C) Distribution of GUS activity in seeds. The *pASD-GUS* fusion construct, in which the GUS-coding sequence was transcriptionally fused to the *ASD* gene promoter sequence covering an approximately 1-kb region upstream of the transcription start site, was transformed into Col-0 plants. GUS activities were detected exclusively in the embryo. (D) Expression pattern during silique development. Total RNA was extracted from the siliques harvested at the indicated time points. Silique development covered from anthesis (0 DAP) to dry seed stage (21 DAP). d, days after pollination (DAP).

Typical AAA proteins contain 3 conserved protein motifs within the AAA domain: Walker A and B motifs and the SRH motif (Karata et al., 1999; Patel and Latterich, 1998). These protein motifs were conserved in *ASD* and its homologues, indicating that the *ASD* protein is an AAA-type protein (Fig. 1C).

The *ASD* gene is highly expressed in seeds

To obtain clues as to the role played by the *ASD* protein, we analyzed the temporal expression pattern of the *ASD* gene using qRT-PCR. The *ASD* gene was expressed predominantly in seeds (Fig. 2A). *ASD* gene expression was particularly high in dry seeds but rapidly decreased as seeds germinate, suggesting that the *ASD* gene plays a role in maintaining seed dormancy or regulating the seed maturation process (Fig. 2B).

We also examined the localized expression pattern of the *ASD* gene in seeds using a promoter-*GUS* gene fusion, in which the GUS-coding sequence was transcriptionally fused to the *ASD* gene promoter sequence covering an approximately 1-kb region upstream of the transcription start site. The fusion construct was transformed into Col-0 plants. The GUS activity was detected in seeds (Fig. 2C) but not in seedlings and fully grown plants (data not shown). Furthermore, *ASD* gene expression was observed only in the embryo but not in the seed coat (Fig. 2C).

The *ASD* gene was highly expressed in seeds, but its ex-




Fig. 3. Seed phenotypes of the 35S:*ASD* transgenic and *asd-1* mutant plants. (A) Mapping of the T-DNA insertion site in the *asd-1* mutant. (B) Transcript levels of the *ASD* gene. Transcript levels were examined by RT-PCR using RNA samples extracted from 2-week-old whole plants grown on MS-agar plates. (C) Seed phenotypes. Mature seeds were photographed. (D) Quantification of percentages of abnormal seeds. Approximately 200 seeds were used from each counting to calculate percentages of abnormal seeds. Bars represent standard error of the mean (*t*-test, **P* < 0.01). (E) Silique development. Images show representative siliques from plants grown in soil under long days for 50 days after germination. Scale bars = 1 mm. (F) Measurements of silique lengths. Silique lengths were calculated using 50 siliques for each plant group. Bars represent standard error of the mean (*t*-test, **P* < 0.01).

pression rapidly decreased in germinating seeds, suggesting that it is regulated by the seed maturation process. We therefore analyzed the expression kinetics of the *ASD* gene during silique development. *ASD* gene expression increased gradually during the seed developmental process and reached a peak at the late developmental stage [21 days after pollination (DAP)] (Fig. 2D), supporting that the *ASD* gene plays a role in seed development, particularly in seed maturation step.

The *ASD* protein is involved in seed maturation

To investigate the physiological role of the *ASD* gene in seed development and maturation, we produced transgenic plants (35S:*ASD*) that overexpress the *ASD* gene under the control of the CaMV 35S promoter. A T-DNA insertional knockout mutant *asd-1* was also obtained from the public database (Fig. 3A). Overexpression of the *ASD* gene in the 35S:*ASD* transgenic plants and lack of gene expression in the *asd-1* mutant were verified by RT-PCR before further analysis (Fig. 3B).

Although the phenotypes of the 35S:*ASD* transgenic plants were indistinguishable from wild-type plant, the transgenic plants produced a higher amount of abnormal seeds with disturbed morphology (Fig. 3C). Whereas approximately 5% of the seeds were morphologically abnormal in Col-0 plants, more than 20% of the seeds exhibited a distorted morphology in the 35S:*ASD* transgenic plants (Fig. 3D), further supporting the role of *ASD* in seed maturation. Seed maturation was not discerni-




Fig. 4. Delayed germination of the 35S:ASD transgenic seeds. Seeds were air-dried for 2 weeks after harvesting and imbibed at 4°C on MS-agar plates for 3 days. The cold-imbibed seeds were allowed to germinate at 22°C under long days. Appearance of visible radicles was used as a morphological marker for germination. Three measurements, each consisting of 80–100 seeds, were averaged. Bars represent standard error of the mean. h, hours after cold imbibition.

bly affected in the *asd-1* mutant, which may be due to functional redundancy among the AAA proteins.

Silique development was also influenced in the 35S:ASD transgenic plants. Length of the 35S:ASD transgenic siliques was shorter than that of Col-0 and *asd-1* mutant siliques (Fig. 3E). Whereas the silique length was reduced by approximately 23% in the 35S:ASD transgenic plants, it was slightly, but reproducibly, elongated in the *asd-1* mutant (Fig. 3F). Seed maturation is essential for seed viability and germination (Lee et al., 2010; Parcy et al., 1994). Consistent with the disturbed seed maturation in the 35S:ASD transgenic plants, germination of the transgenic seeds was significantly delayed (Fig. 4), indicating that the 35S:ASD transgenic seeds had defects in seed maturation processes. Overall, these observations indicated that the ASD gene is involved in the late stages of seed development, such as seed maturation and silique development.

Seed developmental marker genes are influenced in the transgenic and mutant seeds

To look into the molecular mechanisms underlying the function of the ASD gene in seed maturation, we isolated total RNAs from mature seeds and examined transcript levels of key transcription factor genes involved in seed development and maturation by using qRT-PCR. Expression of *LEC1*, *LEC2*, *FUS3*, and *ABI3* genes were reduced 2- to 5-fold in the 35S:ASD transgenic seeds but elevated 2- to 3-fold in the *asd-1* mutant seeds, consistent with the notion that the ASD gene plays a role in seed development and maturation (Fig. 5).

Seed developmental stages are characterized by distinct sets of marker genes (Parcy et al., 1994). The marker genes are grouped into 3 classes according to their temporal expression patterns: class I, maturation (MAT), and LEA. *Cold-regulated 47* (*COR47*), *kold-induced 1* (*KIN1*), *chalcone synthase* (*CHS*), *chalcone flavanone isomerase* (*CHI*), and *dihydroflavonol 4-reductase* (*DFR*) genes constitute the 'class I' gene family (Feinbaum and Ausubel, 1988; Gilmour et al., 1992; Kurkela and Franck, 1990; Parcy et al., 1994; Shirley et al., 1992). These genes were expressed predominantly during the earliest seed developmental stages, covering 0–10 DAP (Parcy et al., 1994). Whereas expression of the 'class I' genes was significantly elevated in the 35S:ASD transgenic seeds, it was downregulated in the *asd-1* mutant seeds (Fig. 5).

The marker genes of the maturation (MAT) phase include




Fig. 5. Transcript levels of seed developmental marker genes in the 35S:ASD transgenic and *asd-1* mutant seeds. Total RNA was extracted from dry seeds. Transcript levels were examined by using qRT-PCR. Biological triplicates were averaged. Bars represent standard error of the mean. The y-axis is presented on a logarithmic scale for better comparison of fold changes.

At2S3, which encodes the *Arabidopsis* napin 3 protein, *cruciferin C* (*CRC*), and *oleosin 1* (*OLEO1*) genes (Parcy et al., 1994). Their expression is initiated at 9 DAP and peaks at 15–18 DAP, functioning in the accumulation of storage proteins and lipids (Guerche et al., 1990; Pang et al., 1988; Parcy et al., 1994; van Rooijen et al., 1992). Expression patterns of the MAT members were distinct from those of the 'class I' marker genes in the 35S:ASD transgenic and *asd-1* mutant seeds. Whereas their expression was downregulated in the 35S:ASD transgenic seeds, it was upregulated in the *asd-1* mutant seeds (Fig. 5).

The LEA genes are expressed from 13 to 18 DAP, and their expression reaches a peak in the final days of silique development. The LEA gene group includes *Arabidopsis thaliana* late embryogenesis abundant 1 (*AtEm1*), *AtEm6*, and *responsive to ABA 18* (*RAB18*) genes (Finkelstein, 1993; Gaubier et al., 1993; Lång and Palva, 1992; Parcy et al., 1994). Expression patterns of the LEA gene group members were similar to those of the MAT genes in the 35S:ASD transgenic and *asd-1* mutant seeds (Fig. 5). Together with the seed phenotypes of the 35S:ASD transgenic plants, these results indicate that the ASD gene plays a role in the seed maturation stage and that seed development is arrested prior to completion of seed maturation




Fig. 6. Mitochondrial localization of the ASD protein in *Arabidopsis* protoplasts. An *ASD-RFP* gene fusion, in which the RFP-coding sequence was fused in-frame to the 3' end of the *ASD* gene, was transiently expressed in *Arabidopsis* protoplasts and visualized by fluorescence microscopy. The mt-yk CD3-989 construct was coexpressed as a mitochondrial marker (Nelson et al., 2007).

in the 35S:*ASD* transgenic plants.

The ASD protein is localized in mitochondria

The AAA proteins are involved in diverse cellular activities (Lupas and Martin, 2002; Patel and Latterich, 1998). Accordingly, their subcellular localizations vary. To determine the subcellular localization of the ASD protein, a red fluorescence protein (RFP)-coding sequence was fused in frame to the 3' end of the *ASD* gene, and the fusion construct was expressed transiently in *Arabidopsis* protoplasts. RFP signals were broadly distributed within the cell, in a manner similar to the distribution of mitochondrial proteins (Nelson et al., 2007).

To examine whether the ASD protein is localized into the mitochondria, organelle-specific marker constructs were coexpressed with the *ASD-RFP* gene fusion. RFP signals were clearly overlaid with the subcellular distribution of the mitochondrial marker CD3-989, which is fused to yellow fluorescence protein (YFP) (Fig. 6), indicating that the ASD protein is a mitochondrial AAA protein.

The ASD protein has ATPase activity

ASD is a mitochondrial AAA protein having conserved protein motifs required for ATPase activity. To examine whether the ASD protein has ATPase activity, a recombinant ASD protein was purified as an MBP fusion from *E. coli* cells. We also generated mutated ASD proteins (mASDs), in which the absolutely conserved lysine residue within the Walker A domain, K256, was mutated to either alanine or arginine, resulting in K256A or K256R. Immunoblot analysis using an anti-MBP antibody revealed that the purified recombinant proteins had an estimated molecular mass of 102 kDa (ASD 60 kDa + MBP 42 kDa = 102 kDa) (Fig. 7A).

The purified recombinant ASD proteins were subject to ATPase activity assays. The MBP-ASD recombinant protein showed remarkable ATPase activity (Fig. 7B). In contrast, the mutated K256A or K256R proteins exhibited significantly suppressed ATPase activities, demonstrating that the ASD protein possesses ATPase activity and that K256 within the Walker A motif is critical for ATP hydrolysis. In addition, ATP hydrolytic activity was greatly enhanced in the presence of Mg^{2+} , showing that the ASD protein is an Mg^{2+} -dependent ATPase (Fig. 7B). MBP protein itself did not exhibit any detectable activity in ATP hydrolysis.

The ASD gene is induced by abiotic stresses in an ABA-dependent manner

To determine the nature of environmental signals regulating the *ASD* gene, we examined the effects of various environmental




Fig. 7. ATPase activity assays of recombinant ASD and mASD proteins. (A) Purified ASD proteins. Recombinant MBP-ASD and -mASD fusion proteins were prepared in *Escherichia coli* cells and partially purified. The recombinant proteins were detected by Coomassie staining (left panel) and Western blot analysis using an anti-MBP antibody (right panel). kDa, kilodaltons. (B) ATPase activity assays. ATP hydrolysis activity of recombinant proteins was estimated in the presence (+) or absence (-) of 3 mM $MgCl_2$. MBP protein was also included in the assays. Three measurements were averaged (*t*-test, **P* < 0.01). Bars represent standard error of the mean.

stress conditions and growth hormones on *ASD* gene expression. *ASD* transcription was upregulated significantly by drought, cold, and salt stresses (Fig. 8A). The *ASD* gene was also induced significantly by ABA but unaffected by SA and MeJA (Fig. 8B), suggesting that the *ASD* gene is involved in abiotic stress responses.

We next examined whether induction of the *ASD* gene by abiotic stresses depends on ABA. Salt induction of the *ASD* gene completely disappeared in the ABA-deficient *aba3-1* mutant (Fig. 8C), indicating that the *ASD* gene is regulated by high salinity in an ABA-dependent manner.

DISCUSSION

Some members of the AAA-type ATPase family possess several biochemical activities, such as metalloprotease activities, in addition to intrinsic ATPase activity and are functional constituents of 26S proteasome complexes. Consistent with the diverse biochemical activities exerted by the AAA-type ATPases, they have been implicated in a variety of cellular and physiological processes, including biogenesis of peroxisomes and mitochondria and HR responses (Olsen, 1998; Sugimoto et al., 2004).

Here, we identified an *Arabidopsis* AAA-type ATPase gene, *ASD*, which is involved in the seed maturation process, thereby




Fig. 8. Effects of abiotic stresses and growth hormones on *ASD* gene expression. Two-week-old plants grown on MS-agar plates were used for treatments with abiotic stresses and growth hormones. Transcript levels were determined by qRT-PCR. Biological triplicates were averaged. Bars indicate standard error of the mean. (A) Effects of abiotic stresses. Plants were exposed to drought, 4°C, or 150 mM NaCl for the indicated time periods before harvesting plant materials for total RNA extraction. (B) Effects of growth hormones. Plants were treated with 20 μ M ABA, 100 μ M SA, or 20 μ M MeJA for the indicated time periods before harvesting plant materials. (C) Effects of high salinity on *ASD* gene expression in the *aba3-1* mutant.

further extending the repertoire of the roles played by AAA-type ATPases in *Arabidopsis*. *ASD* gene expression is regulated by both developmental and environmental cues. Overexpression of the *ASD* gene leads to disrupted seed development and maturation.

The *ASD* gene is expressed predominantly in developing seeds. It is notable that the *ASD* gene is also induced under abiotic stress conditions, which certainly affect seed maturation processes. Seed development is known to be influenced by environmental factors. In cowpea (*Vigna unguiculata*), drought stress reduces seed yields because of defects in seed maturation (Summerfield et al., 1976). Stress regulation of seed development has also been observed in other plant species (Cooper et al., 2003; Martin et al., 2010). Reproduction or seed production is an energy-consuming process. Acquisition of stress

resistance also requires a high input of metabolic energy, implying that proper deposition of energy and metabolites is necessary to complete reproduction under stress conditions. The mitochondrial *ASD* ATPase may play a role in mediating developmental and environmental signals to maintain proper seed maturation process, contributing to plant fitness to energy and nutrient distributions.

The AAA-type ATPases regulate diverse cellular functions with versatile biochemical activities. One subfamily of the AAA-type ATPases possesses metalloprotease activity, as exemplified by FtsH (Karata et al., 1999; Langer, 2000). Membrane-bound organelles serve as membrane-integrated quality-control systems, which are involved in protein folding and selective degradation of non-native polypeptides (Langer, 2000). Mitochondria are representative subcellular organelles that harbor ATP-dependent proteases and chaperone proteins (Langer et al., 2001). The AAA-type proteases form large protein complexes consisting of identical or closely related subunits in association with the membranes. Such protease complexes are thought to be involved in the degradation of mitochondrial membrane proteins (Leonhard et al., 2000). Most of the mitochondrial AAA-type ATPases contain transmembrane (TM) motifs. Whereas the *i*-AAA-type members have 1 TM motif, the *m*-AAA-type members have 2 TM motifs (Urantowka et al., 2005). The N-terminal TM motif is required for oligomerization of the *m*-AAA-type members (Urantowka et al., 2005).

The *ASD* ATPase contains the conserved AAA domain. It also has 1 TM motif in the N-terminal region. The presence of these structural components and the localization of *ASD* in the mitochondria suggest that a certain protease activity resides in the *ASD* protein, although this activity was not determined in this study. According to this view, it is possible that the *ASD* protein may play a role in membrane-mediated protein degradation in mitochondria and may contribute to the proper regulation of seed development and maturation.

Mitochondria play a central role in ATP production via the tricarboxylic acid (TCA) cycle and the integration of carbon and nitrogen metabolism (Pellny et al., 2008; van Aken et al., 2009). Controlled distribution and availability of metabolic energy and nutrients is important for cellular and organismal adaptation under environmental stress conditions. In this regard, mitochondria are the target of environmental stresses. Studies have shown that mitochondria respond to diverse environmental stresses (Grelet et al., 2004; van Aken et al., 2009) and that mitochondria may regulate cellular stress responses (Clifton et al., 2006; van Aken et al., 2009). The prevention of excessive reactive oxygen species (ROS) accumulation and the regulation of energy metabolism are well-known examples. In addition, some mitochondrial proteins, such as alternative oxidases, nicotinamide adenine dinucleotide [NAD(P)H] dehydrogenases, and heat shock proteins, are stress-responsive (van Aken et al., 2009). Although the protein targets of the *ASD* ATPase are still elusive, *ASD*-mediated signaling may be associated with mitochondrial stress responses during seed maturation.

Note: Supplementary information is available on the Molecules and Cells website (www.molcells.org).

ACKNOWLEDGMENTS

This work was supported by the Brain Korea 21 and Biogreen 21 (20080401034001) Programs and by grants from the Plant Signaling Network Research Center, the National Research Foundation of Korea (2009-0087317 and 2007-03415), and from the Agricultural R&D Promotion Center (309017-5), Korea Ministry for Food, Agriculture, Forestry and Fisheries.

REFERENCES

- Clifton, R., Millar, A.H., and Whelan, J. (2006). Alternative oxidases in *Arabidopsis*: a comparative analysis of differential expression in the gene family provides new insights into function of non-phosphorylating bypasses. *Biochim. Biophys. Acta* 1757, 730-741.
- Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16, 735-743.
- Cooper, B., Clarke, J.D., Budworth, P., Kreps, J., Hutchison, D., Park, S., Guimil, S., Dunn, M., Luginbühl, P., Ellero, C., et al. (2003). A network of rice genes associated with stress response and seed development. *Proc. Natl. Acad. Sci. USA* 100, 4945-4950.
- Crouch, M.L. (1987). Regulation of gene expression during seed development in flowering plants. In *Developmental Biology: A Comprehensive Synthesis*, Vol. 5, L.W. Browder, ed. (New York: Plenum Press), pp. 367-404.
- Feinbaum, R.L., and Ausubel, F.M. (1988). Transcriptional regulation of the *Arabidopsis thaliana* chalcone synthase gene. *Mol. Cell. Biol.* 8, 1985-1992.
- Finch-Savage, W.E., and Leubner-Metzger, G. (2006). Seed dormancy and the control of germination. *New Phytol.* 171, 501-523.
- Finkelstein, R.R. (1993). Abscissic acid-insensitive mutations provide evidence for stage-specific signal pathways regulating expression of an *Arabidopsis* late embryogenesis-abundant (*lea*) gene. *Mol. Gen. Genet.* 238, 401-408.
- Finkelstein, R.R., and Somerville, C.R. (1990). Three classes of abscissic acid (ABA)-insensitive mutations of *Arabidopsis* define genes that control overlapping subsets of ABA responses. *Plant Physiol.* 94, 1172-1179.
- Fu, H., Doelling, J.H., Rubin, D.M., and Vierstra, R.D. (1999). Structural and functional analysis of the six regulatory particle triple-A ATPase subunits from the *Arabidopsis* 26S proteasome. *Plant J.* 18, 529-539.
- Gaubier, P., Raynal, M., Hull, G., Huestis, G.M., Grellet, F., Arenas, C., Pagès, M., and Delseny, M. (1993). Two different *Em-like* genes are expressed in *Arabidopsis thaliana* seeds during maturation. *Mol. Gen. Genet.* 238, 409-418.
- Gilmour, S.J., Artus, N.N., and Thomashow, M.F. (1992). cDNA sequence analysis and expression of two cold-regulated genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* 18, 13-21.
- Goldberg, R.B., Barker, S.J., and Perez-Grau, L. (1989). Regulation of gene expression during plant embryogenesis. *Cell* 56, 149-160.
- Grellet, J., Benamar, A., Teyssier, E., Avelange-Macherel, M.H., Grunwald, D., and Macherel, D. (2004). Identification in pea seed mitochondria of a late-embryogenesis abundant protein able to protect enzymes from drying. *Plant Physiol.* 137, 157-167.
- Guerche, P., Tire, C., De Sa, F.G., De Clercq, A., Van Montagu, M., and Krebbers, E. (1990). Differential expression of the *Arabidopsis* 2S albumin genes and the effect of increasing gene family size. *Plant Cell* 2, 469-478.
- Gutierrez, L., Van Wuytswinkel, O., Castelain, M., and Bellini, C. (2007). Combined networks regulating seed maturation. *Trends Plant Sci.* 12, 294-300.
- Gutierrez, L., Mauriat, M., Guénin, S., Pelloux, J., Lefebvre, J.F., Louvet, R., Rusterucci, C., Moritz, T., Guérineau, F., and Bellini, C. (2008). The lack of a systematic validation of reference genes: a serious pitfall undervalued in reverse transcription-polymerase chain reaction (RT-PCR) analysis in plants. *Plant Biotechnol. J.* 6, 609-618.
- Hanson, P.I., and Whiteheart, S.W. (2005). AAA+ proteins: have engine, will work. *Nat. Rev. Mol. Cell Biol.* 6, 519-529.
- Harada, J.J., Dietrich, R.A., Comal, L., and Baden, C.S. (1988). Regulation of gene expression during seed germination and post germinative development. In *Plant Gene Research: Temporal and Spatial Regulation of Plant Genes*, Vol. 5, D.P.S. Verma, and R.B. Goldberg, eds. (Wien: Springer-Verlag), pp. 27-39.
- Holdsworth, M.J., Bentsink, L., and Soppe, W.J. (2008). Molecular networks regulating *Arabidopsis* seed maturation, after-ripening, dormancy and germination. *New Phytol.* 179, 33-54.
- Huang, N., Reinl, S.J., and Rodriguez, R.L. (1992). *RAMY2A*; a novel alpha-amylase-encoding gene in rice. *Gene* 111, 223-228.
- Kagaya, Y., Toyoshima, R., Okuda, R., Usui, H., Yamamoto, A., and Hattori, T. (2005). *LEAFY COTYLEDON1* controls seed storage protein genes through its regulation of *FUSCA3* and *ABSCISIC ACID INSENSITIVE3*. *Plant Cell Physiol.* 46, 399-406.
- Karata, K., Inagawa, T., Wilkinson, A.J., Tatsuta, T., and Ogura, T. (1999). Dissecting the role of a conserved motif (the second region of homology) in the AAA family of ATPases. Site-directed mutagenesis of the ATP-dependent protease FtsH. *J. Biol. Chem.* 274, 26225-26232.
- Keith, K., Kraml, M., Dengler, N.G., and McCourt, P. (1994). *fusca3*: a heterochronic mutation affecting late embryo development in *Arabidopsis*. *Plant Cell* 6, 589-600.
- Kermode, A.R. (1990). Regulatory mechanisms involved in the transition from seed development to germination. *Crit. Rev. Plant Sci.* 81, 280-288.
- Kim, Y.S., Kim, S.G., Park, J.E., Park, H.Y., Lim, M.H., Chua, N.H., and Park, C.M. (2006). A membrane-bound NAC transcription factor regulates cell division in *Arabidopsis*. *Plant Cell* 18, 3132-3144.
- Kurkela, S., and Franck, M. (1990). Cloning and characterization of a cold- and ABA-inducible *Arabidopsis* gene. *Plant Mol. Biol.* 15, 137-144.
- Lång, V., and Palva, E.T. (1992). The expression of a *rab-related* gene, *rab18*, is induced by abscisic acid during the cold acclimation process of *Arabidopsis thaliana* (L.) Heynh. *Plant Mol. Biol.* 20, 951-962.
- Langer, T. (2000). AAA proteases: cellular machines for degrading membrane proteins. *Trends Biochem. Sci.* 25, 247-251.
- Langer, T., Käser, M., Klanner, C., and Leonhard, K. (2001). AAA proteases of mitochondria: quality control of membrane proteins and regulatory functions during mitochondrial biogenesis. *Biochem. Soc. Trans.* 29, 431-436.
- Lee, Y.J., Kim, D.H., Kim, Y.-W., and Hwang, I. (2001). Identification of a signal that distinguishes between the chloroplast outer envelope membrane and the endomembrane system *in vivo*. *Plant Cell* 13, 2175-2190.
- Lee, S.J., Cho, D.I., Kang, J.Y., Kim, M.D., and Kim, S.Y. (2010). *AtNEK6* interacts with *ARIA* and is involved in ABA response during seed germination. *Mol. Cells* 29, 559-566.
- Leonhard, K., Guiard, B., Pellicchia, G., Tzagoloff, A., Neupert, W., and Langer, T. (2000). Membrane protein degradation by AAA proteases in mitochondria: extraction of substrates from either membrane surface. *Mol. Cell* 5, 629-638.
- Lindahl, M., Tabak, S., Cseke, L., Pichersky, E., Andersson, B., and Adam, Z. (1996). Identification, characterization, and molecular cloning of a homologue of the bacterial FtsH protease in chloroplasts of higher plants. *J. Biol. Chem.* 271, 29329-29334.
- Luerssen, H., Kirik, V., Herrmann, P., and Miséra, S. (1998). *FUSCA3* encodes a protein with a conserved VP1/AB13-like B3 domain which is of functional importance for the regulation of seed maturation in *Arabidopsis thaliana*. *Plant J.* 15, 755-764.
- Lupas, A.N., and Martin, J. (2002). AAA proteins. *Curr. Opin. Struct. Biol.* 12, 746-753.
- Martin, R.C., Liu, P.P., Goloviznina, N.A., and Nonogaki, H. (2010). microRNA, seeds, and Darwin?: diverse function of miRNA in seed biology and plant responses to stress. *J. Exp. Bot.* 61, 2229-2234.
- McCarty, D.R., and Carson, C.B. (1991). The molecular genetics of seed maturation in maize. *Physiol. Plant* 81, 267-272.
- Meinke, D.W., Franzmann, L.H., Nickle, T.C., and Yeung, E.C. (1994). Leafy cotyledon mutants of *Arabidopsis*. *Plant Cell* 6, 1049-1064.
- Nambara, E., and Marion-Poll, A. (2003). ABA action and interactions in seeds. *Trends Plant Sci.* 8, 213-217.
- Natesh, S., and Rau, M.A. (1984). The embryo. In *Embryology of Angiosperms*, B.M., Johri, ed. (Berlin: Springer-Verlag), pp. 377-443.
- Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of *in vivo* organelle markers for co-localization studies in *Arabidopsis* and other plants. *Plant J.* 51, 1126-1136.
- Olsen, L.J. (1998). The surprising complexity of peroxisome biogenesis. *Plant Mol. Biol.* 38, 163-189.
- Pang, P.P., Pruitt, R.E., and Meyerowitz, E.M. (1988). Molecular cloning, genomic organization, expression and evolution of 12S seed storage protein genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* 11, 805-820.
- Parcy, F., Valon, C., Raynal, M., Gaubier-Comella, P., Delseny, M.,

- and Giraudat, J. (1994). Regulation of gene expression programs during *Arabidopsis* seed development: roles of the *ABI3* locus and of endogenous abscisic acid. *Plant Cell* 6, 1567-1582.
- Parcy, F., Valon, C., Kohara, A., Miséra, S., and Giraudat, J. (1997). The *ABSCISIC ACID-INSENSITIVE3*, *FUSCA3*, and *LEAFY COTYLEDON1* loci act in concert to control multiple aspects of *Arabidopsis* seed development. *Plant Cell* 9, 1265-1277.
- Patel, S., and Latterich, M. (1998). The AAA team: related ATPases with diverse functions. *Trends Cell Biol.* 8, 65-71.
- Pellny, T.K., Van Aken, O., Dutilleul, C., Wolff, T., Groten, K., Bor, M., De Paepe, R., Reyss, A., Van Breusegem, F., Noctor, G., et al. (2008). Mitochondrial respiratory pathways modulate nitrate sensing and nitrogen-dependent regulation of plant architecture in *Nicotiana glauca*. *Plant J.* 54, 976-992.
- Perlin, D.S., and Spanswick, R.M. (1981). Characterization of ATPase activity associated with corn leaf plasma membranes. *Plant Physiol.* 68, 521-526.
- Rancour, D.M., Dickey, C.E., Park, S., and Bednarek, S.Y. (2002). Characterization of AtCDC48. Evidence for multiple membrane fusion mechanisms at the plane of cell division in plants. *Plant Physiol.* 130, 1241-1253.
- Raz, V., Bergervoet, J.H., and Koornneef, M. (2001). Sequential steps for developmental arrest in *Arabidopsis* seeds. *Development* 128, 243-252.
- Shirley, B.W., Hanley, S., and Goodman, H.M. (1992). Effects of ionizing radiation on a plant genome: analysis of two *Arabidopsis transparent testa* mutations. *Plant Cell* 4, 333-347.
- Snider, J., Thibault, G., and Houry, W.A. (2008). The AAA+ superfamily of functionally diverse proteins. *Genome Biol.* 9, 216.
- Stone, S.L., Kwong, L.W., Yee, K.M., Pelletier, J., Lepiniec, L., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (2001). *LEAFY COTYLEDON2* encodes a B3 domain transcription factor that induces embryo development. *Proc. Natl. Acad. Sci. USA* 98, 11806-11811.
- Sugimoto, M., Yamaguchi, Y., Nakamura, K., Tatsumi, Y., and Sano, H. (2004). A hypersensitive response-induced ATPase associated with various cellular activities (AAA) protein from tobacco plants. *Plant Mol. Biol.* 56, 973-985.
- Summerfield, R.J., Huxley, P.A., Dart, P.J., and Hughes, A.P. (1976). Some effects of environmental stress on seed yield of cowpea (*Vigna unguiculata* (L.) Walp.) cv. Prima. *Plant Soil* 44, 527-546.
- Suzuki, Y., Kawazu, T., and Koyama, H. (2004). RNA isolation from siliques, dry seeds, and other tissues of *Arabidopsis thaliana*. *Biotechniques* 37, 542-544.
- Udvardi, M.K., Czechowski, T., and Scheible, W.R. (2008). Eleven golden rules of quantitative RT-PCR. *Plant Cell* 20, 1736-1737.
- Urantowska, A., Knorpp, C., Olczak, T., Kolodziejczak, M., and Janska, H. (2005). Plant mitochondria contain at least two i-AAA-like complexes. *Plant Mol. Biol.* 59, 239-252.
- van Aken, O., Zhang, B., Carrie, C., Uggalla, V., Paynter, E., Giraud, E., and Whelan, J. (2009). Defining the mitochondrial stress response in *Arabidopsis thaliana*. *Mol. Plant* 2, 1310-1324.
- van Rooijen, G.J., Teming, L.I., and Moloney, M.M. (1992). Nucleotide sequence of an *Arabidopsis thaliana* *oleosin* gene. *Plant Mol. Biol.* 18, 1177-1179.
- Verdier, J., and Thompson, R.D. (2008). Transcriptional regulation of storage protein synthesis during dicotyledon seed filling. *Plant Cell Physiol.* 49, 1263-1271.
- West, M., and Harada, J.J. (1993). Embryogenesis in higher plants: An overview. *Plant Cell* 5, 1361-1369.
- West, M., Yee, K.M., Danao, J., Zimmerman, J.L., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (1994). *LEAFY COTYLEDON1* is an essential regulator of late embryogenesis and cotyledon identity in *Arabidopsis*. *Plant Cell* 6, 1731-1745.
- Wobus, U., and Weber, H. (1999). Seed maturation: genetic programmes and control signals. *Curr. Opin. Plant Biol.* 2, 33-38.
- Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat. Protoc.* 2, 1565-1572.