

## Reaction of 1,3-Dicarbonyl Compounds with Nitril Chloride

### Short Communication

Mladen V. Proštenik\*, Miljenko Dumić, and Nada Bošnjak

PLIVA, Pharmaceutical and Chemical Works, I.L. Ribara 89,  
41000 Zagreb, Croatia, Yugoslavia

(Received 14 February 1980. Accepted 29 February 1980)

Reaction of 1,3-dicarbonyl compounds with nitril chloride through substitution on the activated methylene group, resulted in the formation of chloro, dichloro and oximino derivatives.

(Keywords: 3-Oxobutanoic acid ethyl ester, 2,4-Pentanedione, reaction with nitril chloride)

#### *Die Reaktion der 1,3-Dicarbonylverbindungen mit Nitrilchlorid*

Umsetzung von 1,3-Dicarbonylverbindungen mit Nitrilchlorid ergab die entsprechende Chloro-, Dichloro- und Oximinoderivate in einer Substitutionsreaktion an der aktivierten Methylengruppe.

Nitril chloride reacts with aliphatic<sup>1</sup>, alicyclic<sup>2</sup> and heterocyclic olefins<sup>3</sup> to give mixtures of addition products, such as vicinal dichloro and chloro-nitro compounds, whereas aromatic<sup>4</sup> and heteroaromatic<sup>5</sup> hydrocarbons are substituted and chloro or nitro compounds or their mixtures are formed. Substitution reactions with nitril chloride in the aliphatic series have not been known so far.

We now wish to report that reaction of 1,3-dicarbonyl compounds with nitril chloride gives rise to substitution on the activated methylene group. Various examples with reaction conditions and yields of the products are shown in Table 1. The variation in yields depends on the solvent as well as on the molar ratio of nitril chloride/1,3-dicarbonyl compound, while they are almost temperature independent.

The reaction is carried out by passing nitril chloride through the solution of the dicarbonyl compound in inert solvents. The products were isolated by fractional distillation, and their structure confirmed by spectral data and compared with authentic samples prepared by known procedures.


Table 1. Reaction of 1,3-dicarbonyl compounds **1 a** and **c** with nitryl chloride

	NO <sub>2</sub> Cl/ <b>1 a, c</b> molar ratio	Solvent	Reaction temp. (°C)	Products <sup>a</sup>		
				chloro (%)	dichloro (%)	oxime (%)
<b>1 a</b>	1	CH <sub>2</sub> Cl <sub>2</sub>	40		30 <sup>b</sup>	57 <sup>c</sup>
<b>1 a</b>	2	CH <sub>2</sub> Cl <sub>2</sub>	40		59	15
<b>1 c</b>	1	CH <sub>2</sub> Cl <sub>2</sub>	20	30 <sup>d</sup>	11 <sup>e</sup>	57 <sup>f</sup>
<b>1 c</b>	2	CH <sub>2</sub> Cl <sub>2</sub>	20	36	58	
<b>1 e</b>	1	C <sub>6</sub> H <sub>6</sub>	20	24	12	63
<b>1 c</b>	2	CH <sub>2</sub> Cl <sub>2</sub>	40	60	40	
<b>1 c</b>	2	acetic acid	40		80	19

<sup>a</sup> Yields determined by G.c.; <sup>b</sup> b.p.<sub>18</sub> 85-87 °C (lit.<sup>9</sup> b.p.<sub>18-20</sub> 87 °C); <sup>c</sup> m.p. 72 °C (lit.<sup>10</sup> 75 °C); <sup>d</sup> b.p.<sub>30</sub> 93-98 °C (lit.<sup>11</sup> b.p.<sub>30</sub> 94-96 °C); <sup>e</sup> b.p.<sub>30</sub> 108-109 °C (lit.<sup>11</sup> b.p.<sub>30</sub> 106-108 °C); <sup>f</sup> m.p. 56-58 °C (lit.<sup>12</sup> m.p. 58-58.5 °C).

Special attention was paid to the detection of products having a nitro group but the results were unsatisfactory: High yields of the oxime **2 e** (obtained at a 1:1 molar ratio of nitryl chloride and ethyl acetoacetate, **1 e**) indicate the presence of an ONO<sup>-</sup> species. In some experiments, anhydrous calcium chloride or sodium sulphate was added, thus slightly increasing the yield of the oxime **2 e** from 63% to 69% (in benzene). At higher molar ratios of NO<sub>2</sub>Cl/diketone, chlorination predominated.

The results confirmed earlier observations that chlorination with nitryl chloride through substitution releases nitrite ions<sup>5,6</sup>. The ionic structures of nitryl chloride (cf. *Paul et al.*)<sup>7</sup> do not prefer those with positive chlorine and nitrite ion, but the results presented as well as those reported earlier<sup>5,6,8</sup> indicate that such a structure may exist.



	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
a	CH <sub>3</sub>	Cl	Cl
b	CH <sub>3</sub>	=NOH	
c	OC <sub>2</sub> H <sub>5</sub>	Cl	H
d	OC <sub>2</sub> H <sub>5</sub>	Cl	Cl
e	OC <sub>2</sub> H <sub>5</sub>	=NOH	

## Experimental

Gas-liquid chromatographic analyses were performed on a Perkin Elmer F-11 Model using column  $2 \times 0.004$  m, 2% diethyleneglycol succinate on 80-100 mesh Chromosorb G (AW-DMCS), isothermal at 80 °C (23 min) and programed (10°/min) to 180 °C. Ir spectra were taken on a Perkin Elmer 257 instrument. 3,3-Dichloro-2,4-pentanedione<sup>9</sup>, 3-oximino-2,4-pentanedione<sup>10</sup>, ethyl 2-chloroacetoacetate<sup>11</sup>, ethyl 2,2-dichloroacetoacetate<sup>11</sup>, and ethyl 2-oximinoacetoacetate<sup>12</sup> used as test substances were prepared according to known procedures.

### *General Reaction of 1,3-diketones with Nitryl Chloride*

Nitryl chloride (prepared according to *Schechter et al.*<sup>13</sup>), liquified in an acetone-dry ice bath, was allowed to evaporate into a stirred solution of 0.25 mol of 1,3-diketone **1 a** or **1 c** in 100 ml of solvent during one hour and stirring was then continued for an additional hour. Samples for gas chromatography were taken at this stage. The solvent was evaporated in vacuo on a Büchi Rotavapor. On fractionation the liquid chloro derivatives **1 a, c, d** were obtained, whereas the crystalline oximes **2 b** and **e** were isolated from the residue.

The infrared spectra of the products isolated were identical with those of the authentic samples.

## References

- <sup>1</sup> *J. Ville* and *G. Dupont*, Bull. Soc. Chim. France **1956**, 804; *S. Bresadola*, *P. Canal*, *A. Nenz*, and *E. Gallinella*, Chim. Ind. (Milan) **45**, 937 (1963).
- <sup>2</sup> *C. C. Price* and *C. A. Sears*, J. Amer. Chem. Soc. **75**, 3275 (1953).
- <sup>3</sup> *M. Dumić*, *M. V. Proštenik*, and *I. Butula*, Croat. Chem. Acta **51**, 259 (1978).
- <sup>4</sup> *M. J. Collis* and *D. R. Doddard*, J. Chem. Soc. **1958**, 1952.
- <sup>5</sup> *M. V. Proštenik*, *V. Slukan*, *F. Zorko*, *Z. Hranilović*, and *S. Škreb*, Croat. Chem. Acta **48**, 179 (1976).
- <sup>6</sup> *F. P. Gintz*, *D. R. Doddard*, and *M. J. Collis*, J. Chem. Soc. **1958**, 445.
- <sup>7</sup> *R. C. Paul*, *D. Singh*, and *K. C. Malhotra*, J. Chem. Soc (A) **1969**, 1396.
- <sup>8</sup> *J. Beger*, J. Pract. Chem. **35**, 326 (1967).
- <sup>9</sup> *A. Combes*, Compt. rend. **111**, 273 (1890).
- <sup>10</sup> *V. Wolff*, *P. Block*, *G. Lorentz*, and *P. Trappe*, Ann. Chem. **325**, 134 (1902).
- <sup>11</sup> *D. P. Wyman*, *P. R. Kaufman*, and *W. R. Freeman*, J. Org. Chem. **29**, 2706 (1964).
- <sup>12</sup> *H. Adkins* and *E. W. Reeve*, J. Amer. Chem. Soc. **60**, 1328 (1938).
- <sup>13</sup> *H. Shechter*, *F. Conrad*, *A. L. Daullton*, and *R. B. Kaplan*, J. Amer. Chem. Soc. **74**, 3052 (1952).