

Synthesis of C_1 - and C_{3v} -Symmetric Porphyrin Trimers Based on Triphenylmethane Cores

Cyril A. Papamicaël^{1,*}, Olivier Mongin², and Albert Gossauer³

¹ UMR 6014 IRCOF, CNRS, Université et INSA de Rouen, Mont-Saint-Aignan Cédex, France

² Synthèse et ElectroSynthèse Organiques (CNRS, UMR 6510), Université de Rennes 1, Institut de Chimie, Campus Scientifique de Beaulieu, Rennes Cédex, France

³ Institut für Organische Chemie der Universität, Fribourg, Switzerland

Received January 30, 2007; accepted (revised) February 24, 2007; published online May 23, 2007

© Springer-Verlag 2007

Summary. This work describes the synthesis of a new class of tripodaphyrin derivatives with a triphenylmethane core. Both C_1 - and C_{3v} -symmetric tetrahedral large molecules with covalently linked rigid elements were obtained.

Keywords. Porphyrines; Alkynes; Nanostructures; Chirality; Tripodaphyrin.

Introduction


The design and construction of novel porphyrin architectures with well-defined geometries is an area of increasing current interest [1]. Multi-porphyrins macrorings have attracted enormous interest because porphyrins provide rigid frameworks with relatively easy synthesis approaches [2]. Their biological importance, as it was found that the antenna chlorophylls in photosynthetic bacteria are arranged into macrorings to absorb and transfer solar energy with great efficiency, makes them one of the attracting research targets. In preceding papers [3, 4] we described the synthesis of tripodaphyrins, a new class of porphyrine derivatives. Tripodaphyrins are pyramidal assemblies in which a porphyrin macrocycle, situated on the top of the molecule, is “supported” by three “legs” consisting of linear arrays of covalently

linked rigid constitutive elements. Our principal aims of applications of tripodaphyrins are both studies of these molecules as light-harvesting systems arrays and molecular-based information storage. Indeed, one can envisage that multiple bits of information could be stored through the positioning of (\pm)-chiral tripodaphyrins by means of scanning probe microscopy (STM/AFM). In the present work, we describe the full synthetic details of C_1 - and C_{3v} -symmetric tripodaphyrin derivatives.

Results and Discussion

As a starting point for the design of a C_{3v} -symmetric tripodaphyrin derivative, we choose readily available 4,4',4''-methylidynetrisphenol [5] skeleton **1**. Its reaction with a threefold excess of trifluoromethanesulfonyl chloride afforded the expected compound **2** in 80% yield. Substitution of all three triflate groups by ethynyl groups was achieved in 95% yield on reaction of **2** with triisopropylsilylacetylene under Pd(0)-catalyzed conditions using copper iodide (Scheme 1). Thereafter, cleavage of both trialkylsilyl protecting groups gave the core molecule **4** (70%), which was reacted with [5-(4-iodophenyl)-10,15,20-triphenylporphinato(2-)]zinc [4] to afford the desired C_{3v} -symmetric porphyrin trimer **6** in 49% yield.

* Corresponding author. E-mail: cyril.papamicael@insa-rouen.fr




Scheme 1

The strategy devised for the synthesis of a (\pm)-chiral tripodaphyrin derivative (C_1 -symmetry) was based upon the use of pure [6] tris(4-iodophenyl)methane (**8**), which was prepared in 89% yield starting from **7**. Then, successive replacement of the iodine atoms of triphenylmethane **8** by *Sonogashira* coupling reactions with ethynyltrimethylsilane and compound **10** [8] was possible yielding **9** (41%) and **11** (40%). Thus, a (\pm)-chiral triphenylmethane core was obtained (Scheme 2).

As a precursor of the (\pm)-chiral tripodaphyrin derivative, compound **13** was prepared in 88% yield by coupling **11** and [5-(4-ethynylphenyl)-10,15,20-triphenylporphinato(2-)]zinc (**12**) [4] using triphenyl-

arsine as a ligand of the palladium catalyst instead of triphenylphosphine [7]. The porphyrin trimer **16** with three “arms” of different lengths (C_1 -symmetry) was synthesized after cleavage of both trialkylsilyl protecting groups in 85% yield and coupling with tetraphenylporphyrine **15** elongated with an iodotolane [4] unit (18%) (Scheme 3).

In conclusion, we developed a synthesis to obtain a new class of tripodaphyrins derivatives designed for nanofabrication. On one hand, we synthesized a C_{3v} -symmetric triphenylmethane in which three porphyrin macrocycles are attached to each phenyl ring. On the other hand, by using successive substitution of the three iodine atoms of tris(4-iodophenyl)methane,



Scheme 2

a trimer with three “arms” of different length was also obtained.

Experimental

All air- or water-sensitive reactions were carried out under Ar. Solvents were generally dried and distilled prior to use. Reactions were monitored by thin-layer chromatography on E. Merck silica gel 60F₂₅₄ (0.2 mm) precoated aluminum foils. Column chromatography (CC): E. Merck silica gel 60 (230–400 mesh). Melting points (m.p.) were determined with a hot stage apparatus (Thermovar, C. Reichert AG, Vienna) equipped with a digital thermometer. UV/VIS spectra were recorded on a Hewlett-Packard-8452A diode-array spectrophotometer; λ_{\max} in nm ($\log \epsilon$ in mol⁻¹ dm³ cm⁻¹). NMR: Varian Gemini 200 (¹H: 200.00 MHz; ¹³C: 50.30 MHz), Bruker-AM 360 (¹H: 360.14 MHz; or Bruker Avance DRX 500 (¹H: 500.13 MHz; ¹³C: 125.76 MHz); ¹H and ¹³C chemical shifts (δ) are given in ppm relative to TMS as internal standard, *J* values in Hz. Mass spectra: Vacuum Generators Micromass 7070E instrument equipped with a data system DS 11–250, FAB (fast atom bombardment): in 2-nitrobenzyl alcohol with Ar at 8 kV. Elemental analysis (C, H, N) were conducted by Ciba Specialities Mikrolabor, Marly, Switzerland, their results were found to be in good agreement ($\pm 0.2\%$) with the calculated values. Tetrakis(triphenylphosphine)palladium, tris(dibenzylideneacetone)dipalladium (Pd₂dba₃), triphenylarsine and tetrabutylammonium fluoride (TBAF) were purchased from Aldrich; dimethylformamide (DMF), tetrahydrofuran (THF), trimethylsilylacetylene (TMSA), triisopropylsilylacetylene, and other reagents from Fluka.

Trifluoromethanesulfonic acid methylidyne-4,1-phenylene ester (2) [10]

Trifluoromethanesulfonyl chloride (0.178 cm³, 1.68 mmol) was slowly added under an Ar atmosphere to a stirring mix-


ture of 149 mg 4,4',4''-methylidyne-trisphenol [5] (0.51 mmol), 0.348 cm³ Et₃N (2.5 mmol), and 6.2 mg DMAP (0.051 mmol) in 4.3 cm³ dry acetone at 0°C. After being stirred for 3 h at 0°C, the solvent was removed under reduced pressure and the crude product was purified by CC (CH₂Cl₂/*n*-hexane = 1/4) to yield 280 mg (80%) **2**. M.p.: 82°C; ¹H NMR (200.00 MHz, CDCl₃): δ = 5.64 (*s*, 1H, CH), 7.15 and 7.25 (AA'XX', 2 × apparent *d*, *J* = 8.9 Hz, 12H, H-2, H-3) ppm; ¹³C NMR (50.30 MHz, CDCl₃): δ = 54.7 (CH), 118.7 (*q*, *J* = 321 Hz, CF₃), 121.7 (C-2), 131.0 (C-3), 142.5 (C-4), 148.5 (C-1) ppm; FAB-MS: *m/z* (%) = 687 ([M – H]⁺, 19), 555 (25), 463 (100).

[Methylidyne-tris(4,1-phenylene-2,1-ethynediyl)] tris(triisopropylsilane) (3, C₅₂H₇₆Si₃)

Air was removed from a soln. of 82.5 mg **2** (0.12 mmol) in 7 cm³ DMF/Et₃N (5:1) by blowing Ar for 20 min. Then 41.7 mg Pd(PPh₃)₄ (0.036 mmol), 13.8 mg CuI (0.072 mmol), and 0.134 mm³ ethynyltriisopropylsilane (0.6 mmol) were added, and deaeration was continued for 10 min. Thereafter, the mixture was heated at 45°C for 18 h. The solvent was removed under reduced pressure and the crude product was purified by CC (*n*-hexane) to yield 89.6 mg (95%) **3**. M.p.: 167°C; ¹H NMR (200.00 MHz, CDCl₃): δ = 1.12 (*m*, 63H, Si-Pr₃), 5.49 (*s*, 1H, CH), 6.99 and 7.39 (AA'XX', 2 × apparent *d*, *J* = 8.2 Hz, 12H, H-2, H-3) ppm; ¹³C NMR (50.30 MHz, CDCl₃): δ = 11.3 (SiCH), 18.7 (CH(CH₃)₂), 56.3 (CH), 90.6 (C≡C-Si-Pr₃), 106.8 (C≡C-Si-Pr₃), 121.9 (C-1), 129.2 (C-3), 132.1 (C-2), 143.3 (C-4) ppm; FAB-MS: *m/z* (%) = 786 (M, 57), 742 (100).

Tris(4-ethynylphenyl)methane (4, C₂₅H₁₆)

To a solution of 23.7 mg **3** (30.2 μmol) in 8 cm³ THF, 18 mm³ TBAF (1 M in THF, 18 μmol) were added at 0°C, and the mixture was stirred at 0°C for 10 min. A few grains of CaCl₂ were added and the solvent was evaporated. The crude product was purified by CC (gradient from *n*-hexane to CHCl₃/*n*-hexane = 9/1) to yield 6.7 mg (70%) of **4**. ¹H NMR



Scheme 3

(360.14 MHz, CDCl_3): $\delta = 3.05$ (*s*, 3H, $\text{C}\equiv\text{CH}$), 5.52 (*s*, 1H, CH), 7.02 and 7.42 (AA'XX', $2 \times$ apparent *d*, $J = 8.1$ Hz, 12H, H-2, H-3) ppm; ^{13}C NMR (50.30 MHz, CDCl_3): $\delta = 56.3$ (CH), 77.3 ($\text{C}\equiv\text{C}-\text{H}$), 83.3 ($\text{C}\equiv\text{C}-\text{H}$), 120.5 (C-4), 129.3 (C-2), 132.3 (C-3), 143.6 (C-1) ppm; FAB-MS: m/z (%) = 317 (M^+ , 31), 215 (100).

*C*_{3v}-Symmetric Triporphyrin (**6**, $\text{C}_{157}\text{H}_{94}\text{N}_{12}\text{Zn}_3$)

Air was removed from a soln. of 2.2 mg **4** ($7 \mu\text{mol}$) and 22.7 mg [5-(4-iodophenyl)-10,15,20-triphenylporphyrinato (2-)]zinc (**5**) [**4**] ($28.2 \mu\text{mol}$) in 4.8 cm^3 DMF/*Et*₃N (5/1) by blowing Ar for 20 min. Then 0.79 mg Pd₂(dba)₃ ($0.86 \mu\text{mol}$) and 2.11 mg AsPh₃ ($6.89 \mu\text{mol}$) were added, and deaeration was continued for 10 min. Thereafter, the mixture was heated

at 35°C for 6 h. At that time, the same amount of Pd₂(dba)₃ and AsPh₃ was again added and the stirring continued for 3 h. The solvent was removed under reduced pressure and the crude product was purified by two successive CC ($\text{CHCl}_3/n\text{-hexane}$: gradient from 3/2 to 3/1) to yield 8.1 mg (49%) **6**. UV/VIS (CH_2Cl_2): λ_{max} ($\log \epsilon$) = 298 (4.94), 420 (6.10), 548 (4.69), 578 (3.96), 586 (4.14), 594 (3.69) nm ($\text{mol}^{-1} \text{ dm}^3 \text{ cm}^{-1}$); ^1H NMR (360.14 MHz, CDCl_3): $\delta = 5.72$ (*s*, 1H, CH), 7.28 and 7.70 (AA'XX', $2 \times$ apparent *d*, $J = 8.4$ Hz, 12H, H-phenylene), 7.72–7.81 (*m*, 27H, *m*- and *p*-H-phenyl), 7.96 (apparent *d*, $J = 8.0$ Hz, 6H, H-phenylene on porphyrin), 8.19–8.26 (*m*, 24H, *o*-H-phenyl and H-phenylene on porphyrin), 8.95 (*s*, 12H, β -H on porphyrin outside), 8.98 (*s*, 12H, β -H on porphyrin inside) ppm; ES⁺-MS (in $\text{CHCl}_3/$

HCOOH): $m/z = 1078.0$ ($[M-3Zn + 8H]^{2+}$), 719.2 ($[M-3Zn + 9H]^{3+}$) (calc. avg. mass for C₁₅₇H₉₄N₁₂Zn₃: 2344.69).

Tris(4-iodophenyl)methane (8) [6]

A soln. of 162.4 mg tris(4-iodophenyl)methanol (0.25 mmol) was refluxed in 98% HCOOH for 22 h. Thereafter, the soln. was diluted with CH₂Cl₂ and water. The mixture was neutralized with solid NaHCO₃, extracted three times with CH₂Cl₂, dried (Na₂SO₄), and evaporated. The crude product was purified by CC (CH₂Cl₂/*n*-hexane = 1/19) to yield 140.4 mg (89%) **8**. ¹H NMR (200.00 MHz, CDCl₃): $\delta = 5.35$ (*s*, 1H, CH), 6.80 and 7.61 (AA'XX', 2 × apparent *d*, *J* = 8.2 Hz, 12H, H-phenylene) ppm; ¹³C NMR (50.30 MHz, CDCl₃): $\delta = 55.4$ (CH), 92.3 (C-4), 131.2 (C-2), 137.6 (C-3), 142.3 (C-1) ppm.

[[4-[[Bis(4-iodophenyl)methyl]phenyl]ethynyl]-trimethylsilane (9, C₂₄H₂₂I₂Si)

Air was removed from a soln. of 130.3 mg tris(4-iodophenyl)methane (**8**) [6] (0.21 mmol) in 7.4 cm³ pyridine/*Et*₃N (1/1) by blowing Ar for 20 min. Then, 8.3 mg Pd(PPh₃)₄ (7.2 μmol), 4.4 mg CuI (23 μmol), and 26 mm³ TMSA (0.187 mmol) were added, and the mixture was stirred at 35°C for 2 h. The solvent was removed under reduced pressure and the residue was purified by CC (CH₂Cl₂/*n*-hexane = 1/19) to yield 45.4 mg (41%) **9**. ¹H NMR (360.14 MHz, CDCl₃): $\delta = 0.24$ (*s*, 9H, SiMe₃), 5.39 (*s*, 1H, CH), 6.79 and 7.60 (AA'XX', 2 × apparent *d*, *J* = 8.6 Hz, 8H, H-iodophenyl), 6.97 and 7.38 (AA'XX', 2 × apparent *d*, *J* = 8.2 Hz, 4H, H-phenylene) ppm; ¹³C NMR (50.30 MHz, CDCl₃): $\delta = -0.05$ (SiMe₃), 55.6 (central CH), 92.2 (C-4'), 94.5 (C≡C-SiMe₃), 104.6 (C≡C-SiMe₃), 121.6 (C-1), 129.1 (C-3), 131.3 (C-2'), 132.1 (C-2), 137.6 (C-3'), 142.5 (C1'), 143.0 (C-4) ppm; FAB-MS: m/z (%) = 592 (M⁺, 24), 591 ([M-H]⁺, 46), 577 ([M-CH₃]⁺, 69), 419 ([M-H-I-3CH₃]⁺, 95), 389 (100).

[[4-[[4-Iodophenyl]-4-[[4-[[4-[[triisopropylsilyl]ethynyl]phenyl]ethynyl]phenyl]ethynyl]phenyl]methyl]phenyl]ethynyl]trimethylsilane (11, C₅₁H₅₁ISi₂)

Air was removed from a soln. of 32 mg **9** (54 μmol) and 18.8 mg [[4-[(4-ethynylphenyl)ethynyl]phenyl]ethynyl]triisopropylsilane (**10**) [8] (49 μmol) in 3 cm³ pyridine/*Et*₃N (1/1) by blowing Ar for 20 min. Then, 3 mg Pd(PPh₃)₄ (2.6 μmol) and 2.2 mg CuI (12 μmol) were added, and the mixture was stirred at 35°C for 2 h. The solvent was removed under reduced pressure and the residue was purified by CC (gradient from *n*-hexane to CH₂Cl₂/*n*-hexane = 2/23) to yield 16.7 mg (40%) **11**. M.p.: 73°C; ¹H NMR (360.14 MHz, CDCl₃): $\delta = 0.24$ (*s*, 9H, SiMe₃), 1.13 (*m*, 21H, Si-Pr₃), 5.47 (*s*, 1H, CH), 6.82 and 7.61 (AA'XX', 2 × apparent *d*, *J* = 8.5 Hz, 4H, H-iodophenyl), 7.00 and 7.39 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-2, H-3), 7.03 and 7.44 (AA'XX', 2 × apparent *d*, *J* = 8.0 Hz, 4H, H-2', H-3'), 7.45 (*m*, 4H, H-2''', H-3'''), 7.49 (*m*, 4H, H-2'', H-3''), ¹³C NMR (50.30 MHz, CDCl₃): $\delta = -0.05$ (SiMe₃), 11.3 (SiCH), 18.7 (CH(CH₃)₂), 56.0 (central CH), 89.2, 90.9, 91.0, 91.1 (C≡C), 92.2 (C-4 iodophenyl), 93.0 (C≡C-Si-Pr₃), 94.4 (C≡C-SiMe₃), 104.7 (C≡C-SiMe₃), 106.6 (C≡C-Si-Pr₃), 121.4, 121.6 (C-1, C-4'''), 122.8 (C-1'',

C-4''), 123.2, 123.5 (C-4', C-1'''), 129.2, 129.4 (C-3, C-2'), 131.4 (C-2 iodophenyl), 131.5, 131.6, 131.8 (C-3', C2'', C-3'', C-2'''), 132.0, 132.1 (C-2, C-3'''), 137.6 (C-3 iodophenyl), 142.6 (C-1 iodophenyl), 143.2, 143.3 (C-4, C-1') ppm; FAB-MS: m/z (%) = 847 (M⁺, 100), 803 (53).

[[10,15,20-Triphenyl-5-[[4-[[4-[[4-[[4-[[triisopropylsilyl]ethynyl]phenyl]ethynyl]phenyl]ethynyl]-phenyl]-4-[[trimethylsilyl]ethynyl]phenyl]methyl]phenyl]ethynyl]-phenyl]porphinato(2-)]zinc (13, C₉₇H₇₈N₄Si₂Zn)

Air was removed from a soln. of 6.0 mg **11** (7.1 μmol) and 20 mg [5-(4-ethynylphenyl)-10,15,20-triphenylporphinato(2-)]zinc **12** [4] (28.5 μmol) in 2.4 cm³ of DMF/*Et*₃N (5/1) by blowing Ar for 20 min. Then 0.41 mg Pd₂(dba)₃ (0.45 μmol) and 1.09 mg AsPh₃ (3.56 μmol) were added, and deaeration was continued for 10 min. Thereafter, the mixture was heated at 40°C for 4 h. At that time, the same amount of Pd₂(dba)₃ and AsPh₃ was again added and the stirring continued for 3 h. The solvent was removed under reduced pressure and the crude product was purified by CC (CHCl₃/*n*-hexane: gradient from 2/3 to 11/9) to yield 8.9 mg (88%) **13**. UV/VIS (CH₂Cl₂): λ_{\max} (log ϵ) = 336 (4.86), 360 (4.72), 422 (5.68), 548 (4.34), 588 (3.84) nm (mol⁻¹ dm³ cm⁻¹); ¹H NMR (500.13 MHz, CDCl₃): $\delta = 0.26$ (*s*, 9H, SiMe₃), 1.13 (*s*, 21H, Si-Pr₃), 5.60 (*s*, 1H, CH), 7.09 and 7.45 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-trimethylsilylethynylphenyl), 7.12 and 7.50 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-phenylene-CH in big "leg"), 7.15 and 7.61 (AA'XX', 2 × apparent *d*, *J* = 8.2 Hz, 4H, H-phenylene-CH in porphyrin "leg"), 7.46 (*m*, 4H, H-triisopropylsilylethynylphenyl), 7.51 (*m*, 4H, H-central phenylene), 7.73–7.80 (*m*, 9H, *m*- and *p*-H-phenyl), 7.92 (apparent *d*, *J* = 8.2 Hz, 2H, H-phenylene on porphyrin), 8.21–8.23 (*m*, 8H, *o*-H-phenyl and H-phenylene on porphyrin), 8.96 (*m*, 8H, β -H on porphyrin) ppm; FAB-MS: m/z (%) = 1420.9 (calc. avg. mass for C₉₇H₇₈N₄Si₂Zn: 1421.26).

[5-[[4-[[4-[(4-Ethynylphenyl)-4-[[4-[[4-ethynylphenyl]ethynyl]phenyl]ethynyl]phenyl]methyl]phenyl]ethynyl]-10,15,20-triphenylporphinato(2-)]zinc (14, C₈₅H₅₀N₄Zn)

Compound **14** was obtained in 85% yield (6.4 mg), as described for **4**, from 8.9 mg **13** (6.3 μmol) for 8 min, after purification by CC (CHCl₃/*n*-hexane = 11/9). UV/VIS (CH₂Cl₂): λ_{\max} (log ϵ) = 334 (4.79), 354 (4.62), 420 (5.62), 548 (4.19), 590 (3.47) nm (mol⁻¹ dm³ cm⁻¹); ¹H NMR (360.14 MHz, CDCl₃): $\delta = 3.09$ (*s*, 1H, C≡CH), 3.18 (*s*, 1H, C≡CH in big "leg"), 5.61 (*s*, 1H, CH), 7.12 and 7.48 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-phenylene-CH), 7.13 and 7.50 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-phenylene-CH), 7.16 and 7.62 (AA'XX', 2 × apparent *d*, *J* = 8.3 Hz, 4H, H-phenylene-CH in porphyrin "leg"), 7.48 (*m*, 4H, H-phenylene in big "leg"), 7.51 (*m*, 4H, H-central phenylene in big "leg"), 7.72–7.80 (*m*, 9H, *m*- and *p*-H-phenyl), 7.92 (apparent *d*, *J* = 8.3 Hz, 2H, H-phenylene on porphyrin), 8.20–8.23 (*m*, 8H, *o*-H-phenyl and H-phenylene on porphyrin), 8.96 (*m*, 8H, β -H on porphyrin) ppm; FAB-MS: m/z (%) = 1193.1 (calc. avg. mass for C₈₅H₅₀N₄Zn: 1192.74).

*C*₁-Symmetric Triporphyrin (**16**, C₂₀₅H₁₁₈N₁₂Zn₃)

Air was removed from a soln. of 11 mg **14** (9.2 μmol) and 28.4 mg [5-[4-[4-[(4-iodophenyl)ethynyl]phenyl]ethynyl]phenyl]-10,15,20-triphenylporphinato(2-)]zinc (**15**) [4] (28.2 μmol) in 6.4 cm³ DMF/Et₃N (5:1) by blowing Ar for 20 min. Then 2.08 mg Pd₂(dba)₃ (2.28 μmol) and 5.5 mg AsPh₃ (18 μmol) were added, and deaeration was continued for 10 min. Thereafter, the mixture was heated at 40°C for 4 h. At that time, the same amount of Pd₂(dba)₃ and AsPh₃ was again added and the stirring continued for 3 h. The solvent was removed under reduced pressure and the crude product was purified after two successive CC (CHCl₃/*n*-hexane: gradient from 1/1 to 3/1) to yield 4.8 mg (18%) **16**. UV/VIS (CH₂Cl₂): λ_{max} (log ε) = 354 (5.15), 420 (6.06), 548 (4.69), 586 (4.14) nm (mol⁻¹ dm³ cm⁻¹); ¹H NMR (500.13 MHz, CDCl₃): δ = 5.65 (*s*, 1H, CH), 7.18 and 7.54 (2 × AA'XX', 4 × apparent *d*, *J* = 8.3 Hz, 8H, H-phenylene on central CH in medium and big "legs"), 7.20 and 7.65 (AA'XX', 2 × apparent *d*, *J* = 8.0 Hz, 4H, H-phenylene on central CH in small "leg"), 7.52–7.57 (*m*, 16H, H-phenylene in medium and big "legs"), 7.61 and 7.68 (2 × AA'XX', 4 × apparent *d*, *J* = 8.0 Hz, 8H, H-phenylene on porphyrin in medium and big "legs"), 7.73–7.81 (*m*, 27H, *m*- and *p*-H-phenyl), 7.94 (*m*, 6H, H-phenylene on porphyrin), 8.23 (*m*, 24H, *o*-H-phenyl and H-phenylene on porphyrin), 8.96 (*s*, 12H, β-H on porphyrin outside), 8.97 (*m*, 12H, β-H on porphyrin inside) ppm; ES⁺-MS (in CHCl₃/HCOOH): *m/z* (%) = 1378.7 ([M – 3Zn + 8H]²⁺), 919.3 ([M – 3Zn + 9H]³⁺) (calc. avg. mass for C₂₀₅H₁₁₈N₁₂Zn₃: 2945.41).

Acknowledgements

This work was supported in part by the Swiss National Science Foundation. NMR spectra were recorded on a Bruker

Avance DRX 500 instrument by F. Fehr; mass spectra were recorded by F. Nydegger and I. Müller.

References

- [1] For reviews see a) Burell AK, Officer DL, Plieger PG, Reid DCW (2001) *Chem Rev* **101**: 2751; b) Choi MS, Yamazaki T, Yamazaki I, Aida T (2004) *Angew Chem Int Ed* **43**: 150
- [2] See for example a) Carcel CM, Laha JK, Loewe RS, Tham Yongkit P, Schweikart KH, Misra V, Bocian DF, Lindsey JS (2004) *J Org Chem* **69**: 6739; b) Balasubramanian T, Lindsey JS (1999) *Tetrahedron* **55**: 6771; c) Lee CH, Li F, Iwamoto K, Dadok J, Bothner-By AA, Lindsey JS (1995) *Tetrahedron* **51**: 11645
- [3] Mongin O, Gossauer A (1996) *Tetrahedron Lett* **37**: 3825
- [4] Mongin O, Gossauer A (1997) *Tetrahedron* **53**: 6835
- [5] Vögtle F, Berscheid R, Schnick W (1991) *J Chem Soc Chem Commun* 414
- [6] The synthesis described in the Experimental Section yields tris(4-iodophenyl)methane in higher purity than the procedure reported in Ref. [9] in which direct iodination of triphenylmethane leads in fact to tris(4-iodophenyl)methanol as the major product and a mixture containing tris(4-iodophenyl)methane
- [7] Farina V, Krishnan B (1991) *J Am Chem Soc* **113**: 9585
- [8] Lavastre O, Ollivier L, Dixneuf PH (1996) *Tetrahedron* **52**: 5495
- [9] Popova NI, Skubnevskaya GI, Molin Yu N, Kotlyarevskii IL (1969) *Izv Akad Nauk SSSR, Ser Khim* **11**: 2424
- [10] Satoshi M, Takanobu N, Yoshiaki T (2005) *PCT Int Appl*: 82. WO 2005030829 CAN **142**: 356334 AN 2005: 300497