

Li Tao · Li Jianping · Zhao Zhiwei

Arbuscular mycorrhizas in a valley-type savanna in southwest China

Received: 15 May 2003 / Accepted: 18 September 2003 / Published online: 28 October 2003
© Springer-Verlag 2003

Abstract The arbuscular mycorrhizal (AM) status of 67 plant species in a savanna community in the hot, dry valley of Jinsha River, southwest China was surveyed. It was found that about 95% of the plant species formed AM and 5% possibly formed AM. The composition of AM fungi (AMF) in the rhizosphere soils was also investigated. The AMF spore density ranged from 5 to 6,400 per 100 g soil, with an average of 1,530, and these spores/sporocarps were identified as belonging to six genera. Fungi belonging to the genera *Glomus* and *Acaulospora* were the dominant AMF. High densities of AMF spores in the rhizosphere soils, and the intensive colonization of the plant roots, indicated that plants grown in this valley-type savanna may be highly dependent on AM.

Keywords Arbuscular mycorrhiza · Savanna · Hot-arid ecosystems · Rhizosphere soils

Introduction

Arbuscular mycorrhizas (AM) are important components in savanna ecosystems, as arbuscular mycorrhizal fungi (AMF) can contribute to plant growth by reducing stresses resulting from nutrient deficiencies (mainly P) and drought (Sanginga et al. 1999; Augé 2001). Beneficial effects on plant growth, biomass, drought resistance, and ecological restoration by inoculation with AMF have been reported with herbaceous host plants both in controlled environments using potted plants and in the field (Rao and Tarafdar 1993; Tarafdar and Praveen-Kumar 1996; Estaun et al. 1997; Gemma and Koske 1997; Cuenca et al. 1998; Smith et al. 1998; Frost et al. 2001; Caravaca et al. 2003).

According to the vegetation classification of Walter (1979), a savanna usually develops where the evaporation is about 3–6 times the precipitation, 80–90% of precipitation is concentrated in the wet season, and the annual mean temperature is over 20°C. The characteristics of savanna vegetation are grasslands that are yellow in the dry season of the year, with rare trees. Typical savanna vegetation is distributed mainly in Africa. Valley-type savanna is a special type that develops in river valleys, such as those of different rivers in southwest China. The plant community developed in the hot-dry valley of Jinsha River, Yuamou County (101°35'E-102°06', 25°23'N-26°06') is the most typical of this special type of savanna. The annual average temperature is 21.8°C, the highest up to 43°C, and the mean annual precipitation is 634.0 mm, mainly between June and October. The wet season accounts for over 90% of the precipitation, while annual evaporation reaches 3,847.8 mm, about 6.1 times precipitation (data from Yuanmou County weather station over 24 years). Grasses form the background vegetation, which is withered and yellow in the dry season, and a scattering of trees and shrubs are distributed among the grassland. The dominant grass species are *Heteropogon contortus*, *Bothriochloa pertusa* and *Eriophorum comosum*, and the woody plants are *Phyllanthus emblica*, *Dodonaea angustifolia*, *Vitex negundo f. laxip* and *Jatropha curcas*. The arid soil of this area has the following chemical characteristics: total N 0.10±0.06 (N)%, total P 0.18±0.13 (P₂O₅)%, available P 2.8±0.7 (P₂O₅) mg/100 g soil, total K 2.1±0.7 (K₂O)%, and pH 8.2±0.6 (unpublished data, from 21 soil samples).

The Jinsha River is the upper reach of the Yangtse River, the longest river in China. It winds through the valleys of Yunnan and Sichuan provinces, southwest China, which are densely populated. Human activities have intensively interfered with the valley-type savanna ecosystem and the savanna is now facing desertification (Jin and Ou 2000). Much research has been done on the origin, development, evolution, structure and function of the savanna vegetation (Jin and Ou 2000; Jin 2002), but no reports on AM have been made. Here, we report the

L. Tao · L. Jianping · Z. Zhiwei (✉)
Laboratory for Conservation and Utilization of Bio-resources,
Yunnan University,
650091 Kunming, P.R. China
e-mail: Zhaozhw@ynu.edu.cn
Tel.: +86-871-5034799
Fax: +86-871-5034838

AM status of the plants and the AMF composition in the valley-type savanna of the Jinsha River.

Materials and methods

Roots and their rhizosphere soils were collected to a depth of 5–30 cm in July 2002 (wet season), making sure that the roots were connected to sampled plants and cleaning the trowels between samples. Part of the root system of each plant was fixed in 5 ml formalin, 5 ml acetic acid and 90 ml 70% alcohol, diluted 1:1 with water (1/2 FAA), and stored at 4°C. The remaining roots were air-dried with their rhizosphere soil (about 500 g) for 2 weeks, and then

stored in sealed plastic bags at room temperature for up to 2 months until samples could be treated. Roots were taken from the 1/2 FAA, washed several times in tap water and cleared in 10% (w/v) KOH by heating to approximately 90°C in a water bath for 2–3 h, the time depending on the size/structure of the roots and their pigmentation. The cooled root samples were washed and cut into 0.5–1 cm segments and stained with 0.5% acid fuchsin according to the method of Berch and Kendrick (1982). Fifty 0.5–1 cm root fragments were examined per sample for their AM status under a compound microscope ($\times 160$ – $\times 880$).

The rhizosphere soil samples were wet-sieved for spores using the method described by An et al. (1990). Soil (20 g) from each plant rhizosphere was independently suspended in 150 ml water, stirred with a magnetic stirrer for 10 min, sieved using 40 μ m, 70 μ m, 100 μ m and 150 μ m sieves with tap water, filtered onto a

Table 1 Arbuscular mycorrhizal (AM) status of plants and spore densities of arbuscular mycorrhizal fungi (AMF) in the valley-type savanna soils. *M* AM status of plants (+ mycorrhizal, \pm possibly mycorrhizal), *ASD* AMF spore density (number of AMF spores in 100 g soil from the corresponding plant rhizosphere), *ASD*^a average AMF spore density

Host plants	M	ASD	Host plants	M	ASD ^a
Acanthaceae			Gramineae		
<i>Barleria cristata</i>	+	235	<i>Bothriochloa pertusa</i>	+	2,130
Adiantaceae			<i>Heteropogon contortus</i>	+	555
<i>Adiantum philipense</i>	+	1,240	<i>Neyraudia reynaudiana</i>	+	3,210
Agavaceae			<i>Paspalidium flavidum</i>	+	5
<i>Agave americana</i>	+	525	<i>Tripogon filiformis</i>	+	1,340
Amaranthaceae			Labiatae		
<i>Achyranthes aspera</i>	+	75	<i>Elsholtzia cypriani</i>	+	1,860
<i>Amaranthus spinosus</i>	+	10	<i>Ocimum basilicum</i>	+	1,810
<i>A. viridis</i>	+	1,750	Malvaceae		
Asclepiadaceae			<i>Abutilon indicum</i>	+	1,065
<i>Calotropis gigantea</i>	+	4,355	<i>Sida chinensis</i>	\pm	475
Bignoniaceae			<i>S. szechuensis</i>	+	2,310
<i>Incarvillea arguta</i>	+	115	Mimosaceae		
Bombacaceae			<i>Acacia farnesiana</i>	+	4,695
<i>Bombax malabaricum</i>	+	2,075	<i>Albizia kalkora</i>	+	320
Boraginaceae			Moraceae		
<i>Cynoglossum lanceolatum</i>	+	4,375	<i>Broussonetia papyrifera</i>	+	2,220
Caesalpiniaceae			Myrtaceae		
<i>Tamarindus indica</i>	+	1,555	<i>Eucalyptus citriodora</i>	+	70
Chenopodiaceae			Nyctaginaceae		
<i>Chenopodium ambrosioides</i>	+	595	<i>Boerhavia diffusa</i>	\pm	220
Compositae			Papaveraceae		
<i>Artemisia cononocephala</i>	+	700	<i>Macleaya cordata</i>	+	90
<i>Bidens pilosa</i>	+	1,445	Papilionaceae		
<i>Blainvillea acmella</i>	+	955	<i>Alysicarpus vaginalis</i>	+	580
<i>Conyza blinii</i>	+	1,380	<i>Atylosia scarabaeoides</i>	+	2,965
<i>C. canadensis</i>	+	490	<i>Crotalaria medicaginea</i>	+	80
<i>Laggera pterodonta</i>	+	1,810	<i>Desmodium microphyllum</i>	+	1,125
<i>Parthenium hysterophorus</i>	+	1,295	<i>Tephrosia purpurea</i>	+	535
<i>Siegesbeckia orientalis</i>	+	1,435	<i>Vigna aconitifolius</i>	+	30
<i>Sonchus oleraceus</i>	+	4,560	Polygonaceae		
<i>Xanthium sibiricum</i>	+	245	<i>Polygonum statice</i>	+	330
<i>Zinnia elegans</i>	+	3,255	<i>Rumex hastatus</i>	+	570
Cyperaceae			Rhamnaceae		
<i>Cyperus niveus</i>	+	1,560	<i>Ziziphus mauritiana</i>	+	1,815
<i>C. rotundus</i>	+	2,175	Sapindaceae		
<i>Eriophorum comosum</i>	\pm	2,355	<i>Dodonaea viscosa</i>	+	2,210
Equisetaceae			Selaginellaceae		
<i>Equisetum diffusum</i>	+	2,015	<i>Selaginella pulvinata</i>	+	6,400
Euphorbiaceae			<i>S. mairei</i>	+	2,665
<i>Acalypha acmophylla</i>	+	530	Solanaceae		
<i>Euphorbia heterophylla</i>	+	780	<i>Datura stramonium</i>	+	120
<i>E. hirta</i>	+	975	<i>Solanum khasianum</i>	+	65
<i>E. royleana</i>	+	765	Tiliaceae		
<i>E. thymifolia</i>	+	2,645	<i>Grewia biloba</i>	+	2,900
<i>Phyllanthus urinaria</i>	+	3,330	Verbenaceae		
Gesneriaceae			<i>Vitex negundo</i>	+	1,530
<i>Corallodiscus flabellatus</i>	+	4,505	Zygophyllaceae		
			<i>Tribulus terrester</i>	+	80

Total: plants =67 species; M (“+” =64, about 95%; “ \pm ” =3, about 5%); total spore specimens =102,485; ASD^a=1,530

filter paper, and then placed in a 9 cm Petri dish for examination under a binocular stereomicroscope. AMF spores were counted in the four sieved samples. Some spores were tightly grouped in a sporocarp and it was difficult to count the number of spores per sporocarp; in these cases, a sporocarp was referred to as one spore.

Each spore type was mounted sequentially in water, lactophenol, PVA (polyvinyl lactic acid) and Melzer's reagent (Morton 1988), for identification. Identification was based on spore color, size, surface ornamentation and wall structure, with reference to the descriptions provided by the *International collection of vesicular and arbuscular mycorrhizal fungi* (<http://invam.caf.wvu.edu>) and the original species descriptions (Trappe 1977; Rothwell and Trappe 1979; Nemeček et al. 1981; Walker and Trappe 1981; Smith and Schenck 1985; Morton 1988).

Results

Under the microscope it was noted that hyphae, hyphal coils, vesicles and arbuscules were stained red while the root cortical cells were unstained or were pale red. If at least one root segment was found to contain arbuscules or vesicles, then the plant was noted as an AM plant, recorded as +. If the root cortex was found to be colonized by fungal mycelia, but without arbuscules or vesicles, the corresponding plant was noted as possibly AM, recorded as ±. Plants were recorded as non-mycorrhizal (–) when neither arbuscules/vesicles nor fungal mycelia were detected in their root cortical cells. Of the 67 plant species surveyed, 64 (about 95%) formed AM, and the roots were usually intensively colonized by the hyphae in these species. The possibly mycorrhizal species represented 5% (three samples), and no non-mycorrhizal plants were found. Furthermore, some plant species in families

that are traditionally considered as non-mycorrhizal or rarely forming mycorrhizas were also colonized by AMF. For example, *Rumex hastatus* and *Polygonum sticticum* (Polygonaceae) were typically colonized by AMF (Table 1). Arbuscules were especially rich in the root cortical cells of *Parthenium hysterophorus*, *Zinnia elegans*, *Incarvillea arguta*, *Acalypha acmophylla*, *Abutilon indicum*, *Phyllanthus urinaria* and *Eucalyptus citriodora*. There were vesicles in *Xanthium sibiricum*, *Sida zechuensis*, *Artemisia codonocephala*, *Heteropogon contortus*, *Selaginella pulvinata* and *Bombax malabaricum*. Both arbuscules and vesicles displayed a high diversity in their morphologies.

AMF spores were counted in the 67 rhizosphere soil samples. The spore density was highly uneven, ranging from 5 to 6,400 spores per 100 g rhizosphere soil. The average spore density was 1,530 (Table 1). From the 67 rhizosphere soil samples, 844 AMF spore (or sporocarp) samples were obtained by wet-sieving. Some spore types were identified to species by their morphological characters (Fig. 1) but most of the specimens were identified only to genus level because they lacked distinguishable, fine taxonomic characters. Accordingly, the frequency of AMF occurrence could be statistically analyzed only at the genus level. *Acaulospora* and *Glomus* were the dominant genera; their frequencies of occurrence were 22.4% and 67.0%, respectively, in all the spore samples examined. The frequency of occurrence of AMF from the other four genera was very low (Table 2).

Fig. 1 Some identified species of AMF. 1. *Acaulospora sinosa* 2. *A. bireticulata* 3. *Glomus constrictum* 4. *G. coremioides* 5. *G. sinuosa* 6. *G. rubiformis*. Bar=30µm




Table 2 Identified AMF genera and the frequency of occurrence in rhizosphere soils. *IT* Number of times corresponding AMF isolated from the soils, *F* frequency of occurrence

AMF genus	IT	F (%)
<i>Acaulospora</i>	189	22.4
<i>Archaeospora</i>	16	1.9
<i>Entrophospora</i>	9	1.1
<i>Gigaspora</i>	21	2.5
<i>Glomus</i>	566	67.0
<i>Scutellospora</i>	43	5.1
Total	844	100

Discussion

It has been demonstrated that indigenous AMF can increase the growth of their host plants in hot-dry areas (Requena et al. 2001). Water is a major stress for plants grown in hot-dry valley ecosystems. Though mycorrhizal effects on plant water relations are not as direct and consistent as those on P acquisition and host growth, the results of this study suggest that plants in the valley-type savanna may have a strong dependency on AM, as 95% of the surveyed plants formed AM, and their roots were intensively colonized. There were apparently more arbuscules and vesicles in the roots of the hot-dry valley plants than generally found in previous studies in tropical areas. Onguene and Kuyper (2001) reported that 79% of the samples surveyed in the rain forest of South Cameroon were colonized by AMF. AM plant species were only 56% of the total number of species sampled in Xishuangbanna rain forest in southwest China (Zhao et al. 2001). In the present study, most of the plants surveyed in the families of Brassicaceae, Polygonaceae, Cyperaceae, etc., which are usually considered to be non-mycorrhizal, also commonly formed AM; examples were *Cyperus rotundus*, *Polygonum stitice* and *Rumex hastatus*. A similar situation has been found in the savanna of Venezuela (Lovera and Cuenca 1996) and in Indian semi-arid tropical grassland (Muthukumar and Udaiyan 2002), and indicates that plants grown in these stress environments may depend more on AM.

It has been reported that both spore production and species richness of AMF are lower in arid climates than in other ecosystems (Rose 1981; Pond et al. 1984), and decrease as aridity increases (Stahl and Christensen 1982; Stutz and Morton 1996). However, a short-term, transient decrease in soil moisture may promote spore production (Jacobson 1997; Guadarrama and Alvarez-Sanchez 1998). An average spore density of 1,530 spores per 100 g soil in our study was higher than previous reports in some other areas (Walker et al. 1982; Sylvia 1986; Friese and Koske 1991; Bever et al. 1996). Our results strongly support the hypothesis of Redhead (1975) that the amount of soil water optimal for plant growth may also be optimal for AMF sporulation. The samples were collected in the wet season (plant growth season) in this study, and the abundance of AMF spores and colonization has been reported to be coordinated with growth stages of plants

during the long stress of a strong regimen of drought and rain (Kennedy et al. 2002). Some researchers have also shown that much less sporulation occurs in either extremely dry soils (Stahl and Christensen 1982; Stutz and Morton 1996) or flooded and permanently water-logged soils (Khan 1974).

AMF in the genera *Acaulospora* and *Glomus* are dominant in the valley-type savanna of the Jinsha River, with frequencies of occurrence around 90%. *Glomus* species with sporocarp and periderm, such as *Glomus sinuosa*, and the small-spored *Acaulospora* and *Glomus* species were the most common. Stutz et al. (2000) also reported that AMF in arid regions of southwestern North America and Namibia, Africa were limited mostly to small-spored fungi in Glomaceae and Acaulosporaceae. It is possible that the plants in valley-type savanna and arid regions have a short growth period in the year, and that formation of small spores takes a shorter time than that of large spores in AMF in Gigasporaceae, which may not have time to form and mature prior to the dry season (Brundrett et al. 1999; Boddington and Dodd 2000).

Acknowledgements We thank the two anonymous reviewers for suggesting improvements in the manuscript. Prof. F.A. Smith (University of Adelaide, Australia) is thanked for his constructive suggestions and helping us with improving the English. The authors gratefully acknowledge Xia Yongmei, associate researcher of Xishuangbanna Tropical Botanical Garden, CAS, who helped us to identify all plant specimens. We also are very grateful to Cheng Lizhong, Sha Tao, Wang Guohua and Yang Lin who helped collect some samples. This research was financially supported by the National Natural Science Foundation of China (NSFC: 30360003) and Natural Science Foundation of Yunnan Province (2001C0001Z).

References

- An Z-Q, Hendrix JW, Hershman DE, Henson GT (1990) Evaluation of the "most probable number" (MPN) and wet-sieving methods for determining soil-borne populations of endogoneous mycorrhizal fungi. *Mycologia* 82:516–518
- Augé RM (2001) Water relation, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza* 11:3–42
- Berch SM, Kendrick B (1982) Vesicular-arbuscular mycorrhizae of southern Ontario ferns and fern-allies. *Mycologia* 74:769–776
- Bever JD, Morton JB, Antonovics J, Schultz PA (1996) Host-dependent sporulation and species diversity of arbuscular mycorrhizal fungi in a mown grassland. *J Ecol* 84:71–82
- Boddington CL, Dodd JC (2000) The effect of agricultural practices on the development of indigenous arbuscular mycorrhizal fungi. I. Studies in an Indonesian ultisol. *Plant Soil* 218:137–144
- Brundrett M, Abbott L, Jasper D (1999) Glomalean mycorrhizal fungi from tropical Australia. I. Comparison of the effectiveness and specificity of different isolation procedures. *Mycorrhiza* 8:305–314
- Caravaca F, Barea JM, Palenous J, Figueroa D, Alguacil MM, Roldan A (2003) Establishment of shrub species in a degraded semiarid site after inoculation with native or allochthonous arbuscular mycorrhizal fungi. *Appl Soil Ecol* 22:103–111
- Cuenca G, Andrade ZD, Escalante G (1998) Arbuscular mycorrhizae in the rehabilitation of fragile degraded tropical lands. *Biol Fertil Soils* 26:107–111
- Estau V, Save R, Biel C (1997) AM inoculation as a biological tool to improve plant revegetation of a disturbed soil with

- Rosmarinus officinalis* under semi-arid conditions. *Appl Soil Ecol* 6:223–229
- Friese CF, Koske RE (1991) The spatial dispersion of spores of vesicular-arbuscular mycorrhizal fungi in a sand dune: microscale patterns associated with the root architecture of American beachgrass. *Mycol Res* 95:952–957
- Frost SM, Stahl PD, Williams SE (2001) Long-term reestablishment of arbuscular mycorrhizal fungi in a drastically disturbed semiarid surface mine soil. *Arid Land Res Manage* 15:3–12
- Gemma JN, Koske RE (1997) Arbuscular mycorrhizae in sand dune plants of the North Atlantic coast of the United States: field and greenhouse inoculation and presence of mycorrhizae in planting stock. *J Environ Manage* 50:251–264
- Guadarrama P, Alvarez-Sanchez FJ (1998) Abundance of arbuscular mycorrhizal fungi spores in different environments in a tropical rain forest, Veracruz, Mexico. *Mycorrhiza* 8:267–270
- Jacobson KM (1997) Moisture and substrate stability determine VA-mycorrhizal fungal community distribution and structure in an arid grassland. *J Arid Environ* 35:59–75
- Jin Z-z (ed) (2002) Floristic features of dry-hot and dry-warm valleys, Yunnan and Sichuan. Yunnan Science and Technology Press, Kunming, pp 1–12
- Jin Z-z, Ou X-k (eds) (2000) Yuanjiang, Nujiang, Jinshajiang, Lancangjiang Vegetation of dry-hot valley. Yunnan University Press, Kunming, pp 141–302
- Kennedy LJ, Tiller RL, Stutz JC (2002) Associations between arbuscular mycorrhizal fungi and *Sporobolus wrightii* in riparian habitats in arid South-western North America. *J Arid Environ* 50:459–475
- Khan AG (1974) The occurrence of mycorrhizas in halophytes, hydrophytes and xerophytes, and of *Endogone* spores in adjacent soils. *J Gen Microbiol* 81:7–14
- Lovera M, Cuenca G (1996) Arbuscular mycorrhizal infection in Cyperaceae and Gramineae from natural, disturbed and restored savannas in La Gran Sabana, Venezuela. *Mycorrhiza* 6:111–118
- Morton JB (1988) Taxonomy of VA mycorrhizal fungi: classification, nomenclature and identification. *Mycotaxon* 32:267–324
- Muthukumar T, Udaiyan K (2002) Seasonality of vesicular-arbuscular mycorrhizae in sedges in a semi-arid tropical grassland. *Acta Oecologica Int J Ecol* 23:337–347
- Nemec S, Menge JA, Platt RG, Johnson ELV (1981) Vesicular-arbuscular mycorrhizal fungi associated with citrus in Florida and California and Notes on their distribution and ecology. *Mycotaxon* 73:112–127
- Onguene NA, Kuyper TW (2001) Mycorrhizal associations in the rain forest of South Cameroon. *For Ecol Manage* 140:277–287
- Pond EC, Menge JA, Jarrel WM (1984) Improved growth of tomato in salinized soil by vesicular-arbuscular mycorrhizal fungi collected from saline soils. *Mycologia* 76:74–84
- Rao AV, Tarafdar IC (1993) Role of VAM fungi in nutrient-uptake and growth of clusterbean in an arid soil. *Arid Soil Res Rehabil* 7:275–280
- Redhead JF (1975) Endotrophic mycorrhizas in Nigeria: some aspects of the ecology of the endotrophic mycorrhizal association of *Khaya grandifolia* C.D.D. In: Sanders FE, Mosse B, Tinker PB (eds) *Endomycorrhizas*. Academic Press, London, pp 447–459
- Requena N, Perez-Solis E, Azcon-Aguilar C, Jeffries P, Barea JM (2001) Management of indigenous plant-microbe symbioses aids restoration of desertified ecosystems. *Appl Environ Microbiol* 67:495–498
- Rose SL (1981) Vesicular-arbuscular mycorrhizal associations of some desert plants of Baja California. *Can J Bot* 59:1056–1060
- Rothwell FM, Trappe JM (1979) *Acaulospora bireticulata* sp. nov. *Mycotaxon* 8:471–475
- Sanginga N, Carsky RJ, Dashiell K (1999) Arbuscular mycorrhizal fungi respond to rhizobial inoculation and cropping systems in farmers' fields in the Guinea savanna. *Biol Fertil Soils* 30:179–186
- Smith GS, Schenck NC (1985) Two new dimorphic species in the endogonaceae: *Glomus ambisporum* and *Glomus heterosporum*. *Mycologia* 77:566–574
- Smith MR, Charvat I, Jacobson RL (1998) Arbuscular mycorrhizae promote establishment of prairie species in a tallgrass prairie restoration. *Can J Bot* 76:1947–1954
- Stahl PD, Christensen M (1982) Mycorrhizal fungi associated with *Bouteloua* and *Agropyron* in Wyoming sagebrush-grassland. *Mycologia* 74:877–885
- Stutz JC, Morton JB (1996) Successive pot cultures reveal high species richness of arbuscular endomycorrhizal fungi in arid ecosystems. *Can J Bot* 74:1883–1889
- Stutz JC, Copeman R, Martin CA, Morton JB (2000) Patterns of species composition and distribution of arbuscular mycorrhizal fungi in arid regions of southwestern North America and Namibia, Africa. *Can J Bot* 78:237–245
- Sylvia DM (1986) Spatial and temporal distribution of vesicular-arbuscular mycorrhizal fungi association with *Uniola paniculata* in Florida foredunes. *Mycologia* 78:728–734
- Tarafdar JC, Praveen-Kumar (1996) The role of vesicular arbuscular mycorrhizal fungi on crop, tree and grasses grown in an arid environment. *J Arid Environ* 34:197–203
- Trappe JM (1977) Three new endogonaceae: *Glomus constrictum*, *Sclerocystis clavisporea*, and *Acaulospora scrobiculata*. *Mycotaxon* 6:359–366
- Walker C, Trappe JM (1981) *Acaulospora spinosa* sp. nov. with a key to the species of *Acaulospora*. *Mycotaxon* 12:515–521
- Walker C, Mize CW, McNabb HS (1982) Population of endogonaceous fungi at two locations in central Iowa. *Can J Bot* 60:2518–2529
- Walter H (1979) *Vegetation of earth-ecological systems of the geobiosphere*, 2nd edn. Springer, New York Berlin Heidelberg, pp 72–130
- Zhao Z-W, Xia Y-M, Qin X-Z, Li X-W, Cheng L-Z, Sha T, Wang G-H (2001) Arbuscular mycorrhizal status of plants and the spore density of arbuscular mycorrhizal fungi in the tropical rain forest of Xishuangbanna, southwest China. *Mycorrhiza* 11:159–162