

## Short Communication

***Westerdykella globosa*, a proposal for a new combination**

Tadayoshi Ito and Akira Nakagiri

Institute for Fermentation, Osaka, 17-85, Juso-honmachi 2-chome, Yodogawa-ku, Osaka 532, Japan

Accepted for publication 3 June 1995

The surface ornamentation of ascospores of *Preussia globosa* was compared in an isolate from paddy soil in Japan and a culture derived from the holotype. The ascospores of two cultures were characterized by the surface ornamentation of a single, semicircular spiral ridge. This new finding strongly suggested that the fungus should be transferred to the genus *Westerdykella*. Therefore, the morphological and cultural characters of the fungus were re-examined, and the new combination *Westerdykella globosa* is herein proposed.

Key Words—*Preussia globosa*; *Pseudeurotium globosum*; soil fungus; *Westerdykella globosa*.

Rai and Tewari (1962) originally described and precisely illustrated the species *Pseudeurotium globosum* Rai & Tewari (described as "*Pseudoeurotium*") having multispore asci and large globose ascospores. They soon transferred this species to *Preussia globosa* (Rai & Tewari) Rai & Tewari (1963) because the ascospore cells are readily separable at maturity. Other species in the genus *Preussia* Fuckel, however, have transverse septa and obvious germ slits as described by Cain (1961) and Arx and Storm (1967).

During an ecological study of Japanese soil fungi, a strain isolated from paddy soil was first identified as *P. globosa* (IFO 32630).

The Japanese isolate and a culture derived from the holotype of *Preussia globosa* (IFO 32588=IMI 082625) were recently compared in their morphology. It was found that their ascospores have a single, semicircular spiral ridge on the surface. Furthermore, their asci were confirmed to contain 32 ascospores by dissecting with a Skerman-type micromanipulator.

The above characteristics of *Preussia globosa* were identical with the generic concept of *Westerdykella* (Stolk, 1955), the type species *Westerdykella ornata* Stolk, which produces 32 ascospores in an ascus which are ornamented on the surface with a spiral ridge. We concluded, therefore, that *Preussia globosa* should be transferred to the genus *Westerdykella*.

***Westerdykella globosa*** (Rai & Tewari) T. Ito & Nakagiri, comb. nov. Figs. 1–4

≡ *Pseudeurotium globosum* Rai & Tewari, Mycopathol. Mycol. Appl. 16: 90. 1962.

≡ *Preussia globosa* (Rai & Tewari) Rai & Tewari, Proc. Indian Acad. Sci. B 57: 53. 1963.

Colonies on oatmeal agar growing rapidly, attaining a diameter of 90 mm within 3 weeks at 24°C, velvety, thin, partly immersed, pale olive-buff (Rayner, 1970);


reverse pale olive-buff. Ascumata discrete or confluent, non-ostiolate, maturing within one month, globose to subglobose, 90–260 µm in diam, carbonaceous, dark brown; peridium 1–2-layered, consisting of brown, thick-walled, pseudoparenchymatous cells; ascumatal initials appeared in the form of intercalary segments of the vegetative hyphae, consisting of thick-walled, dictyoid cells. Asci in short chains, 32-spored, subglobose to ovoid, 20–24 × 14–17 µm, evanescent. Ascospores bound together in groups at first, becoming separate at maturity, dark brown, globose to subglobose, 4–6 µm in diam, thick-walled, smooth, with a single, semicircular spiral ridge on the surface.

At 37°C, growth is nil.

Cultures examined: IFO 32630 (T. Ito H2-3-5-20), a culture isolated from paddy soil, Ikeda, Osaka, Japan, April 1990 and a culture derived from the holotype of *Preussia globosa* (IFO 32588=IMI 082625).

Colonies on the following media after three-week incubation at 24°C: On potato-carrot agar growing rapidly, attaining a diameter of 90 mm, velvety, thin, partly immersed, hyaline; reverse uncolored. Ascumata are produced abundantly on the surface or submerged in the medium. On malt extract agar growing rapidly, attaining a diameter of 80 mm, velvety, thin, immersed, pale olive-buff; reverse pale olive-buff. On yeast phosphate soluble starch agar growing rapidly, attaining a diameter of 80 mm, floccose, white to pale gray at the center; reverse pale olive-buff to gray or mouse gray. Abundant ascumata are produced on the surface or submerged in the medium.

The fungus was first isolated from soil of a stream bank in India by Rai and Tewari. The Japanese isolate was found in paddy soil in Osaka as mentioned above. Recently, two strains (T. Ito H6H24-106-2, H645-114-5; 19 Oct. 1994) were isolated from paddy and pasture soils at Iriomote Isl. in Okinawa. Places where these



Figs. 1–4. *Westerdykella globosa* (IFO 32630). 1. Ascoma. 2. Asci. 3. Ascospores with a single, semicircular spiral ridge (arrows). 4. Ascospores of the strain derived from the holotype (IFO 32588). Bars: 1=50  $\mu\text{m}$ , 2=20  $\mu\text{m}$ , 3 and 4=5  $\mu\text{m}$ .

strains have been detected belong to the tropical, subtropical and temperate zones, respectively. The fungus seems to be of worldwide distribution.

**Acknowledgement**—We are indebted to the International Mycological Institute, Kew, England for providing a culture der-

ived from the holotype of *Preussia globosa*.

**Literature cited**

Arx, J. A. von, and Storm, P. K. 1967. Über einige aus dem Erdboden isolierte, zu *Sporormia*, *Preussia* und *Westerdykella* gehorende Ascomyceten. *Persoonia* 4: 407–415.

- Cain, R. F. 1961. Studies of coprophilous Ascomycetes VII. *Preussia*. Can. J. Bot. **39**: 1633-1666.
- Rai, J. N. and Tewari, J. P. 1962. *Pseudoeurotium globosum*, a new Ascomycete from Indian soils. Mycopathol. Mycol. Appl. **16**: 90-96.
- Rai, J. N. and Tewari, J. P. 1963. On some isolates of the genus *Preussia* Fuckel from Indian soils. Proc. Indian Acad. Sci. B **57**: 45-55.
- Rayner, R. W. 1970. "A mycological colour chart," Commonwealth Mycological Institute, Kew, Surrey and British Mycological Society.
- Stolk, A. C. 1955. *Emericellopsis minima* and *Westerdykella ornata*. Trans. Br. Mycol. Soc. **38**: 419-424.