

Available online at www.sciencedirect.com**MYCOSCIENCE**

ISSN 1340-3540 (print), 1618-2545 (online)

journal homepage: www.elsevier.com/locate/myc**Full paper**

Diversity and ubiquity of xylariaceous endophytes in live and dead leaves of temperate forest trees

Takashi Osono^{a,*}, Osamu Tateno^b, Hayato Masuya^c^aCenter for Ecological Research, Kyoto University, Otsu, Shiga 520-2113, Japan^bLaboratory of Forest Ecology, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan^cForestry & Forest Products Research Institute, Tsukuba, Ibaraki 305-8687, Japan

ARTICLE INFO

Article history:

Received 13 December 2011

Received in revised form

17 April 2012

Accepted 15 May 2012

Available online 13 September 2012

Keywords:

Fungi

Lignin

Bleached litter

Xylariaceae

ABSTRACT

To test the hypothesis that xylariaceous endophytes were ubiquitous on live and dead leaves of various tree species in the field, xylariaceous fungi were isolated from live leaves and bleached and nonbleached portions of dead leaves of a total of 94 tree species in a cool temperate forest in Japan. The biodiversity of xylariaceous endophytes was evaluated as the richness of operational taxonomic units (OTUs) determined by phylogenetic analysis of the nucleotide sequence of the D1/D2 region of the LSU rDNA of fungal isolates. A total of 326 isolates of xylariaceous fungi were isolated from live and dead leaves and classified into 15 OTUs. The three major OTUs, *Xylaria* sp.1, *Nemania* sp., and *Biscogniauxia* sp., accounted for 94% (308 isolates) of the total number of isolates, and were isolated from various live and dead leaves. *Xylaria* sp.1 was frequently encountered on bleached portions (which were produced due to the selective decomposition of lignin) of dead leaves of broad-leaved deciduous tree species. The results suggest that xylariaceous endophytes did not show host specificity and had a saprobic phase on dead leaves in their life cycles and that *Xylaria* sp.1 was capable of decomposing lignin in the field conditions.

© 2012 The Mycological Society of Japan. Published by Elsevier B.V. All rights reserved.

1. Introduction

Endophytic fungi are defined as those that can colonize internal plant tissues at some time in their life without causing apparent harm to their host (Sieber 2007). Endophytic fungi on leaves of forest trees play ecological roles as presumed mutualists (Saikkonen 2007), latent pathogens (Sieber 2007), and saprobic decomposers after leaf death (Osono 2006; Promputtha et al. 2007). Previous studies have shown that a few groups of endophytic fungi in the Rhytismataceae and Xylariaceae in Ascomycota take part in the decomposition of lignin (Osono 2002; Koide et al. 2005; Osono and Hirose 2010). As lignin is a major structural component

often limiting decomposition (Hirobe et al. 2004; Osono and Takeda 2005), these ligninolytic endophytes are of particular importance in terms of their roles in carbon turnover and nutrient cycling in forest ecosystems and deserve further studies on their ecology and functioning.

Osono and Hirose (2009) reviewed the ecology of endophytic fungi associated with leaf litter decomposition and recognized two groups of ligninolytic endophytes. The first is Rhytismataceae endophytes, which are relatively host-specific, usually colonize dead leaves for less than one year, and cause lignin decomposition in the initial stage of decomposition. The second is xylariaceous endophytes, which appear to have low host specificity and are found on

* Corresponding author. Tel.: +81 77 549 8252; fax: +81 77 549 8201.

E-mail address: tosono@ecology.kyoto-u.ac.jp (T. Osono).

leaves of various tree species (Whalley 1985, 1996; Petrini and Petrini 1985; Petrini et al. 1995). Xylariaceous endophytes are primarily saprobic and persist until the late stages of decomposition (Osono 2006). Osono and Takeda (2001) demonstrated that an endophyte, *Xylaria* sp., was frequently isolated from bleached portions on dead leaves of Japanese beech (*Fagus crenata*), which were produced due to the selective decomposition of lignin by the fungal colonizer. It is unclear, however, whether the ligninolytic activity of xylariaceous endophytes occur on leaf litter of other tree species with different leaf traits. Further studies are needed regarding the biodiversity, host range, and functioning of xylariaceous endophytes associated with leaf litter decomposition and, particularly, with lignin decomposition.

The purpose of the present study was to evaluate the diversity and ubiquity of xylariaceous endophytes in live and dead leaves of trees in a cool temperate forest. Thus, we isolated xylariaceous endophytes from live and dead leaves for a total of 94 tree species with one of four types of leaf traits (79 broad-leaved deciduous, 8 broad-leaved evergreen, 2 coniferous deciduous, and 5 coniferous evergreen species). The biodiversity of xylariaceous endophytes was evaluated as the richness of operational taxonomic units (OTUs) examined with phylogenetic analysis of the nucleotide sequence of the D1/D2 region of the LSU rDNA of fungal isolates. Fungi in Xylariaceae have been extensively subjected to molecular phylogenetic analysis (i.e., Lee et al. 2000; Davis et al. 2003; Okane et al. 2008; Peláez et al. 2008; Guedegbe et al. 2009) and are suitable for molecular identification of fungal isolates obtained from live and dead leaves of different host trees.

2. Materials and methods

2.1. Study site

Leaf materials used for fungal isolation were collected in Ashiu Experimental Forest of Kyoto University (35°18'N, 135°43'E, 355–660 m a.s.l.), Kyoto Prefecture, central Japan. During the past 29 years, the mean annual temperature was 11.7 °C and mean monthly temperature ranged from 0.4 °C in January to 25.5 °C in August at the office of the Ashiu Experimental Forest at 355 m a.s.l. The mean annual precipitation during the past 29 years was 2353 mm. The study area is covered with snow from December to April. The Ashiu Experimental Forest is in a mountainous area, with natural stands of warm temperate forests dominated by evergreen oaks *Quercus salicina* and *Q. acuta*, below approximately 600 m a.s.l., and natural stands of cool temperate forests dominated by a deciduous beech, *Fagus crenata*, and a deciduous oak, *Q. crispula*, above the warm temperate region. The area is thus an ecotone of two climatic regions and hence has high richness of plant species, including 243 tree species recorded in the Ashiu Experimental Forest.

2.2. Sample collection

Live and dead leaves of a total of 94 tree species in 38 plant families were collected in the study site during the growing season from May to November in 2008 (Table 1). Broad-leaved

deciduous tree species accounted for 84% (79 species) of the 94 species examined. Live, healthy-looking leaves of 74 tree species were collected in May, August, and November, mostly in August. On each sampling occasion, a total of 10 live, healthy-looking leaves were harvested for each tree species from two randomly chosen trees, two branches per individual tree, at an approximate height of 3–4 m. Two types of dead leaves were collected: those bearing bleached portions on the surfaces and those that were not bleached. The presence of bleached portions is associated with fungal colonization within leaf tissues and decomposition of lignin (Osono 2007). In the present study, the bleached portions were observed on dead leaves of 15 tree species, including 12 deciduous broad-leaved, 2 evergreen broad-leaved, and one evergreen coniferous tree species (Table 1). The bleached dead leaves were collected in May and July. Dead leaves without obvious bleached portions (denoted as nonbleached dead leaves) were collected for 63 tree species in May, June, and November. Bleached and nonbleached dead leaves were collected from the forest floor for each tree species on each sampling occasion. Sampling of a total of 1840 leaves was carried out during the study period. The leaves were placed in paper bags and taken to the laboratory. The leaves were processed within 24 h after the collection. One leaf disk was punched out from the central part of each sample leaf, avoiding the primary vein, with a sterile cork borer (5.5 mm in diameter). A total of 10 leaf disks were used for each tree species, each leaf type, and on each sampling occasion, making a total of 1840 disks for the isolation of xylariaceous fungi.

2.3. Fungal isolation

A surface disinfection method (Kinkel and Andrews 1988) was used to isolate xylariaceous fungi. The leaf disks were submerged in 70% ethanol (v/v) for 1 min to wet the surface, then surface-disinfected for 30 sec in a solution of 15% hydrogen peroxide, and then submerged again for 1 min in 70% ethanol. The disks were rinsed with sterile, distilled water, transferred to sterile filter paper in Petri dishes (9 cm in diameter), and dried for 24 h to suppress vigorous bacterial growth after plating (Widden and Parkinson 1973). The disks were placed in 9-cm Petri dishes containing lignocellulose agar (LCA) modified by Miura and Kudo (1970), two disks per plate. LCA contains glucose 0.1%, KH₂PO₄ 0.1%, MgSO₄·7H₂O 0.02%, KCl 0.02%, NaNO₃ 0.2%, yeast extract 0.02%, and agar 1.3% (w/v). Note that the modified LCA of Miura and Kudo (1970) does not contain lignin or other recalcitrant compounds. The modified LCA was used because its low glucose content suppresses the overgrowth of fast-growing fungal species (Osono and Takeda 1999). Plates were incubated at 20 °C in the dark and observed at 1, 4, and 8 weeks after surface disinfection. Putative xylariaceous fungi that produced on the plates conidia and conidiophores of anamorphic Xylariaceae, such as *Xylocoremium*, *Geniculosporium*, and *Nodulisporium*, and/or dark pseudosclerotinial plates in submerged hyphae were subcultured on fresh LCA to establish pure cultures.

2.4. Determination of OTUs

The pure cultures obtained were identified by molecular analysis. When fungal structures such as spores and sporocarps were produced on the medium, their morphological

Table 1 – Tree species examined in the present study, month of collection in 2008, and the number of isolates of xyleriaceous fungi on live leaves and bleached and nonbleached portions of dead leaves. Tree names are arranged in alphabetical order of abbreviated codes.

Tree species	Family ^a	Abbr.	Live leaves		Bleached portions of dead leaves		Nonbleached portions of dead leaves	
			Month ^b	Number of isolates	Month	Number of isolates	Month	Number of isolates
Broad-leaved, deciduous								
<i>Actinidia arguta</i>	At	Aa	A	0	–	–	–	–
<i>Acer carpinifolium</i>	Ac	Ac	A	2	–	–	N	0
<i>Acer nikoense</i>	Ac	Ae	A	2	–	–	–	–
<i>Alnus firma</i>	Be	Af	M, A, N	0, 7, 0	–	–	–	–
<i>Actinidia polygama</i>	At	Ag	A	2	–	–	N	1
<i>Acer japonicum</i>	Ac	Aj	A	4	–	–	N	4
<i>Acer micranthum</i>	Ac	Ak	A	1	–	–	–	–
<i>Aralia elata</i>	Ar	Al	A	1	–	–	–	–
<i>Acer mono</i>	Ac	Am	A	2	M	3	N	0
<i>Acer nipponicum</i>	Ac	An	A	3	–	–	–	–
<i>Acanthopanax sciadophylloides</i>	Ar	Ap	A	0	–	–	N	1
<i>Akebia quinata</i>	Lr	Aq	A	1	–	–	–	–
<i>Acer rufinerve</i>	Ac	Ar	A	3	M	4	N	1
<i>Acer sieboldianum</i>	Ac	As	A	4	M	1	N	0
<i>Aesculus turbinata</i>	Hi	At	A	1	M	2	N	0
<i>Acer amoenum</i> var. <i>matsumurae</i>	Ac	Ay	–	–	–	–	N	1
<i>Benthamidia cousa</i>	Co	Bc	A	3	–	–	N	3
<i>Betula grossa</i>	Be	Bg	M, A, N	1, 2, 7	M	2	N	1
<i>Boehmeria spicata</i>	Ur	Bs	A	1	–	–	–	–
<i>Clethra barvinervis</i>	Er	Cb	A	3	–	–	M, N	0, 0
<i>Castanea crenata</i>	Fa	Cc	A	1	M	1	N	0
<i>Corylus sieboldiana</i>	Be	Ch	A	0	–	–	–	–
<i>Carpinus tschonoskii</i>	Be	Ci	A	8	–	–	N	2
<i>Cercidiphyllum japonicus</i>	Cr	Cj	A	6	–	–	M, N	0, 1
<i>Carpinus japonica</i>	Be	Ck	A	8	–	–	N	4
<i>Carpinus laxiflora</i>	Be	Cl	M, A, N	0, 1, 4	M	3	N	0
<i>Carpinus cordata</i>	Be	Cs	A	4	–	–	N	0
<i>Clerodendrum trichotomum</i>	Ve	Ct	A	0	–	–	–	–
<i>Cladrastis sikokiana</i>	Fb	Cy	A	3	–	–	N	0
<i>Deutzia crenata</i>	Sx	Dc	A	3	–	–	N	0
<i>Eunomyces alatus</i> f. <i>subtriflorus</i>	Ce	Ea	A	3	–	–	N	1
<i>Evodiopanax innovans</i>	Ar	El	A	1	–	–	N	0
<i>Elliottia paniculata</i>	Er	Ep	–	–	–	–	N	0
<i>Fagus crenata</i>	Fa	Fc	A	3	M	2	N	0
<i>Fraxinus sieboldiana</i>	Ol	Fs	A	5	–	–	–	–
<i>Hydrangea hirta</i>	Sx	Hh	A	2	–	–	N	3
<i>Hamamelis japonicus</i> var. <i>obtusata</i>	Ha	Hj	A	4	–	–	M, N	0, 3
<i>Hydrangea petiolaris</i>	Sx	Hp	A	1	–	–	M, N	1, 1
<i>Hovenia tomentella</i>	Rh	Ht	–	–	–	–	N	0
<i>Hydrangea paniculata</i>	Sx	Hy	A	4	–	–	–	–
<i>Ilex macropoda</i>	Aq	Im	A	3	–	–	N	3
<i>Juglans mandshurica</i> var. <i>sachaliensis</i>	Ju	Jm	A	3	–	–	–	–
<i>Kalopanax pictus</i>	Ar	Kp	–	–	–	–	M, N	2, 0
<i>Lindera erythrocarpa</i>	La	Le	A	3	–	–	Jun, N	0, 0
<i>Lyonia ovalifolia</i>	Er	Lo	A	0	M	3	N	2
<i>Lindera triloba</i>	La	Lt	A	2	–	–	N	0
<i>Lindera umbellata</i>	La	Lu	A	2	–	–	N	0
<i>Mallotus japonicus</i>	Eu	Mj	A	1	–	–	N	6
<i>Meliosma myriantha</i>	Sa	Mm	A	0	–	–	N	5
<i>Magnolia obovata</i>	Ma	Mo	A	3	M	3	N	0
<i>Magnolia salicifolia</i>	Ma	Ms	A	2	–	–	M, N	1, 0
<i>Malus tschonoskii</i>	Ro	Mt	A	2	–	–	N	0

Table 1 – (continued)

Tree species	Family ^a	Abbr.	Live leaves		Bleached portions of dead leaves		Nonbleached portions of dead leaves	
			Month ^b	Number of isolates	Month	Number of isolates	Month	Number of isolates
<i>Prunus grayana</i>	Ro	Pg	A	6	–	–	N	1
<i>Pterostyrax hispida</i>	St	Ph	A	0	–	–	N	0
<i>Pterocarya rhoifolia</i>	Ju	Pr	A	5	–	–	M, N	0, 0
<i>Populus sieboldii</i>	Sl	Ps	A	5	–	–	–	–
<i>Paulownia tomentosa</i>	Pa	Pt	A	1	–	–	–	–
<i>Prunus jamasakura</i>	Ro	Py	A	5	–	–	N	4
<i>Quercus crispula</i>	Fa	Qc	A	4	M	1	N	1
<i>Quercus serrata</i>	Fa	Qs	A	0	M	2	N	2
<i>Rhus javanica</i> var. <i>roxburghii</i>	An	Rj	A	2	–	–	–	–
<i>Rhus trichocarpa</i>	An	Rt	A	1	–	–	–	–
<i>Sorbus alnifolia</i>	Ro	Sa	A	1	–	–	M, N	1, 0
<i>Swida controversa</i>	Co	Sc	A	1	–	–	N	1
<i>Schizophragma hydrangeoides</i>	Sx	Sh	A	2	–	–	–	–
<i>Styrax japonica</i>	St	Sj	A	5	–	–	–	–
<i>Sorbus commixta</i>	Ro	Sn	A	1	–	–	N	2
<i>Styrax obassia</i>	St	So	A	1	–	–	N	0
<i>Stachyurus praecox</i>	Sp	Sp	A	3	–	–	N	6
<i>Symplocos chinensis</i>	Sy	Ss	A	3	–	–	N	7
<i>Symplocos coreana</i>	Sy	St	A	2	–	–	–	–
<i>Ulmus parvifolia</i>	Ul	Up	A	3	–	–	N	1
<i>Viburnum furcatum</i>	Ca	Vf	A	3	–	–	–	–
<i>Viburnum plicatum</i> var. <i>tomentosum</i>	Ca	Vp	A	3	–	–	M, N	0, 0
<i>Viburnum wrightii</i>	Ca	Vw	A	0	–	–	–	–
<i>Wisteria floribunda</i>	Fb	Wf	A	0	–	–	–	–
<i>Weigela hortensis</i>	Ca	Wh	A	4	–	–	N	1
<i>Zanthoxylum piperitum</i>	Ru	Zp	A	0	–	–	–	–
<i>Zelkova serrata</i>	Ul	Zs	A	0	–	–	N	0
Broad-leaved, evergreen								
<i>Daphniphyllum macropodium</i> var. <i>humile</i>	Da	Dm	M	4	–	–	M	0
<i>Eurya japonica</i>	Th	Ej	A	2	–	–	–	–
<i>Ilex pedunculosa</i>	Aq	Ip	M	2	–	–	–	–
<i>Ilex sugerokii</i>	Aq	Is	A	2	–	–	–	–
<i>Pieris japonica</i>	Er	Pj	M	0	–	–	M	0
<i>Quercus acuta</i>	Fa	Qa	–	–	Jul	0	–	–
<i>Quercus salicina</i>	Fa	Ql	A	3	Jul	0	–	–
<i>Trochodendron aralioides</i>	Tr	Ta	M	2	–	–	M	2
Coniferous, deciduous								
<i>Larix gmelinii</i>	Pi	Lg	A	0	–	–	M, N	1, 0
<i>Metasequoia glyptostroboides</i>	Cu	Mg	A	2	–	–	N	0
Coniferous, evergreen								
<i>Abies firma</i>	Pi	Ai	–	–	Jul	0	–	–
<i>Chamaecyparis obtusa</i>	Cu	Co	A	1	–	–	–	–
<i>Cryptomeria japonica</i>	Cu	Cr	A	1	–	–	M, N	1, 3
<i>Picea abies</i>	Pi	Pa	A	1	–	–	M	1
<i>Pinus densiflora</i>	Pi	Pd	A	3	–	–	M	1
Number of tree species examined		94		88		15		63
Number of leaves examined		1840		940		150		750
Number of isolates of xylariaceous fungi		326		216		27		83
Number of tree species from which xylariaceous fungi were isolated		82		74		12		36
(% total number of tree species)		(87%)		(84%)		(80%)		(57%)

a Tree family: Ac, Aceraceae; An, Anacardiaceae; Aq, Aquifoliaceae; Ar, Araliaceae; At, Actinidiaceae; Be, Betulaceae; Ca, Caprifoliaceae; Ce, Celastraceae; Co, Cornaceae; Cr, Cercidiphyllaceae; Cu, Cupressaceae; Da, Daphniphyllaceae; Er, Ericaceae; Eu, Euphorbiaceae; Fa, Fagaceae; Fb, Fabaceae; Ha, Hamamelidaceae; Hi, Hippocastanaceae; Ju, Juglandaceae; La, Lauraceae; Lr, Lardizabalaceae; Ol, Oleaceae; Pa, Paulowniaceae; Pi, Pinaceae; Rh, Rhamnaceae; Ro, Rosaceae; Ru, Rutaceae; Sa, Sabiaceae; Sl, Salicaceae; Sx, Saxifragaceae; Sp, Stachyuraceae; St, Styracaceae; Sy, Symplocaceae; Th, Theaceae; Tr, Trochodendraceae; Ul, Ulmaceae; Ur, Urticaceae; Ve, Verbenaceae.

b Month: M, May; Jun, June; Jul, July; A, August; N, November.

characteristics were observed with a Nikon Optiphot microscope (Nikon Inc., Tokyo, Japan). For molecular analysis, a small amount of mycelial tips from each culture were picked, crushed in 24 µl of distilled water in a tube, microwaved for 12 sec, and used as templates for PCR. A reaction mixture (50 µl) containing 25 µl Qiagen GoTaq premix (Qiagen, Ontario, Canada) and 10 pmol of each primer and distilled water was added to the templates. The oligonucleotide primer-pair NL1 and NL4 (O'Donnell 1993) were used for PCR of ribosomal DNA large subunit D1/D2 region. The reactions were initiated with 4 min of denaturation at 95 °C, followed by 40 cycles of two-step PCR, consisting of 20 s at 94 °C and 60 s at 56 °C with a final extension for 10 min at 72 °C on a GeneAmp 9700 thermal cycler (Perkin–Elmer Applied Biosystems, California, USA). Amplification products were purified using a QIAquick PCR Purification Kit (Qiagen, Ontario, Canada), and sequenced with a Big Dye Terminator Cycle Sequencing FS Ready Reaction kit ver. 3.1 and an ABI PRISM 3100 genetic analyzer (Perkin–Elmer Applied Biosystems, California, USA). Both strands of a fragment were sequenced. Sequence data sets were manually truncated both ends and edited using the program BioEdit sequence editor version 5.09 (Hall 1999). Homology searches were performed using each obtained sequences data on a BLAST program at the National Center for Biotechnology Information (NCBI). Neighbor joining trees were also constructed using MEGA version 5 (Tamura et al. 2011) with related sequences from NCBI database. Isolates with more than 99% homology of sequence and within the same cluster were treated as OTUs with tentative codes for data analysis. In the case that obtained sequences contained polymorphic sites, they are treated as the same OTUs with close relatives.

2.5. Statistical analysis

To assess the affinity of major fungal OTUs to leaf traits of host trees, Fisher's exact probability test was performed to examine the differences in the number of tree species from which the major OTUs were isolated between broad-leaved and coniferous trees and between deciduous and evergreen trees.

3. Results

3.1. Operational taxonomic units of xylariaceous fungi

A total of 326 isolates of xylariaceous fungi were isolated from live and dead leaves of 82 (87%) of the 94 tree species examined (Table 1). Xylariaceous fungi were isolated from live leaves of 74 (84%) of 88 tree species, from bleached dead leaves of 12 (80%) of 15 tree species, and from nonbleached dead leaves of 36 (57%) of 63 tree species (Table 1).

The fungal isolates were classified into 15 OTUs (Table 2). *Xylaria* sp.1 was the most dominant OTU (135 isolates), followed by *Nemania* sp. (123 isolates) and *Biscogniauxia* sp. (50 isolates). These three OTUs (308 isolates) accounted for 94% of the total number of isolates (326 isolates). The other 12 OTUs

Table 2 — BLAST search result of 15 operational taxonomic units (OTUs) of xylariaceous fungi on live leaves and bleached and nonbleached portions of dead leaves, and the number of tree species from which the 15 OTUs were isolated.

OTU	Accession number	BLAST search result	Identity (%)	Number of isolates				Number of tree species			
				Live [216]		Dead leaves		Live [88]		Dead leaves	
				Bleached [27]	Nonbleached [83]	Bleached [15]	Nonbleached [63]	Bleached [15]	Nonbleached [63]		
<i>Xylaria</i> sp.1	AB686646	Xylariales cf. JP14-3, Q906954	99	68 (31%)	25 (93%)	42 (51%)	43 (49%)	12 (80%)	22 (35%)		
<i>Nemania</i> sp.	AB669031	Fungal sp. mh337.6, GU552551	100	89 (41%)	0	34 (41%)	51 (58%)	0	26 (41%)		
<i>Biscogniauxia</i> sp.	AB686647	Xylariaceae sp., GU048581	98	44 (20%)	1 (4%)	5 (6%)	24 (27%)	1 (7%)	5 (8%)		
<i>Annulohypoxylon</i> sp.	AB669044	<i>Annulohypoxylon moriforme</i> , DQ840058	99	3 (1.4%)	0	1 (1.2%)	2 (2%)	0	1 (2%)		
<i>Daldinia</i> sp.	AB669047	<i>Daldinia chlidiaae</i> , EF562505	99	2 (0.9%)	0	0	2 (2%)	0	0		
<i>Xylaria</i> sp.2	AB686648	Fungal sp. mh1111.61, GU552542	99	2 (0.9%)	0	0	2 (2%)	0	0		
<i>Rosellinia</i> sp.	AB686649	<i>Rosellinia corticium</i> , DQ840078	99	2 (0.9%)	0	0	2 (2%)	0	0		
<i>Xylaria hypoxylon</i>	AB686650	<i>Xylaria hypoxylon</i> , NG_027599	97	1 (0.5%)	0	0	1 (1%)	0	0		
<i>Nodulisporium</i> sp.	AB686651	<i>Annulohypoxylon moriforme</i> , DQ840057	99	0	1 (4%)	0	0	1 (7%)	0		
<i>Hypoxylon</i> cf. <i>fragiforme</i>	AB686652	<i>Hypoxylon fragiforme</i> , AY083829	99	0	0	1 (1.2%)	0	0	1 (2%)		
<i>Xylaria</i> sp.4	AB686653	Fungal endophyte, EU687185	97	1 (0.5%)	0	0	1 (1%)	0	0		
<i>Xylaria</i> sp.3	AB686654	Xylariaceae sp., AB376751	98	1 (0.5%)	0	0	1 (1%)	0	0		
<i>Xylaria</i> sp.5	AB686655	Xylariales cf. JP14-3, GQ906954	94	1 (0.5%)	0	0	1 (1%)	0	0		
<i>Xylaria</i> sp.6	AB669070	<i>Anthostomella leucospermi</i> , EU552100	99	1 (0.5%)	0	0	1 (1%)	0	0		
<i>Xylariaceae</i> sp.	AB669071	Xylariaceae sp., GU048581	95	1 (0.5%)	0	0	1 (1%)	0	0		

Note: Total numbers of isolates or tree species examined are shown in square brackets, and the percentages relative to the total number of tree species are shown in parentheses.

were isolated only infrequently, with the number of isolates ranging from 1 to 4.

The number of OTUs isolated from live leaves was 13, and those from bleached and nonbleached dead leaves were 3 and 5, respectively (Table 2). *Nemania* sp. was the most dominant OTU on live leaves, followed by *Xylaria* sp.1 and *Biscogniauxia* sp. *Xylaria* sp.1 accounted for 93% of the total number of isolates from bleached dead leaves. *Xylaria* sp.1 was the most dominant OTU on nonbleached dead leaves, followed by *Nemania* sp. and *Biscogniauxia* sp.

Xylaria sp.1, *Nemania* sp., and *Biscogniauxia* sp. were isolated from live leaves of 43 (49%), 51 (58%), and 24 (27%), respectively, of 88 tree species examined (Table 2). *Xylaria* sp.1 was isolated from bleached dead leaves of 12 tree species (Table 2). *Xylaria* sp.1, *Nemania* sp., and *Biscogniauxia* sp. were isolated from nonbleached dead leaves of 22 (35%), 26 (41%), and 5 (8%) of 63 tree species examined, respectively (Table 2). The number of tree species from which *Xylaria* sp.1, *Nemania* sp., and *Biscogniauxia* sp. were isolated from both of live and dead leaves (bleached or nonbleached) was 20, 16, and 1 species, respectively.

3.2. Patterns of occurrence of major OTUs

The number of tree species from which the three major OTUs were isolated was summarized in Table 3 with respect to four types of leaf traits (i.e., broad-leaved deciduous, broad-leaved evergreen, coniferous deciduous, and coniferous evergreen). When live leaves were considered, the number of tree species from which *Xylaria* sp.1, *Nemania* sp., and *Biscogniauxia* sp. were isolated was not significantly different between broad-leaved and coniferous trees ($P = 0.25$, $P = 0.31$, and $P = 0.17$, respectively) or between deciduous and evergreen trees ($P = 0.17$, $P = 0.24$, and $P = 0.15$, respectively) (Table 3).

Xylaria sp.1 was isolated from bleached dead leaves of all of the 12 broad-leaved deciduous tree species examined, but not on those of broad-leaved or coniferous evergreen tree species (Table 3). *Nemania* sp. was not isolated from bleached dead

leaves (Table 3). *Biscogniauxia* sp. was isolated from bleached dead leaves of one broad-leaved deciduous tree species (Table 3).

When nonbleached dead leaves were considered, the number of tree species from which *Xylaria* sp.1 and *Nemania* sp. were isolated was not significantly different between broad-leaved and coniferous trees ($P = 0.32$ and $P = 0.24$, respectively) or between deciduous and evergreen trees ($P = 0.34$ and $P = 0.17$, respectively) (Table 3). *Biscogniauxia* sp. was isolated from nonbleached dead leaves of 2 out of 58 broad-leaved tree species, which were significantly ($P = 0.002$) lower than in coniferous trees (three out of five tree species) (Table 3). The number of tree species of which *Biscogniauxia* sp. was isolated from nonbleached dead leaves was not significantly different between deciduous and evergreen trees ($P = 0.06$).

4. Discussion

In previous studies at the present study site, xylariaceous fungi were isolated from live and dead leaves of two major tree species, *Fagus crenata* and *Swida controversa* (Osono 2002; Osono et al. 2004). In the present study, 15 OTUs of xylariaceous fungi were found from live and dead leaves of 94 tree species, indicating that they are major components of endophytic and litter-inhabiting fungi in the cool temperate forest. The community structure of the fungal OTUs was highly skewed, with the top three OTUs accounting for 94% of the total number of isolates (Table 2). In a similar study of endophytic Xylariaceae from Thailand, Okane et al. (2008) isolated from live leaves of 25 tree species a total of 273 isolates that were assigned to 25 OTUs according to their 28S rDNA D1/D2 sequence. The top three OTUs with respect to the number of fungal isolates accounted for 31% of the total number of isolates in the study of Okane et al. (2008). Thus, the diversity of xylariaceous endophytes was lower in the cool temperate forest in the present study than in the tropical forest of Okane et al. (2008) in terms of the dominance of a few major OTUs and the lower prevalence of rare OTUs.

Table 3 – Number of tree species from which the 15 OTUs of xylariaceous fungi and three major OTUs were isolated, as related to the life form of tree species.

	Total number of tree species examined	Xylariaceae, 15 OTUs	<i>Xylaria</i> sp.1	<i>Nemania</i> sp.	<i>Biscogniauxia</i> sp.
Live leaves					
Broad-leaved, deciduous	75	63 (84%)	38 (51%)	43 (57%)	21 (28%)
Broad-leaved, evergreen	7	6 (86%)	3 (43%)	4 (57%)	1 (14%)
Coniferous, deciduous	2	1 (50%)	1 (50%)	1 (50%)	0 (0%)
Coniferous, evergreen	4	4 (100%)	1 (25%)	3 (75%)	0 (0%)
Bleached portions of dead leaves					
Broad-leaved, deciduous	12	12 (100%)	12 (100%)	0 (0%)	1 (8%)
Broad-leaved, evergreen	2	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Coniferous, evergreen	1	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Nonbleached portions of dead leaves					
Broad-leaved, deciduous	55	31 (56%)	20 (36%)	25 (45%)	2 (4%)
Broad-leaved, evergreen	3	1 (33%)	1 (33%)	0 (0%)	0 (0%)
Coniferous, deciduous	2	1 (50%)	0 (0%)	0 (0%)	1 (50%)
Coniferous, evergreen	3	3 (100%)	1 (33%)	1 (33%)	2 (67%)

Note: No bleached portions of dead leaves were examined for coniferous deciduous tree species.

Xylaria sp.1 and *Nemania* sp. occurred on live and dead leaves of multiple tree species (Table 3), regardless of leaf traits (i.e. broad-leaved vs coniferous, deciduous vs evergreen), suggesting the low host specificity, which is consistent with previous studies of xylariaceous endophytes (Petrini et al. 1995; Cannon and Simmons 2002; Murali et al. 2007; Okane et al. 2008). Previous studies have shown that *Xylaria* sp.1 was also isolated from live and dead twigs (Fukasawa et al. 2009) and cupules (Fukasawa et al. in press) of *F. crenata*, indicating its low tissue specificity. *Biscogniauxia* sp. was isolated from nonbleached dead leaves more frequently (in terms of the number of tree species isolated with respect to the total number of tree species examined) for coniferous than for broad-leaved tree species (Table 3), but the number of coniferous tree species examined was too low (i.e. five species) to be conclusive with the affinity of this OTU to the dead coniferous leaves.

Relating the endophytic fungal OTUs from live and dead leaves to their fruiting bodies is needed to evaluate their ecology and life cycles. Unfortunately, however, the teleomorphic states of the major OTUs in the present study have not yet been collected at the study site, making it difficult to evaluate the ecology and life cycle of the leaf-associated xylariaceous fungi in detail. Only a few rare OTUs have been phylogenetically related to teleomorphic states fruiting on woody tissues (i.e., *Xylaria hypoxylon* and *Hypoxylon fragiforme* in Table 2). Thus, further efforts are needed to search for fruiting bodies to identify the OTUs and to clarify their ecology and host- and tissue-specificity at the study site. It might also be important to take into consideration the possibility that the endophytic life stage of xylariaceous fungi is 'a dead-end' of the life cycle as it rarely ends with sexual reproduction on the leaf. Alternatively, some xylariaceous endophytes with *Geniculosporium* and *Nodulisporium* anamorphs can establish from conidia and grow and reproduce endophytically as anamorphic fungi (Rogers 1985).

The three major OTUs were isolated from not only live leaves but also bleached and nonbleached dead leaves of broad-leaved deciduous tree species (Table 2), indicating that these xylariaceous endophytes have a saprobic phase in their life cycles. The isolation of *Xylaria* sp.1 and *Nemania* sp. from both live and dead leaves of the same tree species suggested that these OTUs could persist in dead leaves from live leaves. *Xylaria* sp.1 was isolated from both bleached and nonbleached dead leaves, whereas *Nemania* sp. and *Biscogniauxia* sp. were mostly isolated from nonbleached dead leaves (Table 2). Because the lignin content is lower in bleached than in nonbleached portions (Osono 2007), *Xylaria* sp.1 is probably capable of decomposing lignin more actively than the other two OTUs. This is consistent with a pure culture test showing that *Xylaria* sp.1 contained isolates that decomposed lignin in *F. crenata* leaves more actively than *Nemania* sp. (as *Geniculosporium* sp., Osono and Takeda 2002). However, we cannot exclude a possibility that *Xylaria* sp.1 preferred bleached to nonbleached portions as substrata for colonization.

Xylaria sp.1 was isolated from bleached dead leaves of deciduous broad-leaved trees but not from those of evergreen broad-leaved or coniferous trees, despite its occurrence on live and nonbleached dead leaves of these evergreen trees (Table 3). However, the number of evergreen tree species

examined for bleached dead leaves in the present study was too low to determine whether *Xylaria* sp.1 was truly absent from bleached dead leaves of evergreen trees. Further studies are needed to examine bleached dead leaves of evergreen trees for the occurrence of xylariaceous fungi and to explore possible mechanisms relating to the reduction of xylariaceous fungi in these leaves.

Acknowledgments

We thank members of the Ashiu Experimental Forest of Kyoto University for help with fieldwork; Dr. Norio Sahashi, Dr. Kunihiko Hata, and Dr. Izumi Okane for useful discussions; and Dr. Elizabeth Nakajima for critical reading of the manuscript. This work was supported by the Global Environmental Research Fund (RF-086) of the Ministry of the Environment, Japan, and by Global COE Program A06 of Kyoto University.

REFERENCES

- Cannon PF, Simmons CM, 2002. Diversity and host preference of leaf endophytic fungi in the Iwokrama Forest Reserve, Guyana. *Mycologia* 94: 210–220.
- Davis EC, Franklin JB, Shaw AJ, Vilgalys R, 2003. Endophytic *Xylaria* (Xylariaceae) among liverworts and angiosperms: phylogenetics, distribution, and symbiosis. *American Journal of Botany* 90: 1661–1667.
- Fukasawa Y, Osono T, Takeda H, 2009. Effects of attack of saprobic fungi on twig litter decomposition by endophytic fungi. *Ecological Research* 24: 1067–1073.
- Fukasawa Y, Tateno O, Hagiwara Y, Hirose D, Osono T, 2012. Fungal succession and decomposition of beech cupule litter. *Ecological Research* 27: 735–743.
- Guedegbe HJ, Miambi E, Pando A, Hounnandan P, Rouland-Lefevre C, 2009. Molecular diversity and host specificity of termite-associated *Xylaria*. *Mycologia* 101: 686–691.
- Hall TA, 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hirobe M, Sabang J, Bhatta BK, Takeda H, 2004. Leaf-litter decomposition of 15 tree species in a lowland tropical rain forest in Sarawak: decomposition rates and initial litter chemistry. *Journal of Forest Research* 9: 341–346.
- Kinkel LL, Andrews JH, 1988. Disinfection of living leaves by hydrogen peroxide. *Transactions of the British Mycological Society* 91: 523–528.
- Koide K, Osono T, Takeda H, 2005. Colonization and lignin decomposition of *Camellia japonica* leaf litter by endophytic fungi. *Mycoscience* 46: 280–286.
- Lee JS, Ko KS, Jung HS, 2000. Phylogenetic analysis of *Xylaria* based on nuclear ribosomal ITS1-5.8S-ITS2 sequences. *FEMS Microbiology Letters* 187: 89–93.
- Miura K, Kudo M, 1970. An agar-medium for aquatic hyphomycetes (in Japanese with English abstract). *Transactions of the Mycological Society of Japan* 11: 116–118.
- Murali TS, Suryanarayanan TS, Venkatesan G, 2007. Fungal endophyte communities in two tropical forests of southern India: diversity and host affiliation. *Mycological Progress* 6: 191–199.
- O'Donnell K, 1993. *Fusarium* and its near relatives. In: Reynolds DR, Taylor JW (eds), *The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*. CAB International, Wallingford, UK, pp 225–233.

- Okane I, Srikitikulchai P, Toyama K, Læssøe T, Sivichai S, Hywel-Jones N, Nakagiri A, Potacharoen W, Suzuki K, 2008. Study of endophytic Xylariaceae in Thailand: diversity and taxonomy inferred from rDNA sequence analyses with saprobes forming fruit bodies in the field. *Mycoscience* 49: 359–372.
- Osono T, 2002. Phyllosphere fungi on leaf litter of *Fagus crenata*: occurrence, colonization, and succession. *Canadian Journal of Botany* 80: 460–469.
- Osono T, 2006. Role of phyllosphere fungi of forest trees in the development of decomposer fungal communities and decomposition processes of leaf litter. *Canadian Journal of Microbiology* 52: 701–716.
- Osono T, 2007. Ecology of ligninolytic fungi associated with leaf litter decomposition. *Ecological Research* 22: 955–974.
- Osono T, Takeda H, 1999. A methodological survey on incubation of fungi on leaf litter of *Fagus crenata* (in Japanese with English abstract). *Applied Forest Science Kansai* 8: 103–108.
- Osono T, Takeda H, 2001. Effects of organic chemical quality and mineral nitrogen addition on lignin and holocellulose decomposition of beech leaf litter by *Xylaria* sp. *European Journal of Soil Biology* 37: 17–23.
- Osono T, Takeda H, 2002. Comparison of litter decomposing ability among diverse fungi in a cool temperate deciduous forest in Japan. *Mycologia* 94: 421–427.
- Osono T, Takeda H, 2005. Decomposition of lignin, holocellulose, polyphenol and soluble carbohydrate in leaf litter of 14 tree species in a cool temperate forest. *Ecological Research* 20: 41–49.
- Osono T, Hirose D, 2009. Ecology of endophytic fungi associated with leaf litter decomposition. In: Rai M, Bridge P (eds), *Applied Mycology*. CAB International, Oxon, UK, pp 92–109.
- Osono T, Hirose D, 2010. Colonization and lignin decomposition of pine needle litter by *Lophodermium pinastri*. *Forest Pathology* 41: 156–162.
- Osono T, Bhatta BK, Takeda H, 2004. Phyllosphere fungi on living and decomposing leaves of giant dogwood. *Mycoscience* 45: 35–41.
- Peláez F, González V, Platas G, Sánchez-Ballesteros J, Rubio V, 2008. Molecular phylogenetic studies within the Xylariaceae based on ribosomal DNA sequences. *Fungal Diversity* 31: 111–134.
- Petrini L, Petrini O, 1985. Xylariaceous fungi as endophytes. *Sydowia* 38: 216–234.
- Petrini O, Petrini LE, Rodrigues KF, 1995. Xylariaceous endophytes: an exercise in biodiversity. *Fitopatologia Brasileira* 20: 531–539.
- Promptutha I, Lumyong S, Dhanasekaren V, McKenzie EHC, Hyde KD, Jeewon R, 2007. A phylogenetic evaluation of whether endophytes become saprotrophs at host senescence. *Microbial Ecology* 53: 579–590.
- Rogers JD, 1985. Anamorphs of *Xylaria*: taxonomic considerations. *Sydowia* 38: 255–262.
- Saikkonen K, 2007. Forest structure and fungal endophytes. *Fungal Biology Reviews* 2: 67–74.
- Sieber TN, 2007. Endophytic fungi in forest trees: are they mutualists? *Fungal Biology Reviews* 2: 75–89.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S, 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739.
- Whalley AJS, 1985. The Xylariaceae: some ecological considerations. *Sydowia* 38: 369–382.
- Whalley AJS, 1996. The xylariaceous way of life. *Mycological Research* 100: 897–922.
- Widden P, Parkinson D, 1973. Fungi from Canadian coniferous forest soils. *Canadian Journal of Botany* 51: 2275–2290.