

Reply to Fulton and Barrett

Reply: Response to Letter to the Editor

Mehmet Sofuoglu*¹, Sonah Yoo¹, Kevin P Hill¹ and Marc Mooney²¹Department of Psychiatry, VA Connecticut Healthcare System, School of Medicine, Yale University, West Haven, CT, USA; ²Department of Psychiatry, Tobacco Use Research Center, University of Minnesota, Minneapolis, MN, USA

Neuropsychopharmacology (2008) 33, 2044; doi:10.1038/sj.npp.1301546

Sir

This letter is in response to the letter by Fulton and Barrett, which argued that our paper (Sofuoglu *et al*, 2007) does not support nicotine's reinforcing effects. Fulton and Barrett stated that since in our intravenous self-administration model we used an inactive placebo (saline) for nicotine, greater nicotine self-administration over saline could be due to smokers' 'expectations' of receiving nicotine rather than nicotine's reinforcing effects.

Although expectations of drug effects can confound clinical studies (Mooney *et al*, 2004), it is unlikely to explain our findings. First, in our study, nicotine self-administration and subjective nicotine effects were dependent on the nicotine dose (0.1, 0.4, and 0.7 mg). This would be unexpected if self-administration was based solely on 'expectation' of receiving nicotine. Second, if the 'expectation' of receiving nicotine was driving nicotine self-administration, then one would expect that smokers would choose nicotine over placebo for all routes of administration. This has not been the case. Nicotine products that lead to slower nicotine delivery, such as nicotine gum, are not reliably chosen over placebo gum, compared to the intravenous route, which provides faster nicotine delivery and is chosen over placebo (Hughes *et al*, 2000; Harvey *et al*, 2004). The most likely explanation of these findings is greater reinforcement with faster nicotine delivery (de Wit *et al*, 1992). Lastly, there is a wide range of preclinical literature on nicotine's reinforcing effects,

including those demonstrating the molecular and behavioral mechanisms of nicotine reinforcement (Picciotto and Corrigall, 2002; Le Foll and Goldberg, 2006). On the basis of this large body of evidence, it is hard to escape the conclusion that—like cocaine, heroin, and alcohol—nicotine is reinforcing.

REFERENCES

- de Wit H, Bodker B, Ambre J (1992). Rate of increase of plasma drug level influences subjective response in humans. *Psychopharmacology (Berl)* 107: 352–358.
- Harvey DM, Yasar S, Heishman SJ, Panlilio LV, Henningfield JE, Goldberg SR (2004). Nicotine serves as an effective reinforcer of intravenous drug-taking behavior in human cigarette smokers. *Psychopharmacology (Berl)* 175: 134–142.
- Hughes JR, Rose GL, Callas PW (2000). Do former smokers respond to nicotine differently from never smokers? A pilot study. *Nicotine Tob Res* 2: 255–262.
- Le Foll B, Goldberg SR (2006). Nicotine as a typical drug of abuse in experimental animals and humans. *Psychopharmacology (Berl)* 184: 367–381.
- Mooney M, White T, Hatsukami D (2004). The blind spot in the nicotine replacement therapy literature: assessment of the double-blind in clinical trials. *Addict Behav* 29: 673–684.
- Picciotto MR, Corrigall WA (2002). Neuronal systems underlying behaviors related to nicotine addiction: neural circuits and molecular genetics. *J Neurosci* 22: 3338–3341.
- Sofuoglu M, Yoo Y, Hill KP, Mooney M (2007). Self-administration of intravenous nicotine in male and female cigarette smokers. *Neuropsychopharmacology* (in press).

*Correspondence: Dr M Sofuoglu, Department of Psychiatry, Yale University, VA Connecticut Healthcare System, 950 Campbell Ave., Bldg. 36/116A4, West Haven, CT 06516, USA, Tel: +203 937 4809, Fax: +203 937 3478, E-mail: mehmet.sofuoglu@yale.edu