

Convergent dendron approach using a phenol triallyl building block: synthesis of a phenol dendron containing 27 allyl termini

Sylvain Nlate,^{a*} Jean-Claude Blais^b and Didier Astruc^{a*}

^a Groupe Nanosciences Moléculaires et Catalyse, LCOO (UMR CNRS N° 5802), Université Bordeaux I, 351 Cours de la Libération, Talence Cedex, France. E-mail: s.nlate@lcoo.u-bordeaux.fr; d.astruc@lcoo.u-bordeaux.fr

^b Laboratoire de Chimie Structurale Organique Biologique (EP CNRS N° 103), Université Paris VI, 4 Place Jussieu, 75252, Paris, France. E-mail: blaisjc@moka.ccr.jussieu.fr

Received (in Montpellier, France) 14th October 2002, Accepted 25th November 2002

First published as an Advance Article on the web 6th December 2002

A successful convergent dendron synthesis is achieved using a phenol triallyl (AB₃) building block by a sequence of reactions consisting of hydrosilylation-phenol protection. The key feature is the hydrosilylation using dimethylchloromethylsilane that avoids the side dehydrohalogenation previously found with linear branches containing β -hydrogens. The advantage of the AB₃ building block is the multiplication of the dendron branches by three at each generation, which allows the synthesis of the second-generation 27-allyl phenol dendron.

Since Vögtle's seminal report of the first iteration 25 years ago,¹ dendrimer syntheses have been achieved using a large variety of strategies and frameworks, and numerous potential applications are forecast.^{2–4} The divergent synthetic method has been the one used most because it is the easiest one to carry out, and also because it is the only one that can lead to large dendrimers even if defects cannot be discarded. The convergent method appeared in the early 90's with the reports of the groups of Fréchet,⁵ Miller⁶ and Moore⁷ and has been recently reviewed.⁸ It is elegant, and its great advantages are that it can lead to pure dendrons that are bifunctional: they can be decorated on the periphery for a given function or property, and they still bear another functional group at the focal point for attachment onto a dendritic core, nanoparticle, polymer, surface, etc. Although dendrons have a limited size because their synthesis cannot be extended due to steric bulk around the focal point that needs to be reactive, it is likely that they will play a key role in future nanotechnology involving molecular components. We have already shown that small dendrons bearing ferrocenyl-silyl or amidoferrocenyl termini are efficient sensors for the dihydrogenophosphate anion when, as part of thiolate ligands, they are attached to gold nanoparticles.⁹ So far, Fréchet's dendrons (AB₂ units) have been widely used in the literature, and AB₃ units are relatively scarce.⁸ We now report another convergent dendron synthesis based on a phenoltriallyl building block up to the second generation, i.e. 27 allyl groups. This molecular brick has already been used to synthesize dendrimers, but these attempts of convergent syntheses of dendrons have so far been frustrated by side reactions.¹⁰ Indeed the synthesis involved the nucleophilic substitution of the iodo group of the iodoalkyl branches by the phenol triallyl, but dehydrohalogenation became dominant. The present strategy prevents this side reaction by removing β -hydrogens on the termini.

Experimental section

Trichloro-trisilicium aryl sulfide 2

A mixture of triallyl sulfide dendron **1** (0.50 g, 1.93 mmol) and SiHMe₂(CH₂Cl) (0.945 g, 8.706 mmol, 1.1 mL), in Et₂O, was

stirred for 48 h at ambient temperature. The solution was filtered on Celite, and the solvent was removed under vacuum. After column chromatography on silica gel with a mixture of pentane : Et₂O (95 : 5) as eluent, **2** was obtained as a colorless oil (1.020 g, 1.74 mmol, 90%). Elemental analysis calcd for C₂₆H₄₉Si₃Cl₃S: H 8.45, C 53.44; found: H 8.35, C 53.00%. ¹H NMR (CDCl₃): δ_{ppm} 7.21 (s, C₆H₄, 4H); 2.72 (s, ClCH₂, 6H); 2.50 (s, SCH₃, 3H); 1.60 (m, CH₂, 6H); 1.06 (m, CH₂, 6H); 0.57 (m, SiCH₂, 6H); 0.05 (s, SiCH₃, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 143.3 (C_q, ArS); 140.0 (C_q, Ar); 127.4 (CH_{Ar}); 114.8 (CH_{Ar}); 43.1 (C_q-CH₂); 41.9 (CH₂); 30.2 (ClCH₂); 17.5 (CH₂CH₂Si); 15.9 (SCH₃); 14.2 (CH₂Si); -4.0 (SiMe).

Triiodo-trisilicium aryl sulfide 3

A mixture of **2** (0.500 g, 0.855 mmol) and NaI (1.77 g, 12.8 mmol), in 2-butanone, was stirred for 24 h at 80 °C. After removal of the solvent under vacuum, the residue was extracted with Et₂O (50 \times 3 mL). This solution was washed with 3 \times 50 mL of a saturated aqueous solution of Na₂S₂O₃. The solvent was removed under vacuum, which gave **3** as a yellow oil (0.624 g, 0.726 mmol 85%). Elemental analysis calcd for C₂₆H₄₉Si₃I₃S: H 5.75, C 36.37; found: H 5.55, C 35.82%. ¹H NMR (CDCl₃): δ_{ppm} 7.21 (s, C₆H₄, 4H); 2.50 (s, SCH₃, 3H); 1.96 (s, ICH₂); 1.60 (m, CH₂, 6H); 1.06 (m, CH₂, 6H); 0.60 (m, SiCH₂, 6H); 0.09 (s, SiCH₃, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 143.1 (C_q, ArS); 140.6 (C_q, Ar); 127.5 (CH_{Ar}); 114.8 (CH_{Ar}); 43.1 (C_q-CH₂); 41.9 (CH₂); 17.5 (CH₂CH₂Si); 16.00 (SCH₃); 15.7 (CH₂Si); -3.39 (SiMe); -13.60 (CH₂I).

9-allyl-trisilicium aryl sulfide 5

A mixture of triallylmethylphenol **4** (0.598 g, 2.620 mmol) and K₂CO₃ (0.413 g, 2.990 mmol), in 20 mL DMF, was stirred for 30 min at ambient temperature. The triiodo-trisilicium aryl sulfide **3** (0.500 g, 0.582 mmol) in solution in 5 mL DMF was added, and the reaction mixture was stirred at 80 °C 48 h. The solvent was removed under vacuum and the product was extracted with Et₂O. The organic layer was washed several times with water in order to remove DMF, then dried over

sodium sulfate, filtered and evaporated under vacuum. After column chromatography on silica gel using a mixture of pentane : Et₂O (9 : 1) as the eluent, **5** was obtained as a colorless oil (0.410 g, 0.353 mmol, 60%). Elemental analysis calcd for C₇₄H₁₀₆Si₃O₃S: H 9.21, C 76.62; found: H 9.23, C 76.52%. MALDI TOF mass spectrum, *m/z*: 1160.04 [*M*]⁺ (calcd 1159.98). ¹H NMR (CDCl₃): δ_{ppm} 7.19 (d, C₆H₄, 10H); 6.84 (d, C₆H₄, 6H); 5.54 (m, CH₂=CH, 9H); 5.00 (m, CH₂=CH, 18H); 3.49 (s, CH₂O, 6H); 2.45 (s, SCH₃, 3H); 2.43 (d, CH₂=CH-CH₂, 18H); 1.66 (m broad, CH₂, 6H); 1.20 (m broad, CH₂, 6H); 0.61 (m broad, CH₂, 6H); 0.05 (s, SiMe, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 159.41 (C_q, ArO); 145.00 (C_q, ArS); 137.25 (C_q, Ar); 134.78 (CH₂=CH); 134.62 (C_q, Ar); 127.46 (CH_{Ar}); 127.09 (CH_{Ar}); 117.43 (CH₂=CH); 114.64 (CH_{Ar}); 113.55 (CH_{Ar}); 60.15 (CH₂O); 43.53 (C_q-CH₂); 42.65 (C_q-CH₂); 42.01 (CH₂); 17.70 (CH₂); 16.01 (SCH₃); 14.64 (CH₂); -4.56 (SiMe).

Trichloro-trisilicium phenol dendron 6

A mixture of triallyl dendron **4** (1 g, 4.380 mmol) and SiH-Me₂(CH₂Cl) (2.279 g, 24.091 mmol, 1.2 mL), in Et₂O, was stirred for 48 h at ambient temperature. The solution was filtered on Celite, and the solvent was removed under vacuum. After column chromatography on silica gel with a mixture of pentane : Et₂O (95 : 5) as eluent, **6** was obtained as a colorless oil (2.209 g, 3.985 mmol, 91%). Elemental analysis calcd for C₂₅H₄₇Si₃Cl₃O: H 8.55, C 54.18; found: H 8.35, C 54.62%. ¹H NMR (CDCl₃): δ_{ppm} 7.12 (d, C₆H₄, 2H); 6.76 (d, C₆H₄, 2H); 4.65 (s, OH, 1H); 2.72 (s, ClCH₂, 6H); 1.60 (m, CH₂, 6H); 1.06 (m, CH₂, 6H); 0.57 (m, SiCH₂, 6H); 0.05 (s, SiCH₃, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 155.0 (C_q, ArO); 140.0 (C_q, Ar); 127.5 (CH_{Ar}); 114.8 (CH_{Ar}); 43.1 (C_q-CH₂); 41.9 (CH₂); 30.5 (ClCH₂); 17.5 (CH₂CH₂Si); 14.5 (CH₂Si); -4.1 (SiMe).

Triiodo-trisilicium phenol dendron 7

A mixture of **6** (2.178 g, 3.930 mmol) and, NaI (2.279 g, 58.945 mmol), in 2-butanone, was stirred for 24 h at 80 °C. After removal of the solvent under vacuum, the residue was extracted with Et₂O (50 × 3 mL). This solution was washed with 3 × 50 mL of a saturated aqueous solution of Na₂S₂O₃. The solvent was removed under vacuum, which gave **7** as a yellow oil (3.234 g, 3.903 mmol, 99%). Elemental analysis calcd for C₂₅H₄₇Si₃I₃O: H 5.72, C 36.24; found: H 5.64, C 37.01%. ¹H NMR (CDCl₃): δ_{ppm} 7.12 (d, C₆H₄, 2H); 6.76 (d, C₆H₄, 2H); 4.76 (s, OH, 1H); 1.94 (s, ICH₂, 6H); 1.60 (m, CH₂, 6H); 1.05 (m, CH₂, 6H); 0.58 (m, SiCH₂, 6H); 0.07 (s, SiCH₃, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 155.1 (C_q, ArO); 140.6 (C_q, Ar); 127.5 (CH_{Ar}); 114.8 (CH_{Ar}); 43.1 (C_q-CH₂); 41.9 (CH₂); 17.5 (CH₂CH₂Si); 15.7 (CH₂Si); -2.95 (SiMe); -13.1 (CH₂I).

Triiodo-trisilicium phenol propionate 8

13.35 mL (1.776 g, 9.655 mmol) of a fresh solution of C₂H₅COI, prepared by slowly adding 2.4 mL C₂H₅COCl to 5 g of Me₃SiI and 22.6 mL CH₂Cl₂, was added to a solution of **7** (2 g, 2.414 mmol) in 30 mL CH₂Cl₂. Then, NEt₂Prⁱ (0.030 g, 0.214 mmol) was also added. The reaction mixture was stirred for 16 h at ambient temperature. After removal of the solvent under vacuum, the residue was extracted with CH₂Cl₂ (3 × 40 mL). The organic solution was washed with a saturated aqueous solution of sodium bicarbonate, then with an aqueous solution saturated with Na₂S₂O₃ and dried over sodium sulfate. After removing the solvent under vacuum, the protected phenol dendron **8** was obtained as an orange-yellow oil. Elemental analysis calcd for C₂₈H₅₁Si₃I₃O: H 5.92, C 36.72; found: H 5.72, C 36.21%. (1.943 g, 2.196 mmol, 91%). ¹H NMR (CDCl₃): δ_{ppm} 7.24 (d, C₆H₄, 2H); 7.03 (d, C₆H₄,


2H); 2.59 (q, CH₃CH₂CO, 2H); 1.96 (s, ICH₂, 6H); 1.65 (m, CH₂, 6H); 1.05 (m, CH₂, 6H); 0.60 (m, SiCH₂, 6H); 0.08 (s, SiCH₃, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 175.8 (CO); 150.0 (C_q, ArO); 145.1 (C_q, Ar); 127.3 (CH_{Ar}); 120.8 (CH_{Ar}); 43.6 (C_q-CH₂); 41.8 (CH₂); 27.6 (CH₂); 17.6 (CH₂CH₂Si); 15.7 (CH₂Si); 9.1 (CH₃CH₂CO); -3.0 (SiMe); -13.1 (CH₂I).

9-Allyl-trisilicium phenol dendron 9


A mixture of triallylmethylphenol **9** (0.680 g, 2.978 mmol) and, K₂CO₃ (0.417 g, 2.978 mmol), in 20 mL DMF, was stirred for 30 min at ambient temperature. The protected triiodo-trisilicium phenol propionate **8** (0.440 g, 0.497 mmol) in solution in 5 mL DMF was added, and the reaction mixture was stirred at 80 °C for 48 h. Then 0.360 g K₂CO₃ and 0.600 mL water were added, and the reaction mixture was stirred at 80 °C for 24 h. The solvent was removed under vacuum and the product extracted with Et₂O. The organic layer was washed several times with water in order to remove DMF, then dried over sodium sulfate, filtered and evaporated under vacuum. After column chromatography on silica gel using a mixture of pentane : Et₂O (9 : 1) as the eluent, **9** was obtained as a colorless oil (0.380 g, 0.336 mmol, 68%). Elemental analysis calcd for C₇₃H₁₀₄Si₃O₄: H 9.28, C 77.60; found: H 9.31, C 77.09%. MALDI TOF mass spectrum, *m/z*: 1151.77 [*M* + Na]⁺ (calcd 1152.87). ¹H NMR (CDCl₃): δ_{ppm} 7.20 (d, C₆H₄, 6H); 7.09 (d, C₆H₄, 2H); 6.87 (d, C₆H₄, 6H); 6.67 (d, C₆H₄, 2H); 5.56 (m, CH₂=CH, 9H); 5.03 (m, CH₂=CH, 18H); 4.62 (s, OH, 1H); 3.52 (s, CH₂O, 6H); 2.43 (d, CH₂=CH-CH₂, 18H); 1.61 (m broad, CH₂, 6H); 1.14 (m broad, CH₂, 6H); 0.60 (m broad, CH₂, 6H); 0.06 (s, SiMe, 18H). ¹³C NMR (CDCl₃): δ_{ppm} 159.38 (C_q, ArO); 152.84 (C_q, ArO); 139.77 (C_q, Ar); 137.17 (C_q, Ar); 134.74 (CH₂=CH); 127.51 (CH_{Ar}); 127.40 (CH_{Ar}); 117.40 (CH₂=CH); 114.65 (CH_{Ar}); 113.52 (CH_{Ar}); 60.08 (CH₂O); 43.13 (C_q-CH₂); 42.60 (C_q-CH₂); 41.96 (CH₂); 17.63 (CH₂); 14.58 (CH₂); -4.61 (SiMe).

27-Allyl-9-siliciumphenol dendron 10

A mixture of 9-allyl-3-silicium phenol dendron **9** (0.500 g, 0.442 mmol) and, K₂CO₃ (0.070 g, 0.500 mmol), in 10 mL DMF, was stirred for 30 min. at ambient temperature. The protected triiododendron **8** (0.087 g, 0.098 mmol) in solution in 3 mL DMF was added, and the reaction mixture was stirred at 80 °C 72 h. Then, 0.052 g K₂CO₃ and 0.090 mL water were added, and the reaction mixture was stirred at 80 °C for 48 h. The solvent was removed under vacuum and the product was extracted with Et₂O. The organic layer was washed several times with water in order to remove DMF, then dried over sodium sulfate, filtered and evaporated under vacuum. After chromatography on silica gel using a mixture of pentane : Et₂O (9 : 1) as the eluent, **10** was obtained as a colorless oil (0.169 g, 0.044 mmol, 45%). Elemental analysis calcd for




Scheme 1




Scheme 2

$\text{C}_{244}\text{H}_{356}\text{Si}_{12}\text{O}_{13}$: H 9.36, C 76.43; found: H 9.54, C 75.72. MALDI TOF mass spectrum, m/z : 3856.37 [$M + \text{Na}$] $^+$ (calcd 3857.51). ^1H NMR (CDCl_3): δ_{ppm} 7.20 (d, C_6H_4 , 24H); 7.01 (d, C_6H_4 , 2H); 6.88 (d, C_6H_4 , 24H); 6.67 (d, C_6H_4 , 2H); 5.55 (m, $\text{CH}_2=\text{CH}$, 27H); 5.02 (m, $\text{CH}_2=\text{CH}$, 54H); 4.67 (s,

OH, 1H); 3.51 (s, CH_2O , 6H); 2.43 (d, $\text{CH}_2=\text{CH}-\text{CH}_2$, 54H); 1.63 (m broad, CH_2 , 18H); 1.15 (m broad, CH_2 , 18H); 0.61 (m broad, CH_2 , 18H); 0.06 (s, SiMe, 54H). ^{13}C NMR (CDCl_3): δ_{ppm} 159.27 (C_q , ArO); 159.00 (C_q , ArO); 139.08 (C_q , Ar); 137.07 (C_q , Ar); 134.63 ($\text{CH}_2=\text{CH}$); 127.30



Scheme 3



Scheme 4

(CH_{Ar}); 127.05 (CH_{Ar}); 117.32 ($CH_2=CH$); 113.40 (CH_{Ar}); 113.31 (CH_{Ar}); 60.02 (CH_2O); 42.92 (C_q-CH_2); 42.49 (C_q-CH_2); 41.86 (CH_2); 17.54 (CH_2); 14.43 (CH_2); -4.69 ($SiMe$).

Results and discussion

Dendritic growth has been carried out in parallel using the arylsulfide triallyl **1** and the phenol triallyl **4** that is an AB_3 unit for dendritic or hyperbranched polymer construction. The model **1** serves to test the divergent reaction of the allyl termini since it does not need protection. The compounds **1** and **4** have been reported, and are accessible by perallylation of the $CpFe^+$ complexes of the aryl ether and aryl sulfide respectively^{10b,11} (Scheme 1).

The growing strategy is based on the hydrosilylation of the allyl branches using chloromethyldimethylsilane that has

already been successfully used by Seyferth for dendrimer synthesis.¹² The chloromethyl termini are transformed into iodo-methyl by reaction with sodium iodide in refluxing butanone, then nucleophilic substitution by phenol triallyl leads to the next generation. With **1**, these reactions can be carried out in a straightforward way without any protection (Scheme 2).

With the phenol triallyl brick, it is of course necessary to protect the phenol group during the dendron construction, *i.e.* this synthesis is now a real convergent strategy for the construction of a dendron that is ultimately deprotected either to react on a protected dendron or onto another framework. Thus, after introduction of the iodo groups on the branches, the phenol function is protected by the propionate before synthesis of the first generation nona-allyl dendron (Scheme 3). The protection allows the condensation of the phenol triallyl AB_3 unit, and later, of the phenol nona-allyl AB_9





Fig. 1 MALDI-TOF mass spectrum of the 27-allyl dendron **10** (molecular peaks $[M + Na]^+$ at 3856.37).



Scheme 5

first-generation dendron. This convergent synthesis involving protection–deprotection has been carried out until the synthesis of the 27-allyl phenol that has been obtained in its deprotected form, *i.e.* as the second-generation dendron (Scheme 4). The first generation 9-allyl and second generation 27-allyl dendrons have been characterized by their standard spectroscopic and analytical data, but also especially by the prominent molecular peak in the MALDI TOF mass spectrum. In these mass spectra (see Fig. 1), impurities due to unfinished reactions are detected but are present only in relatively small amounts. The side peak at 3814.32 Dalton that is close to the molecular peak results from a small impurity corresponding to the phenoltriallyl AB₃ unit **4** in which one allyl group is missing (unfinished synthesis during the CpFe⁺-induced allylation reaction). The second side peak at 3557.27 Dalton corresponds to a dendron in which the ninth tripod is missing (again uncompleted introduction of the phenoltriallyl group **4**). The third side peak at 3515.36 Dalton corresponds to the combination of both impurities (unfinished triallylation and unfinished introduction of the phenoltriallyl group **4**). The peak at 1277.77 Dalton corresponds to exactly one third of the molecular peak, and is thus attributed to this molecular peak with a tricationic charge (m/z with $z = 3$).

This successful synthesis of the 27-allyl dendron phenol contrasts with the preceding attempt using a triiodoalkyl phenol bearing hydrogens in the position β relative to the iodo group. In that case, nucleophilic substitution proceeded correctly for two of the branches because it was faster than dehydrohalogenation. A rather pure 19-allyl phenol dendron was obtained (as shown in Scheme 5), however, the third

nucleophilic substitution was considerably slowed down due to the steric bulk of the two other nearby covalently attached dendrons. Thus, dehydrohalogenation of the third iodoalkyl branch proceeded quantitatively (Scheme 5). With the new triiodoalkyl dendron, the lack of β positions inhibits the dehydrohalogenation reaction.

Conclusion

The arylsulfide **1** and phenol derivative **4** are easily available in relatively good yields in only three steps from *p*-chlorotoluene via CpFe⁺-mediated synthesis. Thus, the chemistry reported here is easily carried out and should prove useful. The arylsulfide is a good model for the convergent synthesis from **4**. It is also a good model for the synthesis of aryl thiol dendrons that could bind gold nanoparticles and surfaces. The convergent synthesis of the 27-allyl phenol dendron from **4** was successful and proceeds in relatively good yields. Its advantage is that it multiplies the number of branches by three at each generation, allowing a very rapid growth up to 27 allyl branches in the second-generation dendron.

Acknowledgements

We are grateful to the Institut Universitaire de France (IUF), the Centre National de la Recherche Scientifique (CNRS) and the Universities Bordeaux I and Paris VI for financial support.

References

- 1 E. Buhleier, W. Wehner and F. Vögtle, *Synthesis*, 1978, 155.
- 2 (a) For a pioneering early review, see: D. A. Tomalia, A. N. Naylor and W. A. Goddard III, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 138; (b) For a comprehensive early review, see: N. Ardoin and D. Astruc, *Bull. Soc. Chim. Fr.*, 1995, **132**, 875.
- 3 For recent reviews, see ref. 4 and: (a) O. A. Matthews, A. N. Shipway and J. F. Stoddart, *Prog. Polym. Chem.*, 1998, **23**, 1; (b) V. Balzani, S. Campagna, G. Denti, A. Juris, S. Serroni and M. Venturi, *Acc. Chem. Res.*, 1998, **31**, 26; (c) H. Frey, *Angew. Chem., Int. Ed.*, 1998, **37**, 2193; (d) A. Archut and F. Vögtle, *Chem. Soc. Rev.*, 1998, **27**, 233; (e) D. K. Smith and F. Diederich, *Angew. Chem. Eur. J.*, 1998, **4**, 1353; (f) H.-F. Chow, T. K.-K. Mong, M. F. Nongrum and C.-W. Wan, *Tetrahedron*, 1998, **4**, 453; (g) J. P. Majoral and A.-M. Caminade, *Chem. Rev.*, 1999, **99**, 845; (h) A. W. Bosman, E. W. Jansen and E. W. Meijers, *Chem. Rev.*, 1999, **99**, 1665; (i) M. Fisher and F. Vögtle, *Angew. Chem., Int. Ed.*, 1999, **38**, 884; K. Inoue, *Prog. Polym. Sci.*, 2000, **25**, 453; (j) S. Hecht and J. M. J. Fréchet, *Angew. Chem., Int. Ed. Engl.*, 2001, **40**, 74; (k) V. Balzani, P. Ceroni, A. Juris, M. Venturi, S. Campagna, F. Puntoriero and S. Serroni, *Coord. Chem. Rev.*, 2001, **219–221**, 545; (l) Sekiguchi, V. Y. Lee and M. Nanjo, *Coord. Chem. Rev.*, 2001, **210**, 11.
- 4 For recent books, see: (a) G. R. Newkome, C. N. Moorefield and F. Vögtle, *Dendrimers and Dendrons. Concepts, Syntheses, Application*, Wiley-VCH, Weinheim, 2001; (b) *Dendrimers and Other Dendritic Polymers*, eds. J. M. J. Fréchet and D. A. Tomalia, Wiley, Chichester, 2001.
- 5 (a) C. J. Hawker and J. M. J. Fréchet, *J. Chem. Soc., Chem. Commun.*, 1990, 1010; (b) J. M. J. Fréchet, *Science*, 1994, **263**, 1710.
- 6 T. M. Miller and T. X. Neenan, *Chem. Mater.*, 1990, **2**, 346.
- 7 J. S. Moore and Z. Xu, *Macromolecules*, 1991, **24**, 5893.
- 8 (a) For a recent review on dendrons, see: S. M. Grayson and J. M. J. Fréchet, *Chem. Rev.*, 2001, **101**, 3819; (b) for a review of AB₃ units, see ref. 4, chapters 5–7 and, for selected examples, see: C. Cardona and A. E. Kaifer, *J. Am. Chem. Soc.*, 1998, **120**, 4023; D. K. Smith and F. Diederich, *Chem. Commun.*, 1998, 2501; V. Percec, W.-D. Cho and G. Ungar, *J. Am. Chem. Soc.*, 2000, **122**, 10 273; V. Percec, W.-D. Cho, G. Hungar and D. J. P. Yearley, *J. Am. Chem. Soc.*, 2001, **123**, 1302.
- 9 M.-C. Daniel, J. Ruiz, S. Nlate, J. Palumbo, J.-C. Blais and D. Astruc, *Chem. Commun.*, 2001, 2000.
- 10 (a) V. Sartor, L. Djakovitch, J.-L. Fillaut, F. Moulines, F. Neveu, V. Marvaud, J. Guittard, J.-C. Blais and D. Astruc, *J. Am. Chem. Soc.*, 1999, **121**, 2929; (b) V. Sartor, S. Nlate, J.-L. Fillaut, L. Djakovitch, F. Moulines, V. Marvaud, F. Neveu and J.-C. Blais, *New J. Chem.*, 2000, **24**, 351; (c) a divergent dendritic synthesis close to that reported in refs. 2a,b has been successfully achieved up to the 9th generation using the building block **4** and the poly-iodomethylsilyl dendrimers: J. Ruiz, G. Lafuente, S. Marcen, J.-C. Blais, S. Lazarre and D. Astruc, submitted for publication.
- 11 Cleavage of the aryl ether, but not of the aryl sulfide occurs during the CpFe⁺-induced perallylation and both complexes undergo iron-arene cleavage also by reaction of excess *t*-BuOK to provide the organic derivatives **1** and **2**^{10b}.
- 12 S. W. Krsda and D. Seyferth, *J. Am. Chem. Soc.*, 1998, **120**, 3604.