


This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Chemoselective Deprotection of α -Indole and Imidazole Ribonucleosides

Tilak Chandra^a; Kenneth L. Brown^a

^a Department of Chemistry and Biochemistry, Ohio University, Athens, Ohio, USA

To cite this Article Chandra, Tilak and Brown, Kenneth L.(2007) 'Chemoselective Deprotection of α -Indole and Imidazole Ribonucleosides', *Nucleosides, Nucleotides and Nucleic Acids*, 26: 1, 1 – 8

To link to this Article: DOI: 10.1080/15257770601052216

URL: <http://dx.doi.org/10.1080/15257770601052216>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

CHEMOSELECTIVE DEPROTECTION OF α -INDOLE AND IMIDAZOLE RIBONUCLEOSIDES

Tilak Chandra and Kenneth L. Brown □ *Department of Chemistry and Biochemistry, Ohio University, Athens, Ohio, USA*

□ *A series of 2',3'-isopropylidene and 5'-trityl-protected α -indole and α/β -benzimidazole and imidazole ribonucleosides were deprotected with different acids. Selectivity was achieved for 5'- versus 2',3'- deprotection by using formic acid in the α -indole ribonucleoside series. Treatment of α -indole ribonucleosides with a mixture of formic acid and ether at room temperature afforded 2',3'-deprotected α -ribonucleosides, whereas treatment of the α -benzimidazole ribonucleosides with the same acid afforded the 5'-deprotected ribonucleoside without any 2', 3'-deprotected products. The structures of these ribonucleosides were elucidated with 2D (NOESY, COSY, and HMQC) NMR spectroscopy.*

Keywords Chemoselective; coenzyme B₁₂; biosynthesis; glycosylation


INTRODUCTION

Semisynthesis of cobalamins^[1–6] with altered axial nucleotide ligands requires cobyrinic acid^[7] and an α -ribonucleotide^[8] as precursors. Analogs of coenzyme B₁₂ (5'-deoxyadenosylcobalamin, AdoCbl) with altered axial nucleotide ligands, such as B3-deazaAdoCbl (Figure 1) are of interest as probes of the function, if any, of the axial nucleotide in the enzymatic activation of AdoCbl for carbon-cobalt bond homolysis^[9,10]. Literature reports on the synthesis and deprotection of indole α -ribonucleosides are limited. Deprotection is a crucial step for these unstable α -ribonucleosides,^[11–18] in order to retain the α -configuration and to achieve the synthesis of AdoCbl. Trityl (Tr), dimethoxytrityl (DMTr), and isopropylidene groups are commonly used for the protection of 5-hydroxyl and 2, 3-hydroxyl groups in both carbohydrate and nucleoside chemistry.^[19] Strong protic

Received 7 December 2005; accepted 15 June 2006.


This research was supported by the National Institute of General Medical Sciences, Grant GM 48858.

Address correspondence to Kenneth L. Brown, Department of Chemistry and Biochemistry, Ohio University, Athens, OH 45701-3132. E-mail: brownk3@ohio.edu

**SCHEME 1** Deprotection of α -indole ribonucleosides.

using 2-fluoromethylpyridinium tosylate as a condensing reagent^[22] or by direct glycosylation.^[14] 2D NMR and X-ray crystallography confirmed the anomeric configuration of these indoline ribonucleosides.^[15] Protected indole ribonucleosides^[13] (**1-3**, Scheme 1) were prepared from the corresponding indoline α -ribonucleosides^[14,15] by manganese dioxide oxidation^[13] at elevated temperature. Benzimidazole, 5,6-dimethylbenzimidazole and imidazole α -ribonucleosides^[15] (**11 & 12**) and (**13 & 14**) were similarly prepared using 2-fluoromethyl pyridinium tosylate as a condensing reagent^[22] (Scheme 2). These glycosylation reactions produce mixtures of α - (70–85%) and β -ribonucleosides (15–30%).

Upon treatment with formic acid/ether (2:3) at room temperature (Scheme 1) the indole ribonucleosides afforded the 2',3'-deprotected indole ribonucleoside in fairly good yield, without affecting the trityl group. In contrast, the α -ribonucleosides of dimethylindole, indole and 5-bromoindole, on treatment with aqueous acetic acid at 50–60°C, afforded the corresponding 5'-deprotected α -ribonucleoside without affecting the isopropylidene group or causing deglycosylation. At higher temperature and longer time we observed deglycosylation and charring without any desired product. While it is known that 2',3'-O-isopropylidene- β -D-ribofuranosyl-2,5,6-trichloroindole^[23] and other halogenated indole ribonucleosides produce 2',3'-deprotected indole ribonucleosides upon prolonged heating with acetic acid and water (4:1), for the current α -indole ribonucleosides we do not observe any 2',3'-deprotected products. Instead,



Reagents and conditions:

- (a) TMS-protected bases, 2-fluoromethyl pyridinium tosylate, DIEA, methylene chloride, -30 °C to 0 °C
 (b) formic acid:ether, (2:3), room temp, 3-4 h, or 1% TFA, methylene chloride, room temp, 30 min or aqueous AcOH (80:20), 50-60 °C

SCHEME 2 Preparation and deprotection of α/β -benzimidazole/imidazole ribonucleosides.

prolonged heating of these α -indole ribonucleosides at higher temperature in acetic acid produced only charring.

The 1-(2,3-O-isopropylidene-5'-O-triphenylmethyl- α -D-ribofuranosyl)-benzimidazole, and 5,6-dimethylbenzimidazole ribonucleosides (**11** & **12**), however, react differently (Scheme 2). The trityl group can be easily removed by treating the protected ribonucleoside with formic acid/ether, heating with aqueous acetic acid, or using 1% TFA in methylene chloride, without affecting the isopropylidene.

A series of protected α and β ribosides were prepared by the Mukaiyama^[22] method and isolated in pure form and thoroughly characterized by two-dimensional NMR techniques (COSY, HMQC, and NOESY). All protected α -ribonucleosides have a NOE between the isopropylidene methyl protons and imidazole proton, whereas the β -anomer does not have a NOE between these protons. The methyl protons of the isopropylidene had a NOE with the imidazole proton in 5'-deprotected α -benzimidazole ribonucleoside **15**, but the β -benzimidazole does not have a NOE between those protons (2D NOESY).

Structural proof for the α -anomer was also confirmed with the help of NOE difference spectroscopy. Irradiation of the isopropylidene methyl protons of 5'-deprotected benzimidazole ribonucleoside, **15**, caused a 10% positive NOE enhancement of the imidazole proton, and the precursor

11 (protected ribonucleoside) shows a 12% NOE enhancement upon irradiation of the methyl protons. The signals for the methyl protons of the isopropylidene in the benzimidazole ribonucleoside (2',3'-O-isopropylidene- α -D-ribofuranosyl) benzimidazole, **15**, were visible at δ 1.279 and 1.382 ppm ($d = 0.103$). The anomeric proton signal appeared at δ 6.33 ppm with a J value of 4.5 Hz, which further confirms the α -configuration of the ribonucleoside, whereas the anomeric proton signal for compound **17** was at δ 6.07 ppm with J value of 3.2 Hz and a methyl proton separation of 0.269 ppm. Similarly, the deprotected indole ribonucleosides also showed the same NMR pattern as the protected α -ribonucleosides.

EXPERIMENTAL SECTION

All reactions were carried out in oven or flame-dried glassware under a nitrogen atmosphere. Solvents were distilled prior to use. Dichloromethane, benzene and ether were distilled from calcium hydride. Purification of reaction products was carried out by flash chromatography using silica gel (230–400 mesh). All reagents were commercially available and used without further purification. All reactions were monitored by thin-layer chromatography (TLC) using Merck silica gel 60, F-254. Column chromatography was conducted using flash silica gel.

Physical Measurements

^1H NMR and ^{13}C NMR spectra were recorded on a Varian INOVA-500 or VXR-400 NMR spectrometers using the residual proton resonance of the solvent as an internal reference at 25°C. Two-dimensional NMR (COSY, NOESY) spectra were obtained at 25°C on a Varian INOVA-500 NMR spectrometer using TMS as an internal reference. The multiplicities of the ^{13}C NMR signals were determined by the HMQC and DEPT technique. Mass data were obtained at University of Illinois using a micromass Quattro-I mass spectrometer.

1-(5-O-Triphenylmethyl- α -D-ribofuranosyl)indole (4): A mixture of the protected ribonucleoside **1**^[13] (0.700 g, 1.3 mmol) in ether:formic acid (3:2) (10 mL) was stirred for 3–4 hours at room temperature while TLC (ethyl acetate:hexane 1:1) was used to monitor the progress of the reaction. Upon completion, the solvent was removed on a rotary evaporator, the residue was dissolved in ethyl acetate, and washed with saturated sodium bicarbonate solution. The ethyl acetate layer was concentrated and the residue was purified by flash chromatography. Yield: 60%; R_f : 0.2; white solid; ^1H NMR (DMSO- d_6): δ 3.57 (dd, $J = 2.0, 5.5$ Hz, 1H, 5'), 3.70 (d, $J = 7.7$ Hz, 1H, 5''), 3.84 (bt, 1H, 2'), 4.0 (bs, 1H), 5.0 (d, $J = 3.4$ Hz, 3'), 5.49 (d, $J = 9.5$ Hz, 1H, 1'), 6.47 (d, $J = 8.0$ Hz, 1H, Ar), 6.70 (t, $J = 7.5$ Hz, 1 H, Ar), 6.93 (s, 1H, Ar), 7.1 (d, $J = 7.5$ Hz, 6H, Ar), 7.2 (t, 3H,

trityl), 7.27 (t, $J = 7.7$ Hz, 6H, trityl), 7.44 (d, 1H, $J = 8.0$ Hz, Ar); ^{13}C NMR (DMSO- d_6): 65.0, 66.64, 68.45, 71.43, 81.98 (C1'), 110.8 (CH), 119.0 (CH), 121.1 (CH), 121.7 (CH), 121.8 (Cquat), 126.0 (CH), 126.48, 127.48 (CH), 128.0 (Cquat), 130.1 (CH), 137.25, 145.9 (Cquat); HRMS: m/z : 492.2166 (calcd for $\text{C}_{32}\text{H}_{30}\text{NO}_4$; 492.2174) (M + H); MS: (FAB) m/z 492, 394, 324, 243.

1-(2, 3-isopropylidene- α -D-ribofuranosyl)5,6-dimethylindole (8): Compound **2**^[13] (150 mg, 0.26 mmol) was dissolved in aqueous acetic acid (5 mL) and heated for 5 hours at 50–60°C in a oil bath. After complete conversion of the starting material, the reaction mixture was cooled and then concentrated to dryness to give a residue. The residue was purified by flash column chromatography (benzene:ether; 50:50) to give 70 mg of compound **8** as a viscous oil. Yield: 80%; R_f : 0.7; ^1H NMR (CDCl_3): δ 1.38 (s, 3H, CH_3 , isopropylidene), 1.64 (s, 3H, CH_3 , isopropylidene), 2.35 (s, 3H, CH_3), 2.40 (s, 3H, CH_3), 3.73 (dd, $J = 4.0, 8.0$ Hz, 1H, 5'), 3.82 (dd, $J = 3.24, 8.9$ Hz, 1H, 5'), 4.22 (q, 1H, CH, 4'), 4.93–4.95 (m, 1H, CH, 3'), 5.03–5.05 (m, 1H, CH, 2'), 6.06 (d, $J = 3.5$ Hz, 1H, CH, 1'), 6.46 (d, $J = 3.0$ Hz, 1H, indole), 7.10 (d, $J = 3.0$ Hz, 1H, indole), 7.27 (s, 1H, Ar), 7.39 (s, 1H, Ar); ^{13}C NMR (CDCl_3): δ 19.97 (CH_3), 20.70 (CH_3), 25.36 (CH_3), 27.26 (CH_3), 62.52 (CH_2 , 5'), 80.18 (C 3'), 84.03 (C 2'), 84.83 (C 4'), 91.15 (C 1'), 103.13 (CH, Ar), 110.52 (CH), 114.70 (Cquat, isoprop), 121.25 (CH), 123.72 (CH), 126.01 (Cquat), 127.35 (Cquat), 130.75 (Cquat), 146.22 (Cquat); HRMS: m/z : 318.1704 (calcd for $\text{C}_{18}\text{H}_{24}\text{NO}_4$; 318.1704) (M + H).

1-(2,3-O-isopropylidene- α -D-ribofuranosyl)benzimidazole (15): The benzimidazole ribonucleoside^[15,17] **11**, was similarly deprotected as described for the indole nucleoside **4** using either the mixture of (formic acid:ether) or aqueous acetic acid (method described for the compound **8**) Yield: 80–90%; R_f : 0.3; White foam; ^1H NMR (CDCl_3): δ 1.279 (s, 3H, CH_3 , isopropylidene), 1.382 (s, 3H, CH_3 , isopropylidene), 3.85 (dd, $J = 3.5, 8.5$ Hz, 1H, 5'), 3.96 (dd, $J = 2.0, 9.5$ Hz, 1H, 5'), 4.42 (bt, 1H, CH, 4'), 4.92 (t, $J = 6.5$ Hz, 1H, CH, 2'), 5.01 (d, $J = 6$ Hz, 1H, CH, 3'), 6.33 (d, $J = 4.5$ Hz, 1H, CH, 1'), 7.21–7.28 (m, 3H, Ar), 7.76 (d, $J = 7.5$ Hz, 1H, Ar), 8.24 (s, 1H, imidazole); ^{13}C NMR (CDCl_3): δ 24.18 (CH_3), 25.65 (CH_3), 63.87 (CH_2 , 5'), 80.06 (C 2'), 82.35 (C 3'), 83.45 (C 4'), 87.21 (C 1'), 109.90 (CH, Ar), 113.34 (Cquat, isoprop), 119.61 (CH), 122.47 (CH), 123.06 (CH), 132.69 (Cquat), 142.30 (CH), 142.58 (Cquat); HRMS: m/z : 291.1344 (calcd for $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}_4$; 291.1343) (M + H); MS: (FAB) m/z 291, 195, 152, 119.

1-(2,3-O-isopropylidene- β -D-ribofuranosyl)benzimidazole (17): Yield: 80–90%; R_f : 0.3; White crystalline solid; ^1H NMR (CDCl_3): δ 1.40 (s, 3H, CH_3 , isopropylidene), 1.67 (s, 3H, CH_3 , isopropylidene), 3.85 (dd, $J = 2.83, 8.91$ Hz, 1H, 5'), 4.01 (dd, $J = 2.43, 9.72$ Hz, 1H, 5'), 4.51 (d, 1H, CH, 4'), 4.65 (bs, 1H, OH), 4.98–4.99 (m, 1H, CH, 2'), 5.05 (dd, $J = 2.43, 3.64$ Hz, 1H, CH, 3'), 6.07 (d, $J = 3.24$ Hz, 1 H, CH, 1'), 7.29–7.32 (m, 2H, Ar), 7.52 (d, $J = 7.29$ Hz, 1H, Ar), 7.75 (d, 1H, $J = 6.89$ Hz, 1H, Ar), 8.39 (s, 1H,

imidazole); ^{13}C NMR (CDCl_3): δ 25.25 (CH_3), 27.28 (CH_3), 62.05 (CH_2 , 5'), 81.50 (C 3'), 85.36 (C 2'), 86.52 (C 4'), 92.81 (C 1'), 110.46 (CH, Ar), 114.18 (Cquat, isoprop), 119.87 (CH), 122.82 (CH), 123.44 (CH), 132.41 (Cquat), 142.83 (CH), 143.33 (Cquat); HRMS: m/z : 291.1346 (calcd for $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}_4$; 291.1343) (M + H); MS: (FAB) m/z 291, 195, 152, 119.

REFERENCES

1. Brown, K.L.; Li, J. Activation parameters for the carbon-cobalt bond homolysis of coenzyme B₁₂ induced by the B₁₂-dependent ribonucleotide reductase from *Lactobacillus Leichmannii*. *J. Am. Chem. Soc.* **1998**, 120, 9466–9474.
2. Brown, K.L. Chemistry and enzymology of vitamin B₁₂. *Chem. Rev.* **2005**, 105(6), 2075–2149. Washington, DC, United States.
3. Brown, K.L. The enzymatic activation of coenzyme B₁₂. *Dalton Trans.* **2006**, 9, 1123–1133.
4. Hamza, M.S.A.; Zou, X.; Banka, R.R.; Brown, K.L.; van Eldik, R. Kinetic and thermodynamic studies on ligand substitution reactions and base-on/base-off equilibria of cyanoimidazolylcobamide, a vitamin B₁₂ analog with an imidazole axial nucleoside. *Dalton Trans.* **2005**, (4), 782–787.
5. Brown, K.L.; Zou, X.; Li, J.; Chen, G. Enzymatic activity of coenzyme B₁₂ derivatives with altered axial nucleotides: Probing the mechanochemical triggering hypothesis in ribonucleotide reductase. *Inorg. Chem.* **2001**, 40, 5942–5947.
6. Brown, K.L.; Cheng, S.; Zou, X.; Li, J.; Chen, G.; Valente, E.J.; Zubkowski, J.D.; Marques, H.M. Structural and enzymic studies of a new analog of coenzyme B₁₂ with an α -adenosyl upper axial ligand. *Biochemistry* **1998**, 37, 9704–9715.
7. (a) Renz, P. Intermediates in the biosynthesis of vitamin B₁₂, *Methods in Enzymology* 1971, 18 C, 82//;92. (b) Bonnett, R.; Godfrey, J.M., Redman, D.G. Cobyric acid and related compounds from vitamin B₁₂. *J. Chem. Soc. [section C organic]* **1969**, 8, 1163–1166.
8. Friedrich, W.; Gross, G.; Bernhauer, K.; Zeller, P. Syntheses in the vitamin B₁₂ series. IV. Partial synthesis of vitamin B₁₂. *Helv. Chim. Acta.* **1960**, 43, 704–712.
9. Hay, B.P.; Finke, R.G. Thermolysis of the cobalt-carbon bond of adenosylcobalamin. 2. Products, kinetics, and cobalt-carbon bond dissociation energy in aqueous solution. *J. Am. Chem. Soc.* **1986**, 108, 4820–4829.
10. Hay, B.P.; Finke, R.G. Thermolysis of the cobalt-carbon bond in adenosylcorrins. 3. Quantification of the axial base effect in adenosylcobalamin by the synthesis and thermolysis of axial base-free adenosylcobinamide. insights into the energetics of enzyme-assisted cobalt-carbon bond homolysis. *J. Am. Chem. Soc.* **1987**, 109, 8012–8018.
11. Stevens, J.D.; Ness, R.K.; Fletcher, H.G., Jr.; Hewitt, G., Jr. Syntheses with partially benzylated sugars. XI. Studies on the synthesis of the anomeric 5,6-dimethyl-1-D-ribofuranosylbenzimidazoles ribazoles. Comparison of the condensation of 2, 3, 5-tri-O-benzoyl-D-ribofuranosyl bromide and 2,3,5-tri-O-benzyl-D-ribofuranosyl chloride with 5,6-dimethylbenzimidazole. *J. Org. Chem.* **1968**, 33(5), 1806–1810.
12. Holly, F.W.; Shunk, C.H.; Peel, E.W.; Cahill, J.H.; Lavigne, J.B.; Folkers, K. Vitamin B₁₂. XIX. Synthesis of the 1-D-ribosides of 5, 6-dimethylbenzimidazole. *J. Am. Chem. Soc.* **1952**, 74, 4521–4525.
13. Chandra, T.; Zou, S.; Brown, K.L. Low temperature dehydrogenation of α -indoline nucleosides. *Tetrahedron Lett.* **2004**, 45, 7783–7786.
14. Chandra, T.; Brown, K.L. Direct glycosylation: Synthesis of α -indoline ribonucleosides. *Tetrahedron Lett.* **2005**, 46(12), 2071–2074.
15. Brown, K.L.; Chandra, T.; Zou, S.; Valente, E. Regioselective glycosylation: synthesis of α -indoline nucleosides. *Nucleosides, Nucleotides and Nucleic Acids* **2005**, 24(8), 1147–1165.
16. Chandra, T.; Brown, K.L. Deprotection of α -imidazole/benzimidazole ribonucleosides by catalytic carbon tetrabromide initiated photolysis. *Tetrahedron Lett.* **2005**, 46(49), 8617–8619.
17. Chandra, T.; Zou, S.; Valente, E.J.; Brown, K.L. Regio- and stereo-selective glycosylation: Synthesis of 5-haloimidazole α -ribonucleosides. *J. Org. Chem.* **2006**, 71(31), 5000–5003.
18. Pichumani, K.; Chandra, T.; Xiang, Z.; Brown, K.L. Experimental observation of cross correlated relaxation between H_{1'} chemical shift anisotropy and H_{1'}-H_{2'} dipolar relaxation mechanisms in

- ribonucleosides: Application to characterize their anomeric configuration. *J. Phys. Chem. B.* **2006**, 106, 5–8.
19. Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*. 3rd edn., Wiley, New York, 1999.
 20. Bessodes, M.; Komiotis, D. Antonakis K. Rapid and selective detritylation of primary alcohols using formic acid. *Tetrahedron Lett.* **1986**, 27, 579–582.
 21. MacCoss, M.; Cameron, D.J. Facile detritylation of nucleoside derivatives by using trifluoroacetic acid. *Carbohydr. Res.* **1978**, 60, 206–209.
 22. Mukaiyama, T.; Hashimoto, Y.; Hayashi, Y.; Shoda, S-i. Stereoselective synthesis of α -ribonucleosides from 1-hydroxy sugars by using 2-fluoropyridinium tosylate. *Chem. Lett.* **1984**, 57–560.
 23. Chen, J.J.; Wei, Y.; Drach, J.C.; Townsend, L.B. Synthesis and antiviral evaluation of trisubstituted indole *N*-nucleosides as analogues of 2,5,6-trichloro-1-(β -D-ribofuranosyl)benzimidazole (TCRB). *J. Med. Chem.* **2000**, 43, 2449–2456.