

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Aldehydo-oligonucleotides for Bioconjugation

M. Dechamps^a; E. Sonveaux^a

^a Catholic University of Louvain, School of Pharmacy, Bruxelles, Belgium

To cite this Article Dechamps, M. and Sonveaux, E.(1998) 'Aldehydo-oligonucleotides for Bioconjugation', *Nucleosides, Nucleotides and Nucleic Acids*, 17: 4, 697 – 709

To link to this Article: DOI: 10.1080/07328319808004669

URL: <http://dx.doi.org/10.1080/07328319808004669>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ALDEHYDO-OLIGONUCLEOTIDES FOR BIOCONJUGATION

Dechamps, M. and Sonveaux*, E.

Catholic University of Louvain, School of Pharmacy,
UCL 7340, Avenue E. Mounier 73, B-1200 Bruxelles, Belgium

Abstract: the synthesis of a new phosphoramidite building block with a masked aldehyde function is described. It was incorporated in an oligodeoxyribonucleotide. Coupling to amino derivatives by reductive amination was performed in high yield after unmasking the aldehyde function.

The immobilization of single stranded DNA to macromolecules is important for the design of diagnostic kits and biologically active substances. Immobilized oligonucleotides are indeed used to fish out a complementary sequence from a mixture of DNA or RNA^{1,2}. The coupling of oligonucleotide effectors to peptides^{3,4}, basic polymers^{5,6} and proteins^{7,8} enhances their cellular penetration, and could allow their targeting to specific tissues or determine their intracellular routing. The link between the oligonucleotide and its support or vector may be reversible (typically a disulfide bridge) or irreversible (typically a thioether or an amide bond). The formation of an amine link by reductive amination was rarely performed. It is rather surprising, as reductive amination is largely used in protein chemistry and does not interfere with the functions present in DNA. The reason is probably because suitable aldehyde building blocks for automatic DNA synthesis are not currently available. A documented strategy to introduce aldehyde functions at the 3'-end of an oligonucleotide is the sodium periodate oxydation of a terminal ribonucleotide⁵. This procedure may be tricky however as β -elimination in the presence of an amine could be a major side-

reaction (Figure 1). Eliminations of this kind are indeed observed when apurinic DNA is treated by piperidine in the Maxam-Gilbert sequencing protocol⁹.

Oligonucleotides bearing aldehyde functions could be conjugated to another molecule by either the hydrolyzable C=N imine, oxime, hydrazone or semicarbazone bond or the hydrolytically stable C-N bond obtained by reduction of the former. The hydrolytic stability of the C=N bond can be tuned according to the aim pursued (e.g. imines are far less stable than oximes¹⁰). Bioreversible bonds are of prime importance in the antisense research but the sole bioreversible bond available nowadays is the disulfide bridge. An enlargement of the palette of available bond types is thus convenient and in this connection, a reliable access to aldehydo-oligodeoxyribonucleotide is valuable.

We report here the synthesis of a nucleotide analog **8** (Figure 2) that allows the introduction of an aldehyde function in any position of an oligomer and in multiple copies, if necessary. The organocadmium **2**, obtained from 10-bromomagnesium-1-decene¹¹, was reacted with the readily accessible^{12,13} protected deoxyribosyl chloride **1**. The coupling yield was good (85%), but the stereochemistry at the anomeric center was not controlled (R, 65%, S, 35%). The two epimers were separated by column chromatography and the synthesis was pursued with the S one¹⁴. Osmium tetroxide oxidation¹⁵, followed by silyl protection gave the fully protected nucleoside analog **5**. The *t*-butyldimethylsilyl group was used to protect the glycol because it is robust, compatible with the reagents used for automated DNA synthesis and easily removed at the oligonucleotide level. It is indeed the best 2'-O protecting group for RNA synthesis on solid support¹⁶. The selective aminolysis of the ester functions of **5** led to **6**, that was dimethoxytritylated on the primary alcohol function and then phosphitylated¹⁷. Compound **8** is a building block tailored for the automatic phosphoramidite-type DNA synthesis.

We then incorporated **8** in the sequence GTCGTGAC-(**8**)-GGGAAAAC. The trityl-on oligomer was cleaved from the glass beads by concentrated ammonia, detritylated, treated with tetrabutylammonium fluoride in THF to remove the silyl




FIGURE 1

Possible β -elimination in oxidized 3' terminal ribonucleotide

protections and desalted. The resulting oligomer **9** was homogeneous in reversed phase C₁₈ HPLC and PAGE. The oxidation of the glycol function of **9** by sodium periodate to give **10** (Figure 3) was conveniently followed by reversed phase HPLC. The scintillation counting after reduction of **10** by NaBT₄ proved that the oxidation was quantitative.

Oligonucleotide **10** was reductively coupled (NaBH₃CN) with amino derivatives of biotin, namely biotin hydrazide to give **11** and D-biotinyl-1,8-diamino-3,6-dioxaoctane, to give **12**. These couplings were also monitored by reversed phase HPLC, the conjugates being retarded by 1-1.5 minutes as compared to the aldehyde-oligomer **10**. The reactions were practically quantitative.

To get a further proof of the incorporation of one mole of biotin per mole of oligonucleotide, several mixtures of **12** + **9** (used as an internal standard) and avidin in different molar ratios were prepared. The mixtures were ultrafiltrated with a membrane cutoff that allowed the separation of the free oligonucleotides from avidin. They were quantified in the filtrate by reversed phase HPLC. The result of this titration of the conjugate **12** by avidin proved that indeed one molecule of biotin was immobilized per oligomer (see experimental part).

In conclusion, we described a practical synthesis of **8**. It is a convenient building block to introduce an aldehyde function in oligodeoxyribonucleotides and perform couplings with amino derivatives. Both 3' and 5' ends of glycol-oligonucleotides of type **9** remain free, so that attachment of other molecules at the extremities as well as 5' ³²P labeling remain feasible.




FIGURE 2




FIGURE 3

EXPERIMENTAL

The proportions in solvent and eluent mixtures are expressed as volumic fractions. The NMR spectra were recorded on a 500 MHz (^1H , 125 MHz ^{13}C , 81 MHz ^{31}P) instrument, except when otherwise stated. Chemical shifts are given in ppm relative to TMS (^1H , ^{13}C) or phosphoric acid (^{31}P). Biotin hydrazide was from Sigma and D-biotinyl-1,8-diamino-3,6-dioxaoctane from Boehringer Mannheim.

(2-R,3-S,5-S)-1-O,3-O-di(p-chlorobenzoyl)-1,2,3,5-tetrahydroxy-2,5-anhydropentadec-14-ene 3

The reaction was performed under an argon atmosphere. A solution of 10-bromo-1-decene (40 mmoles) in dry THF (10 ml) was added portionwise to magnesium turnings (41 mmoles) magnetically stirred in dry THF (50 ml) containing 2-3 drops of

1,2-dibromoethane. The rate of addition was such that a constant reflux was maintained. Ten minutes after the end of addition, more THF (20 ml), and solid cadmium chloride (21 mmol, dried at 100°C for 1 h, then kept under argon) was added. The mixture was kept for 1/2 h at room temperature. Dry hexane (30 ml) was added, followed by solid chloride **1** (10 mmol, in 4-5 fractions). The mixture was warmed (67°C) for 1 h, then poured on cold water (100 ml, 0°C). The pH was adjusted to neutral with acetic acid. The water phase was extracted with toluene. The organic phase was washed (sat. NaHCO₃, H₂O), dried (MgSO₄), filtered and evaporated. The residue was chromatographed on silica (hexane/ethyl acetate, 95/5) (Yield: 85%; R, 65%, S, 35%). S Epimer: R_f = 0.15. 200 MHz ¹H NMR (CDCl₃): 7.96 and 7.41 (2 x m, ClBz), 5.78 (m, CH=CH₂), 5.45 (m, ClBzOCH), 4.95 (m, CH=CH₂), 4.48 (m, ClBzOCH₂CH), 4.32 (m, ClBzOCH₂CH), 4.20 [m, CH(CH₂)₈CH=CH₂], 2.23 [ddd, 1H, ring CH₂ (H exo)], 2.04 [m, CH(CH₂)₇CH₂CH=CH₂], 1.90 [m, ring CH₂ (H endo)], 1.65 [m, CHCH₂(CH₂)₇CH=CH₂], 1.30 [m, CHCH₂(CH₂)₆CH₂CH=CH₂]. ¹³C NMR, 165.4, 139.8, 139.1, 131.1, 129.03 and 128.5 (ClBz and CH=CH₂), 114.2 (CH=CH₂), 82.1 and 79.5 (CHOCH), 77.6 (ClBzOCH), 65.1 (ClBzOCH₂), 38.5 (ring CH₂), 35.3, 33.9, 29.7, 29.6, 29.3, 29.2, 29.1 and 25.8 [(CH₂)₈]. C, H analysis, calc. C, 65.35, H, 6.43; found, C, 65.42, H, 6.53.

(2-R,3-S,5-S)-1-O,3-O-di(p-chlorobenzoyl)-1,2,3,5,14,15-hexahydroxy-2,5-anhydropentadecane **4**

Compound **3** (5 mmol) was dissolved in an acetone/water mixture (9:1, 30 ml) kept at 0°C. *N*-methylmorpholin *N*-oxide (7.5 mmol) was added, followed by osmium tetroxide (0.5 mmol, i.e. 0.5 ml of a 1M solution in *t*-butanol). The mixture was magnetically stirred and the progress of the reaction was followed by tlc (CH₂Cl₂/CH₃OH, 95:5; **3**, R_f = 0.8, **4**, R_f = 0.15). The mixture was evaporated after 6 h and partitioned between water and CH₂Cl₂. The organic phase was dried (MgSO₄), filtered and evaporated. The compound was purified by column chromatography (eluent, CH₂Cl₂). It is a colorless oil (Yield, 90%). ¹H NMR (CDCl₃): 7.97 and 7.42 (2 x m, ClBz), 5.46 (m, ClBzOCH), 4.50 (m, ClBzOCH₂CH), 4.34

(m, ClBzOCH₂CH), 4.25 [m, CH(CH₂)₈CH(OH)-CH₂(OH)], 3.67 [m, CH(CH₂)₈CH(OH)-CH₂(OH)], 3.40 [m, CH(CH₂)₈CH(OH)-CH₂(OH)], 3.05 [large signal, CH(CH₂)₈CH(OH)-CH₂(OH)], 2.22 [ddd, 1H, ring CH₂ (H exo)], 1.90 [m, ring CH₂ (H endo)], 1.75 and 1.40 [2 x m, CH(CH₂)₇CH₂CH(OH)-CH₂(OH) and CHCH₂(CH₂)₇CH(OH)-CH₂(OH)], 1.25-1.35 [m, CHCH₂(CH₂)₆CH₂CH(OH)-CH₂(OH)]. ¹³C NMR, 165.2, 139.5, 130.8, and 128.0 (ClBz), 80.6 and 79.3 (CHOCH), 77.3 (ClBzOCH), 72.1 (CHOH), 66.6 (CH₂OH), 64.6 (ClBzOCH₂), 37.5 (ring CH₂), 35.7, 33.0, 29.5, 29.4, 29.3, 25.8 and 25.4 [(CH₂)₈]. C, H analysis, calc. C, 61.38, H, 6.39; found, C, 61.09, H, 6.38.

(2-R,3-S,5-S)-1-O,3-O-di(p-chlorobenzoyl)-14-O-15-O-di(*t*-butyldimethylsilyl)-1,2,3,5,14,15-hexahydroxy-2,5-anhydropentadecane 5

The reaction was performed under argon with magnetic stirring. The diol **4** (2 mmoles) was dissolved in DMF (5 ml), dried imidazole (5 mmoles) was added, followed by *t*-butyldimethylsilylchloride (3 mmoles). The progress of the reaction was followed by tlc (eluent CHCl₃, R_f **4** = 0.1, R_f **5** = 0.7). After 48 h, the mixture was partitioned between water (20 ml) and CH₂Cl₂ (20 ml). The organic phase was washed (3 x) with water, dried (MgSO₄), filtered and evaporated. A filtration on silica (eluent CHCl₃) gave the product (Yield, 90%). ¹H NMR (CDCl₃): 7.97 and 7.42 (2 x m, ClBz), 5.46 (m, ClBzOCH), 4.52 (m, ClBzOCH₂CH), 4.34 (m, ClBzOCH₂CH), 4.25 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.61 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.41 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 2.18 [ddd, 1H, ring CH₂ (H exo)], 1.90 [m, ring CH₂ (H endo)], 1.70 and 1.40 [2 x m, CH(CH₂)₇CH₂CH(OSiR₃)-CH₂(OSiR₃) and CHCH₂(CH₂)₇CH(OSiR₃)-CH₂(OSiR₃)], 1.25-1.35 [m, CHCH₂(CH₂)₆CH₂CH(OSiR₃)-CH₂(OSiR₃)], 0.85 [s, C(CH₃)₃], 0.0 [s, Si(CH₃)₂]. ¹³C NMR, 165.3, 139.6, 130.9, 128.7 and 128.3 (ClBz), 80.7 and 79.3 (CHOCH), 77.3 (ClBzOCH), 73.1 (CHOSiR₃), 67.4 (CH₂OSiR₃), 64.7 (ClBzOCH₂), 37.6 (ring CH₂), 35.7, 34.3, 29.6, 29.5, 29.4, 25.9, 25.7 and 25.0 [(CH₂)₈ and C(CH₃)₃], 18.2 [C(CH₃)₃], -4.7 [Si(CH₃)₂]. C, H, analysis, calc. C, 61.86, H, 8.10; found, C, 61.90, H, 8.13.

(2-R,3-S,5-S)-14-O-15-O-di(*t*-butyldimethylsilyl)-1,2,3,5,14,15-hexahydroxy-2,5-anhydropentadecane 6

The diester **5** (2 mmoles) was dissolved in a mixture of THF (50 ml) and 24% aqueous trimethylamine (4 ml). After 3 h at 0°C, the solvent was evaporated. The residue was partitioned between water and ether. The ethereal phase was washed with water (5 x). The *N*-methyl-*p*-chlorobenzamide was precipitated by cooling the ethereal solution at -5°C. Yield of **6**, 90%.

¹H NMR (CDCl₃): 4.17 (m, HOCH), 4.05 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.73 (m, HOCH₂CH), 3.58 (m, HOCH₂CH), 3.55 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.40 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 2.9-3.2 (broad signal, 2 x OH), 1.88 [ddd, 1H, ring CH₂ (H exo)], 1.60 [m, ring CH₂ (H endo)], 1.50 and 1.35 [2 x m, CH(CH₂)₇CH₂CH(OSiR₃)-CH₂(OSiR₃) and CHCH₂(CH₂)₇CH(OSiR₃)-CH₂(OSiR₃)], 1.18-1.28 [m, CHCH₂(CH₂)₆CH₂CH(OSiR₃)-CH₂(OSiR₃)], 0.85 [s, C(CH₃)₃], 0.0 [s, Si(CH₃)₂]. ¹³C NMR, 86.6 and 78.7 (CHOCH), 73.2 (HOCH), 67.4 (CHOSiR₃), 63.2 (CH₂OSiR₃), 60.3 (HOCH₂), 41.1 (ring CH₂), 35.6, 34.3, 29.6, 29.5, 29.4, 26.0, 25.9 and 25.0 [(CH₂)₈ and C(CH₃)₃], 18.2 [C(CH₃)₃], -4.7 [Si(CH₃)₂]. C, H analysis, calc. C, 62.49, H, 11.27; found, C, 62.36, H, 11.37.

(2-R,3-S,5-S)-1-O-(4,4'-dimethoxytrityl)-14-O-15-O-di(*t*-butyldimethylsilyl)-1,2,3,5,14,15-hexahydroxy-2,5-anhydropentadecane 7

Compound **6** (10 mmoles) was dissolved in dry pyridine (50 ml). Dimethoxytrityl chloride (12 mmoles) was added. The reaction was monitored by tlc (eluent CH₂Cl₂/N(C₂H₅)₃, 98:2). When the reaction was complete, methanol was added (10 ml). After 10 minutes, the mixture was diluted with CH₂Cl₂ and washed with aqueous Na₂CO₃ (5%). The organic phase was dried (MgSO₄), filtered and evaporated. The product was isolated by column chromatography (eluent CH₂Cl₂/N(C₂H₅)₃, 98:2, R_f = 0.4). Yield 75%. ¹H NMR (CDCl₃): 6.70-7.45 (trityl), 4.22 (m, HOCH), 4.12 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.88 (m, tritylOCH₂CH), 3.73 (s, CH₃O), 3.61 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 3.49 (m,

tritylOCH₂CH), 3,15 [m, CH(CH₂)₈CH(OSiR₃)-CH₂(OSiR₃)], 2.9-3.1 (broad signal, OH), 1.89 [ddd, 1H, ring CH₂ (H exo)], 1.65 [m, ring CH₂ (H endo)], 1.48 and 1.40 [2 x m, CH(CH₂)₇CH₂CH(OSiR₃)-CH₂(OSiR₃) and CHCH₂(CH₂)₇CH(OSiR₃)-CH₂(OSiR₃)], 1.18-1.32 [m, CHCH₂(CH₂)₆CH₂CH(OSiR₃)-CH₂(OSiR₃)], 0.85 [s, C(CH₃)₃], 0.0 [s, Si(CH₃)₂]. ¹³C NMR, 159.0, 145.3, 136.5, 130.5, 128.6, 128.3, 127.3, 113.7 and 86.5 (trityl), 84.0 and 78.8 (CHOCH), 75.3 (HOCH), 73.2 (tritylO-CH₂), 67.8 (CHOSiR₃), 65.3 (CH₂OSiR₃), 55.7 (CH₃O), 41.0 (ring CH₂), 36.0, 34.1, 29.6, 29.5, 29.4, 26.2, 25.9 and 25.0 [(CH₂)₈ and C(CH₃)₃], 18.2 [C(CH₃)₃], -4.7 [Si(CH₃)₂]. C, H analysis, calc. C, 70.26, H, 9.33; found, C, 70.46, H, 9.37.

{3-[(2-R,3-S,5-S)-1-O-(4,4'-dimethoxytrityl)-14-O-15-O-di(*t*-butyldimethylsilyl)-1,2,5,14,15-pentahydroxy-2,5-anhydrolpentadecoxy}-(2-cyanoethoxy)-(N,N-diisopropylamino)-phosphine 8

The reaction was performed under argon. Compound **7** (0.1 mmoles) was coevaporated three times with acetonitrile, dried under high vacuum (3 h) and dissolved in dry CH₂Cl₂ (5 ml) containing diisopropylammonium tetrazolide (0.06 mmoles). Bis(diisopropylamino) (β-cyanoethoxy) phosphine (0.12 mmoles) was added. The mixture was magnetically stirred for 2 h [control by tlc, hexane/N(CH₂CH₃)₃, 6:1]. Sodium bicarbonate (5%, 5 ml) was added. The organic phase was washed with a phosphate buffer (pH 7, 5 ml), then with brine (sat., 5 ml), dried (MgSO₄), filtered and evaporated. The residue was chromatographed on silica [eluent: hexane/N(CH₂CH₃)₃, 6:1, R_f of **8** = 0.7]. Yield, 70 %. ³¹P NMR, δ = 148.811 and 148.641 ppm. C, H analysis, calc. C, 67.02, H, 9.18; found, C, 67.35, H, 8.98.

Synthesis and purification of oligonucleotide 9

The β-cyanoethyl phosphoramidite strategy was used¹⁸. At the end of the trityl-on synthesis, the glass beads were treated with conc. ammonia at 50°C for 24 h. The basic solution was passed

through an Oligo-PakTM reversed phase cartridge (Pharmacia), that retained the tritylated oligomer. The column was washed with water. Treatment with TFA:H₂O (2%) cleaved the trityl protection and allowed to elute the desired product. The residue after lyophilization (0.02 μ moles of oligonucleotide) was dissolved in a solution of tetrabutylammonium fluoride in THF (0.1 M, 10 μ l). After 2 h of shaking (the time necessary for full desilylation was checked by tlc on the monomeric compound **6**), the THF was evaporated and the residue was dissolved in water (1 ml). This solution was desalted on a Pharmacia NAP-10TM microcolumn. The column was equilibrated with phosphate buffer (15 ml, 10 mM, pH 6.8). The sample was then loaded and eluted with the buffer. The first ml corresponded to the dead volume. The oligonucleotide was found in the following 1.5 ml of eluate.

Oxidation of **9** to **10**

A solution of NaIO₄ (120 mM) in phosphate buffer (20 mM, pH 7) was made. It was degazed by helium (to avoid overoxidation of the aldehyde to be obtained). This solution (1 ml) was added to the glycol-oligonucleotide (100 OD in 25 μ l). The reaction was stopped after 30 min in the dark at room temperature by desalting on NAP-10TM. The success of the reaction was checked by reversed phase HPLC [BIORAD Bio-Sil C₁₈ HL 150x4.6 mm column. Linear gradient of eluent B (triethylammonium acetate 0.1 M in CH₃CN:H₂O, 60:40) in eluent A (triethylammonium acetate 0.1 M in CH₃CN:H₂O, 5:95), from 0 to 30% in 25 min. Flow, 1 ml/min.]. The retention times were 15.4 and 16.3 min. for **9** and **10** respectively. The solution of the aldehyde-oligomer was not kept, but immediately used for the next step.

Titration of **10** by NaBT₄

9 (5 nmoles) was dissolved in water in an eppendorf tube. A 5 mM solution of NaIO₄ in 20 mM acetate buffer (pH 4.75) was added (2.5 μ l). The mixture was stirred in the dark for 5 h. A 10 mM solution of NaBT₄ in 40 mM borate buffer (pH 9) was added

(2.5 μ l). The mixture was stirred at 4°C for 5 h, then diluted with water (300 μ l). The yield of tritium incorporation was determined by passing the sample through NAP-5TM cartridges, isolating the high molecular weight fraction and determining the immobilized tritium content by scintillation counting.

Synthesis of **11**

A solution of biotin hydrazide (10 μ l, 15 mM) and acetate buffer (5 μ l, 500 mM, pH 5) and a solution of sodium cyanoborohydride (5 μ l, 100 mM in acetate buffer) were added to the aldehyde-oligonucleotide **10** dissolved in water (1.5 nmoles in 500 μ l). The mixture was desalted on NAP-10 TM after 2 h at room temperature. The conjugate was isolated by semi-preparative HPLC, in the same conditions as indicated above.

Synthesis of **12**

A solution of D-biotinoyl-1,8-diamino-3,6-dioxaoctane (10 μ l, 15 mM) and phosphate buffer (100 μ l, 20 mM, pH 6.8) and a solution of sodium cyanoborohydride (10 μ l, 5 mM in water) were added to the aldehyde-oligonucleotide **10** dissolved in water (1.5 nmoles in 200 μ l). After 4 h at room temperature, the mixture was diluted by water to 500 μ l and desalted on NAP-5 TM. The conjugate was isolated by semi-preparative HPLC, in the same conditions as indicated above.

Titration of **12** by avidin (Figure 4)

Aliquots of a solution of avidin (Sigma, 5.2 nanomoles/ml of sites of complexation) were added to solutions of the biotin-oligonucleotide conjugate **12** (66 picomoles of oligonucleotide in 200 μ l). Each solution was ultrafiltrated (ULTRAFREE-MCTTM from Millipore, membrane UFC3-LTK, cutoff 30.000), and the filtrate checked by HPLC. Oligomer **9** was used as a not retained internal




FIGURE 4

standard. The addition of 12.5 μ l of the solution of avidine (65 picomoles of sites of complexation) was necessary to completely retain the biotinylated oligomer on the filter.

Acknowledgment: The belgian FNRS, FRIA, FDS and the french ANRS are acknowledged for financial support. A financial support was also obtained from the "Actions de recherches concertées de la Direction générale de l'Enseignement supérieur et de la Recherche scientifique, Ministère de l'Education et de la Recherche, Communauté Française de Belgique".

REFERENCES

- 1 Kawai, S., Maekawajiri, S. and Yamane, A., *Anal. Biochem.*, **209**, 63 (1993).
- 2 Ishii, J.K. and Ghosh, S.S., *Bioconjugate Chem.*, **4**, 35 (1993).
- 3 Arar, K., Monsigny, M. and Mayer, R., *Tetrahedron Lett.*, **34**, 8087 (1993).

- 4 Bongartz, J.P., Aubertin, A.M., Milhaud, P.G. and Lebleu, B., *Nucleic Acids Res.*, **22**, 4681 (1994).
- 5 Lemaitre M., Bayard B. and Lebleu B., *Proc. Natl. Acad. Sci. U.S.A.*, **84**, 648 (1987).
- 6 Boussif, O., Lezoualc'h, F., Zanta, M.A., Mergny, M.D., Scherman, D., Demeneix, B. and Behr, J.P., *Proc. Natl. Acad. Sci. U.S.A.*, **92**, 7297 (1995).
- 7 Wu, G.Y. and Wu, C.H., *J. Biol. Chem.*, **267**, 12436 (1992).
- 8 Bonfils, E., Depierreux, C., Midoux, P., Thuong, N. T., Monsigny, M. and Roche, A. C., *Nucleic Acids Res.*, **20**, 4621 (1992).
- 9 Mattes, W. B., Hartley, J. A. and Kohn, K. W., *Biochim. Biophys. Acta*, **868**, 71 (1986).
- 10 Vasseur, J. J., Rayner, B. and Imbach, J. L., *Biochem. Biophys. Res. Commun.*, **134**, 1204 (1986).
- 11 François, P., Muzzin, P., Dechamps, M. and Sonveaux, E., *New J. Chem.*, **18**, 649 (1994).
- 12 Fox, J.J., Yung, N.C., Wempen, I. and Hoffer, M., *J. Am. Chem. Soc.*, **83**, 4066 (1961).
- 13 Kotick, M.P., Szantay, C. and Bardos, T.J., *J. Org. Chem.*, **34**, 3806 (1969).
- 14 We took this epimer because the aldehyde function is not only suitable for coupling with supports, vectors or reporter groups, but could also be used to perform an hybridization-triggered crosslinking with a target mRNA and so reinforce an antisense effect. For this purpose, the aldehyde function incorporated in a duplex has to point towards the amino functions of the nucleic base situated in front of him. The S epimer was considered to be more suitable than the other because it corresponds to the natural β configuration of the nucleotides.
- 15 Jacobsen, E.N., Marko, I., Mungall, W.S., Schroder, G., Sharpless, K.B., *J. Am. Chem. Soc.*, **110**, 1968 (1988).
- 16 Usman, N., Jiang, M.Y., Cedergren, R.J. and Ogilvie, K.K., *J. Am. Chem. Soc.*, **109**, 7845 (1987).
- 17 Nielsen, J., Taagaard, M., Marugg, J.E., van Boom, J.H. and Dahl, O., *Nucleic Acids Res.*, **14**, 7391 (1986).
- 18 Brown, T. and Brown, D.J.S., in Eckstein, F., ed., *Oligonucleotides and Analogues. A Practical Approach*, IRL Press, Oxford, 1991, 1.