

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

SYNTHESIS OF CHIMERIC OLIGONUCLEOTIDES CONTAINING INTERNUCLEOSIDIC PHOSPHODIESTER AND S-PIVALOYLTHIOETHYL PHOSPHOTRIESTER RESIDUES

Andrei P. Guzaev^a; Balkrishen Bhat^a; Guity Balow^a; Muthiah Manoharan^a

^a Department of Medicinal Chemistry, Isis Pharmaceuticals, Carlsbad, California, U.S.A.

Online publication date: 31 March 2001

To cite this Article Guzaev, Andrei P. , Bhat, Balkrishen , Balow, Guity and Manoharan, Muthiah(2001) 'SYNTHESIS OF CHIMERIC OLIGONUCLEOTIDES CONTAINING INTERNUCLEOSIDIC PHOSPHODIESTER AND S-PIVALOYLTHIOETHYL PHOSPHOTRIESTER RESIDUES', *Nucleosides, Nucleotides and Nucleic Acids*, 20: 4, 1015 — 1018

To link to this Article: DOI: 10.1081/NCN-100002481

URL: <http://dx.doi.org/10.1081/NCN-100002481>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

**SYNTHESIS OF CHIMERIC OLIGONUCLEOTIDES
CONTAINING INTERNUCLEOSIDIC
PHOSPHODIESTER AND
S-PIVALOYLTHIOETHYL PHOSPHOTRIESTER
RESIDUES**

**Andrei P. Guzaev,* Balkrishen Bhat, Guity Balow,
and Muthiah Manoharan**

Department of Medicinal Chemistry, Isis Pharmaceuticals,
2292 Faraday Ave., Carlsbad, California 92008

ABSTRACT

Novel oligonucleotide analogs that bear phosphodiester and bioreversible *S*-pivaloyl 2-mercaptoethyl (SPME) phosphate triester internucleosidic linkages are described. Their synthesis employs a novel methodology of oligonucleotide deprotection under mild, non-aqueous conditions.

Antisense oligonucleotides are a novel class of genomic therapeutics. It is well known however that oligonucleotide phosphorothioates are of limited stability in blood and tissues. Besides, being negatively charged molecules, they lack the ability to permeate biological membranes in an efficient manner. Thus, both their oral bioavailability and cellular uptake need further improvement. In order to address this issue, a bioreversible *S*-acyl 2-mercaptoethyl (1) phosphate protecting group has been introduced into synthetic oligonucleotides. The bioreversible oligonucleotides may potentially serve as prodrug forms of antisense oligonucleotides. A search for their reliable and straightforward synthesis is therefore of great interest.

The preparation of oligonucleotide prodrugs using the *S*-acyl 2-mercaptoethyl strategy presents a complex synthetic problem. So far, no conventional deprotection method has been reported in the literature. By an orthogonal protection

*Corresponding author.




Figure 1. Structures of 2-cyanoethyl phosphoramidites and solid supports.

strategy, preparation of short *S*-pivaloyl 2-mercaptoethyl (SPME) protected oligonucleotides has recently been achieved using photolabile nucleic base protecting groups and solid support (2). However, fully modified SPME oligonucleotides are poorly soluble in water (3) and therefore can hardly be considered as drug candidates.

We reported previously synthesis of a phosphoramidite building block 8 (4) (Fig. 2) and oligothymidylates bearing both SPME and phosphodiester linkages (5). In this communication, we report a novel strategy of oligonucleotide deprotection under mild, non-aqueous conditions. The strategy allows one to deprotect nucleic bases and the backbone while SPME phosphotriester groups are kept intact.

Previously we observed that the SPME group was almost indefinitely stable towards 1 M piperidine in MeCN and of a limited stability towards 0.01 M K₂CO₃ in MeOH (23). With this information in hand, we synthesized a number of model oligonucleotides using phosphoramidites 1–3 and solid supports 6–8 (Fig. 1) and studied their deprotection. We found that phenoxyacetyl (PAC) protection of deoxycytosine and deoxyadenosine residues was removed under conditions compatible with the SPME group. In contrast, removal of the PAC protection from




Figure 2. Synthesis of phosphoramidite building blocks 17–21.



deoxyguanosine residue required an unacceptably long treatment with the solution of piperidine (48 h).

This data prompted us to use *N*-unprotected 2-cyanoethyl deoxyguanosine phosphoramidite **4** synthesized as reported earlier (6). In accordance with the previous observations, the use of **4** within the standard protocol of the DNA synthesis demonstrated no deterioration of coupling yields or the purity of synthetic oligonucleotides (6,7).

Having set the deprotection conditions, we synthesized SPME protected nucleoside phosphoramidites **17–21** and evaluated their use in preparation of chimeric oligonucleotides. First, **10** was converted to a bisamidite **11**, which was isolated on a silica gel column as a colorless oil in 90% yield and more than 98% purity (³¹P NMR). Nucleosides **12–16** were treated with **11** in the presence of **1H**-tetrazole to give **17–21** (Fig. 2). These were isolated by column chromatography in 75–95% yield and characterized. The phosphoramidites **17–21** were stable compounds whose shelf life exceeded 2 years. Previously we reported that the half-life of disappearance of **17** in 80% aqueous MeCN was 58 h, which indicated its high hydrolytic stability (4).

Oligonucleotides **22–27** (Table) were assembled using phosphoramidite building blocks **1–5** and **17–21** and solid supports **6–9**. For deprotection, a three-step procedure was developed. Solid support-bound oligonucleotides were treated first with 1 M piperidine in MeCN for 8 h and then extensively washed with dioxane. On the second step, the synthetic columns were replaced on the instrument, and the standard detritylation subroutine was carried out. Finally, oligonucleotide material was released from the support with 0.01 M K₂CO₃ in MeOH for 80 min, and the solution was neutralized with glacial AcOH. The obtained mixtures consisted of chimeric oligonucleotides **22–24** along with products of methanolysis of the SPME group. With oligothymidylates, the extent of methanolysis was minimal (*ca.* 2% per SPME group). In other cases it depended on the base composition of an oligonucleotide and varied between 2 and 10%.

Table. Chimeric Oligonucleotides **22–27** Containing SPME Phosphotriester Groups^a

Compound	Sequence (5' → 3')	Backbone
22	<i>TTT</i> T ₁₃ <i>TTT</i> T	P=S
23	<i>T*T*T*</i> T ₁₃ <i>T*T*T*</i> T* ^b	P=S
24	CCCCAA <i>T</i> T ₁₀	P=S
25	<i>CC</i> CCCACCACTTCCCCT <i>CT</i> C	P=S
26	<i>CCC</i> CCACCACTTCCCC <i>TCT</i> C	P=S
27	<i>AG</i> CTTCTTTGCACATGT AA A	P=S

^a. Nucleotide residues bearing SPME group at the 3'-phosphate are italicized;

^b. T* stands for 2'-O-(2-methoxyethyl)-5-methyluridine residue.



In conclusion, the present methodology of mild, non-aqueous oligonucleotide deprotection allows straightforward preparation of potential prodrug oligonucleotides bearing a desired number of the SPME phosphotriester moieties and, potentially, other types of base-labile oligonucleotides.

REFERENCES

1. (a) Vives, E.; Dell'Aquila, C.; Bologna, J.-C.; Morvan, F.; Rayner, B.; Imbach, J.-L. *Nucleic Acids Res.* **1999**, *27*, 4071–4076; (b) Tosquellas, G.; Alvarez, K.; Dell'Aquila, K.; Morvan, F.; Vasseur, J.-J.; Imbach, J.-L.; Rayner, B. *Nucleic Acids Res.* **1998**, *26*, 2069–2074.
2. Alvarez, K.; Vasseur, J.-J.; Beltran, T.; Imbach, J.-L. *J. Org. Chem.* **1999**, *64*, 6319–6328.
3. Tosquellas, G.; Bologna, J. C.; Morvan, F.; Rayner, B.; Imbach, J.-L. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2913–2918.
4. Guzaev, A.; Boyode, B.; Balow, G.; Tivel, K. L.; Manoharan, M. *Bioorg. Med. Chem. Lett.*, **1998**, *8*, 1123–1126.
5. Guzaev, A.; Balow, G.; Manoharan, M. *Nucleosides & Nucleotides*, **1999**, *18*, 1391–1392.
6. Hayakawa, Y.; Kataoka, M. *J. Am. Chem. Soc.*, **1998**, *120*, 12395–12401.
7. Gryasnov, S. M.; Letsinger, R. L. *Nucleic Acids Res.*, **1992**, *20*, 1879–1882.



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002481>