

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

SYNTHESIS OF 2',3'-DIDEOXY-2',3'-DIDEHYDRO NUCLEOSIDES VIA A SERENDIPITOUS ROUTE

Z. Guo^a; Y. S. Sanghvi^a; L. E. Brammer Jr.^b; T. Hudlicky^b

^a Isis Pharmaceuticals, Inc., Carlsbad, California, U.S.A. ^b University of Florida, Gainesville, Florida, U.S.A.

Online publication date: 31 March 2001

To cite this Article Guo, Z. , Sanghvi, Y. S. , Brammer Jr., L. E. and Hudlicky, T.(2001) 'SYNTHESIS OF 2',3'-DIDEOXY-2',3'-DIDEHYDRO NUCLEOSIDES VIA A SERENDIPITOUS ROUTE', *Nucleosides, Nucleotides and Nucleic Acids*, 20: 4, 1263 – 1266

To link to this Article: DOI: 10.1081/NCN-100002532

URL: <http://dx.doi.org/10.1081/NCN-100002532>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS OF 2',3'-DIDEOXY-2',3'-DIDEHYDRO NUCLEOSIDES VIA A SERENDIPITOUS ROUTE

Z. Guo,¹ Y. S. Sanghvi,^{1,*} L. E. Brammer, Jr.,² and T. Hudlicky²

¹Isis Pharmaceuticals, Inc., 2292 Faraday Avenue, Carlsbad,
California 92008

²University of Florida, Gainesville, Florida 32604

ABSTRACT

This paper describes a “green” synthesis of 2',3'-unsaturated 2',3'-dideoxynucleosides via an electrochemical reaction. Using this approach d4T, d4U, ddA and ddI can be synthesized in high yields.


INTRODUCTION

With the growing success of clinical programs and potential market demand for oligonucleotide-based drugs, it is important that all raw materials for synthesis of these drugs are easily accessible at affordable cost (1). For the assembly of oligonucleotides, 2'-deoxynucleosides (dNs) are the key building-blocks (2). Currently all dNs are in limited supply and remain expensive due to their isolation from natural sources such as fish milt (3). Therefore, we have focused our attention on methods development for the synthesis of dNs, particularly pyrimidine dNs.

RESULTS AND DISCUSSION

Although thymidine can be synthesized (4) via direct glycosylation procedure (Scheme 1), the concomitant formation of α -anomer decreases the overall

*Corresponding author.




Scheme 1. Synthesis of thymidine via glycosylation procedure.

yield and increases the cost of desired β -anomer. Therefore, we elected to use ribonucleosides as starting material for synthesis of dNs due to their availability in bulk at low cost from bacterial fermentation and preformed glycosidic linkage (Scheme 2).


The Mattocks reaction of uridine analog (**1**) has been reported to provide 2'- α -bromo-3'-acetoxy nucleoside (**2**) due to anchimeric assistance of oxygen at the 2-position of the pyrimidine ring (5). Subsequent reduction of the halide with tin reagents furnished dNs (**3**, Scheme 2). Scale-up of tin mediated reduction is not practical due to the toxic nature of the reagent. In order to implement "green" chemistry in our manufacturing processes, we chose an electrochemical method that offers a versatile and nonstoichiometric alternative to metal-based reduction of the halide.

Treatment of acetyl bromide with uridine and 5-methyluridine in acetonitrile gave quantitative yields of 2'-bromo-3',5'-diacetyluridine (**2**, B = U) and 2'-bromo-3',5'-diacetyl-5-methyluridine (**2**, B = T), respectively. Electrochemical reduction (**6**) of **2** in methanol did not furnish the expected 2'-deoxy nucleoside derivatives **3**. Interestingly, both reactions gave 2',3'-unsaturated 2',3'-dideoxynucleosides **6** via elimination of the 3'-acetoxy group (**5**, Scheme 3). Subsequent treatment with base resulted in formation of d4T and d4U. All attempts to reduce 2'- α -bromo-3'-acetoxy nucleosides **2** via electrochemical methods have resulted in formation of elimination products only.




Scheme 2. Synthesis of thymidine via 2'-deoxygenation procedure.





Scheme 3. Serendipitous synthesis of d4T.



Scheme 4. Green synthesis of ddI.

The concept of electrochemical reduction was further extended towards synthesis of ddI (**9**). The 3'-bromo-3'-deoxy nucleoside **7** was conveniently synthesized from inosine following a literature protocol (5). Electrochemical reduction of **7** in MeOH under similar conditions furnished >80% yield of 2',3'-unsaturated nucleoside **8** (Scheme 4). Transformation of **8** to **9** has been reported in the literature (5).

SUMMARY

Interestingly literature search revealed that electrochemical reductions of nucleosides have been tried (7). However the yields of 2',3'-unsaturated nucleosides were somewhat lower than what we report herein (8). It seems that choice of solvent plays a major role during electrochemical reactions and we found that MeOH gave best results. Furthermore, we believe that this is a first "green" synthesis of d4T and may have potential for further scale-up. In conclusion, we have failed to accomplish our original goal of synthesis of 2'-deoxynucleosides by electrochemical reduction.



ACKNOWLEDGMENTS

We are thankful to Dr. Douglas L. Cole for his support and encouragement throughout this project.

REFERENCES AND NOTES

1. Sanghvi, Y.S., Andrade, M., Deshmukh, R.R., Holmberg, L., Scozzari, A.N. and Cole, D.L. In *Manual of Antisense Methodology*, G. Hartmann and S. Endres (eds.), Kluwer Academic Publishers **1999**, pp 3–23.
2. Sanghvi, Y. S. DNA with Altered Backbones in Antisense Applications In *Comprehensive Natural Products Chemistry*, Editors-in-Chief, D.H.R Barton and K. Nakanishi, in Volume 7: *DNA and Aspects of Molecular Biology* E.T. Kool (ed.), Pergamon Press, 258–311 (1999).
3. The bulk price for 2'-deoxynucleosides isolated from fish milt is around \$1,000/Kg.
4. Freskos, J.N. and Senaratne, P.A. US Patent 4,914,233, April 3, **1990**.
5. Shiragami, H., Amino, Y., Honda, Y., Arai, M., Tanaka, Y., Iwagami, H., Yukawa, T. and Izawa, K. *Nucleosides Nucleotides* **1996**, *15*, 31–45.
6. Cell assembly for electrolysis: The cell was home built using a glass beaker fitted with a compartment (fritted disk at the bottom) containing a platinum plate (2 × 2 cm, 0.1 mm thick) as anode and mercury cathode (pool at the bottom). The top was covered with a plate with an inlet and outlet for argon, a thermometer and an Ag/Ag⁺ (1 mm silver wire immersed in 0.1 M AgNO₃ in acetonitrile) reference electrode. Cyclic and linear sweep voltammograms were conducted on the electrolysis solutions prior to the electrolysis to determine the optimum reduction potentials. Electrochemical reductions were performed in methanol or acetonitrile with tetraethylammonium *p*-toluenesulfonate (0.2 M) as electrolyte. The concentration of the electrolyte was maintained in a 5-fold excess relative to the electroactive species. The nucleosidic substrate (5 mmol) was added and the stirred solution was electrolysed under argon at an initial constant potential of -1.9 V for 1 h. The progress of the reaction was followed by TLC and most of the starting material was consumed within 1 h. Upon completion of the reaction, the solution was concentrated under vacuum and the residue purified by short silica gel column chromatography. The yield of the isolated product was always >80%. The deprotected (NH₄OH) products were identical (TLC, UV, NMR) with an authentic sample.
7. Adachi, T., Iwaski, T., Inoue, I. and Miyoshi, M. *J. Org. Chem.* **1979**, *44*, 1404–1409.
8. Johansen, O. Marcuccio, S.M. and Mau, A.W.-H. *Aust. J. Chem.* **1994**, *47*, 1843–1850.



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002532>