

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Recent Advances in the Synthesis of Conformationally Locked Nucleosides and Their Success in Probing the Critical Question of Conformational Preferences by Their Biological Targets

Yongseok Choi^a; Hyung R. Moon^a; Yuichi Yoshimura^a; Victor E. Marquez^{ab}

^a Laboratory of Medicinal Chemistry, Center for Cancer Research, National Cancer Institute at Frederick, Frederick, Maryland, USA ^b Laboratory of Medicinal Chemistry, Center for Cancer Research, National Cancer Institute at Frederick, Frederick, MD, USA

Online publication date: 09 August 2003

To cite this Article Choi, Yongseok , Moon, Hyung R. , Yoshimura, Yuichi and Marquez, Victor E.(2003) 'Recent Advances in the Synthesis of Conformationally Locked Nucleosides and Their Success in Probing the Critical Question of Conformational Preferences by Their Biological Targets', *Nucleosides, Nucleotides and Nucleic Acids*, 22: 5, 547 — 557

To link to this Article: DOI: 10.1081/NCN-120021954

URL: <http://dx.doi.org/10.1081/NCN-120021954>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Recent Advances in the Synthesis of Conformationally Locked Nucleosides and Their Success in Probing the Critical Question of Conformational Preferences by Their Biological Targets

Yongseok Choi, Hyung R. Moon, Yuichi Yoshimura,
and Victor E. Marquez*

Laboratory of Medicinal Chemistry, Center for Cancer Research, National
Cancer Institute at Frederick, NIH, HHS, Frederick, Maryland, USA

ABSTRACT

The present work describes some recent approaches to the syntheses of three classes of locked-North nucleosides: β -D-ribo-, β -D-deoxyribo-, and β -D-dideoxyribonucleosides. The method developed for the latter class permitted access to a novel bicyclo[3.1.0]hexene-type nucleosides structurally similar to D4T and carbovir. A structural analysis and biological activities are discussed.

Key Words: Bicyclo[3.1.0]hexane and bicyclo[3.1.0]hexene; Conformationally locked carbocyclic nucleosides; Pseudorotational analysis; D4T and carbovir analogues; Intramolecular cyclopropanation; Lipase-catalyzed resolution.

*Correspondence: Victor E. Marquez, Laboratory of Medicinal Chemistry, Center for Cancer Research, National Cancer Institute at Frederick, NIH, HHS, 376 Boyles St., Frederick, MD 21702, USA; E-mail: marquez@dc37.nci.nih.gov.






Figure 1. Fixed location of (N)- and (S)-methanocarba nucleosides in the pseudorotational cycle (nucleoside numbering).

INTRODUCTION


Conventional nucleosides or nucleotides equilibrate rapidly in solution between two extreme forms of ring pucker: (1) The North conformation with pseudorotational P values ranging between 342° to 18° (${}_2E \rightarrow {}^3T_2 \rightarrow {}^3E$) and (2) the antipodal South conformation with values of P between 162° to 198° (${}^2E \rightarrow {}^2T_3 \rightarrow {}_3E$) (Fig. 1).^[1,2] Preference for any of these specific conformations in solution is determined by the interplay of important interactions such as anomeric and *gauche* effects.^[2] However, when a nucleoside or nucleotide binds to its target enzyme, only one form is present at the active site. In order to query these enzymes for their conformational preference we have designed and synthesized conformationally locked nucleosides that reside strictly in the normal range of either North or South conformations.

The bicyclo[3.1.0]hexane scaffold provides a convenient way to lock the conformation of carbocyclic nucleosides which are known generically as a methanocarba (MC) nucleosides. Because of its exclusive pseudoboat conformation, a rigid North envelope (${}_2E$) conformation can be constructed when the cyclopropane ring appears fused between carbons $C_{4'}$ and $C_{5'}$. Conversely, fusion of the cyclopropane ring between carbons $C_{1'}$ and $C_{5'}$ provides a rigid South (${}_3E$) envelope conformation (Fig. 1).

RESULTS AND DISCUSSION


The North Scaffold as a Promising Template for Drug Development

North- and South-MC nucleosides with the same nucleobase are very nicely discriminated by specific enzymes and thus function as target-selective agents. Some of the most important compounds with potential as drug candidates are in the North hemisphere. (N)-MC thymidine (**1**) is a very selective antiviral agent which has shown greater potency against HSV-1 and HSV-2 infection and less toxicity than the standard ganciclovir.^[3,4] This compound is also effective in vivo against murine MC38 tumors transduced with the herpes thymidine kinase gene.^[5] MRS2279 (**2**), a 3',5'-biphosphate purine analogue from Dr. Jacobson's laboratory is a very potent and selective high affinity P2Y₁ receptor agonist which does not block nucleotide signaling at most of the other known P2Y receptor subtypes.^[6]




SYNTHETIC METHODS

Most of the published methods for (N)-MC nucleosides rely on a critical hydroxyl-directed cyclopropanation step (Sch. 1) from the allylic position on a cyclopentenol precursor.^[7-9] Although the cyclopropanation step works well, the synthesis of the starting carbocycle is quite laborious and requires expensive starting materials. The resulting pseudosugars can be converted to purine or pyrimidine (N)-MC nucleosides by convergent approaches (from **4**) or by linear approaches (from **6**).



Scheme 1.





Scheme 2.


Normally, a convergent Mitsunobu coupling with the preformed nucleobase works better with purines. On the other hand, for pyrimidines, the formation of abundant *O*-alkylation products makes the linear approach the preferred route.

In the past couple of years we have developed some new approaches using more accessible and less expensive starting materials. At the core of these approaches lies a critical intramolecular cyclopropanation step from a carbene intermediate engendered from a diazo β -keto ester. The retrosynthetic analysis depicted in Sch. 2 illustrates the general strategy.

Steps Ia to VIIa describe the approach to ribo-like, (N)-MC nucleosides, whereas steps Ib to VIIb correspond to the deoxyribo series. The syntheses of the precursor diazo β -keto esters IIIa and IIIb are simple and straightforward, and after metal-catalyzed thermolysis the carbene intermediates IVa and IVb fold intramolecularly into the desired bicyclo[3.1.0]hexane scaffold. The top part in Sch. 2 depicts the desired β -D-like configuration, achieved via a stereospecific synthesis from a chiral precursor (Ia). For the bottom part, the process leading to intermediate VIIb is achiral, and thus the racemic mixture is resolved at this stage via a lipase-catalyzed asymmetric acetylation.

A. Enantiospecific Synthesis of Ribo-type, North-Methanocarba Nucleosides

Starting with inexpensive and plentiful D-isoascorbic acid, the chiral precursor **7**^[10,11] was obtained in 4 steps and 60% overall yield (Sch. 3). After the stepwise oxidation of **7**, first under Swern conditions and then with sodium chlorite, Dieckmann condensation of the activated acid **8** with ethyl 2-lithioacetate afforded β -keto esters **9** (8%) and **10** (59%) as a mixture of separable diastereoisomers. Following diazo transfer with tosyl azide and triethylamine (TEA), β -keto ester **10** was converted to diazo compound **11**, which underwent metal-catalyzed thermolysis to generate the desired bicyclo[3.1.0]hexane scaffold via a carbenoid intermediate (IVa, Sch. 2). Comparisons with authentic compounds previously made from different sources,^[12] confirmed that the major isomer, **12**, had the desired stereochemistry. Reduction of the carbonyl moiety gave diol **14**, which can be protected. In the case of benzyl-protected **15**, acid-catalyzed removal of the acetonide and reaction of **16** with thionyl chloride produced the cyclic sulfite **17**. This compound reacted well with sodium azide to give 82% of **18** and 10% of the alternative regioisomer. Azide **18** was efficiently reduced to amine **19** (overall yield from D-isoascorbic acid = 2%) which is a direct precursor of purine and pyrimidine analogues using well-known linear approaches.^[12]




Scheme 3.



B. Chemico-Enzymatic Approaches to Deoxyribo-type, North-Methanocarba Nucleosides

In this case the process is basically achiral ($\text{Ib} \rightarrow \text{VIIb}$, Sch. 2). A final resolution step using adenosine deaminase worked well only for purines, particularly for the guanosine analogue, which was obtained directly from the deamination of the 2,6-diaminopurine nucleoside.^[13] To overcome this limitation, a resolution step prior to the incorporation of the nucleobase was recently accomplished via an economical lipase-catalyzed asymmetric acetylation performed on diol **VIb** (Sch. 2).^[14]

Commercially available lipase PS-C in the presence of a large excess of vinyl acetate effectively discriminated between the enantiomers of *rac*-**20** (**VIb**, $\text{R} = \text{SiPh}_2\text{CMe}_3$, Schs. 2 and 4) producing diacetate **21** (57%) and monoacetate **22** (42%) after 64 h. Using chiral HPLC, the enantiomeric excess of monoacetate **22** was estimated to be 96% and, although the enantiomeric excess of **21** should have been over 98%, it contained 13% of an impurity (**21'**). Removal of this impurity was important since diacetate **21** corresponded to the enantiomer with the desired D-like configuration.^[14] The structure of **21'** was that of an acetal arising from the reaction of the unrecognized enantiomer **22** with excess acetaldehyde—generated during the reaction—followed by further enzymatic acetylation. Treatment of contaminated **21** with ammonium hydrogenfluoride in aqueous DMF (50°C, overnight) gave an easily separable mixture of **23** and diol **12** in 79% and 11% yields, respectively.




Scheme 4.

tively. The optical purity of **23**, was estimated to be 99% ee by HPLC after its conversion back to **21** with *tert*-butyldiphenylsilyl chloride.^[14] Completing the synthesis of deoxyribo-type, (N)-MC nucleosides from diacetate **23** was simple and straightforward.

C. Enantiospecific Synthesis of Dideoxyribo-type and Dideoxy-didehydroribo-type, North-Methanocarpa Nucleosides. Synthesis of Analogues of D4T and Carbovir


The attempt to remove the secondary, C_{3'}-OH (nucleoside numbering) under radical deoxygenation conditions resulted in the destruction of the fused cyclopropane ring, which collapsed after formation of a radical intermediate at the adjacent carbon. To circumvent this problem, deoxygenation was performed prior to the formation of the bicyclo[3.1.0]hexane template (Sch. 5) starting from a compound readily available in our laboratory, such as **24**.^[8] The corresponding thiocarbonylimidazolidine (**25**) gave under radical deoxygenation conditions an inseparable mixture of cyclic allylic (**26**) and homoallylic (**27**) benzoate esters. The mixture of alcohols (**28** and **29**) obtained after hydrolysis of the benzoate esters was cyclopropanated to give isomeric bicyclo[3.1.0]hexanols, **30** and **31**, that were easily separable by column chromatography. With the exception of optical rotation, the spectral properties of **30** matched exactly those of its previously reported racemate (*rac*-**30**).^[15] From compound **30**, purine nucleosides are readily accessible in a convergent manner by Mitsunobu coupling with the corresponding nucleobase. However, coupling of *rac*-**30** with pyrimidines, such as *N*³-benzoylthymine, was reported to give rather low yields of the desired *N*-alkylated product.^[15]

To improve the coupling yields for pyrimidine nucleosides we postulated that a bicyclo[3.1.0]hexenol, such as **35**, would react more readily (Sch. 6). The synthesis of this precursor required the elimination of the C_{3'}-OH (nucleoside sugar numbering) from a compound, such as **33**, which was easily obtained from **32**^[9] in 3 steps.



Scheme 5.





Scheme 6.

When the secondary alcohol was converted to the mesylate ester **34** for a possible base-catalyzed E2 elimination, only the hydrolyzed product was obtained. Because of the constrained nature of the bicyclo[3.1.0]hexane template, a more favorable *anti*-elimination was sought by attempting to invert the configuration of the leaving group after treatment with $\text{I}_2/\text{Ph}_3\text{P}/\text{imidazole}$. However, the constrained nature of the system forced a double inversion to give instead the thermodynamic product **36** with the iodine axially disposed. Still, elimination occurred by refluxing **36** in toluene in the presence of DBU to produce the desired bicyclo[3.1.0]hexene pseudosugar **37** in 51% yield. Base-catalyzed hydrolysis of the benzoate ester afforded the key intermediate **35**. As predicted, coupling of **35** with N^3 -benzoylthymine under Mitsunobu conditions gave the corresponding *N*-alkylated product as the major isomer in 43% yield. The final D4T analogue **38** was obtained after treatment with NH_4OH and final removal of the TBDPS group. A similar approach was used for the synthesis of the carbovir analogue **39**. Catalytic hydrogenation of **38** and **39** afforded the corresponding dideoxyribo-type analogues in excellent yields.

STRUCTURAL ANALYSIS AND STRUCTURE-ACTIVITY RELATIONSHIP

The use of bicyclo[3.1.0]hexane templates has already allowed us to determine conformational preference for enzymes such as adenosine deaminase,^[9] HIV reverse

Table 1. Pseudorotational parameters for **38**, D4T (**40**), and Carbovir (**41^a** and **41^b**). Numbers represent degrees (°).

	ν_0	ν_1	ν_2	ν_3	ν_4	P	ν_{\max}	χ	γ
38	6.56	-5.83	2.67	1.68	-5.20	293.14	6.81	-119.12	56.17
40	-0.51	0.20	0.19	-0.49	0.60	72.08	0.61	-100.82	52.80
41^a	-26.47	17.89	-1.09	-16.02	25.68	92.23	27.98	-86.24	-177.76
41^b	-27.22	17.21	0.23	-17.56	26.98	89.55	28.91	-52.63	-82.90

transcriptase,^[16] DNA (cytosine-C5) methyl transferase,^[17] and several subtypes of adenosine receptors.^[7] A common structural characteristic of D4T (**40**) and carbovir (**41**) is a double bond, a feature that imparts a certain level of rigidity to the five member ring. Thus, compounds **38** and **39** represent hybrid molecules that incorporate in their bicyclo[3.1.0]hexene template a similar double bond. Compound **38** provided adequate crystals for X-ray analysis. At first glance, a comparison between the X-ray structure of **38** and that of D4T (**40**) obtained from the Cambridge database revealed little differences. Indeed, the superposition of both structures shows a RMS deviation of only 0.039 Å. On the other hand, the pseudorotational parameters for these molecules are quite different, particularly with respect to the value of P (Table 1). Both structures are in the North hemisphere, but they are 140° apart from each other, separated from a perfect North ($P=0^\circ$) by an almost equal number of degrees (ca. 70°) towards the West and East, respectively. Also, the ring of D4T (**40**) is far more planar ($\nu_{\max}=0.61^\circ$) with a mean deviation from planarity of only 0.0025 Å. The ring of **38** is more puckered ($\nu_{\max}=6.81^\circ$) with a mean deviation from planarity of 0.025 Å.

Although we did not obtain a crystal structure of **39**, the rigid bicyclo[3.1.0]-hexene ring is expected to be similar to that of **38** with identical pseudorotational parameters. In contrast, the pseudorotational parameters for the two molecules present in the unit cell of a carbovir X-ray structure are quite different from those of **39** (Table 1). The main differences found are first the degree of flatness (see ν_{\max}) which shows that the five-member ring of **38** (and by analogy that of **39**) is much flatter than that of carbovir. Secondly, the value of P places carbovir in an almost perfect East conformation, 157° away from **38** or **39**. These conformational differences may be associated with some unique biological activity for compounds **38** and **39**, which is currently under investigation.

REFERENCES

1. Saenger, W. Chapter 2, Defining terms for the nucleic acids. In *Principles in Nucleic Acid Structure*; Springer-Verlag: New York, NY, 1984.
2. Thibaudeau, C.; Chattopadhyaya, J. Stereoelectronic effects in nucleoside and nucleotides. In *Stereoelectronic Effects in Nucleosides and Nucleotides and Their Structural Implications*; Uppsala University Press: Uppsala, Sweden, 1999.
3. Marquez, V.E.; Siddiqui, M.A.; Ezzitouni, A.; Russ, P.; Wang, J.; Wagner, R.W.; Matteucci, M. Nucleosides with a twist. Can fixed forms of sugar ring



- pucker influence biological activity in nucleosides and oligonucleotides? *J. Med. Chem.* **1996**, *39*, 3739–3747.
4. Zalah, L.; Huleihel, M.; Manor, E.; Konson, A.; Ford, H., Jr.; Marquez, V.E.; Johns, D.G.; Agbaria, R. Metabolic pathways of N-methanocarbathymidine, a novel antiviral agent in native and herpes simplex virus-infected vero cells. *Antiviral Res.* **2002**, *55*, 63–75.
 5. Noy, R.; Ben-Zvi, Z.; Manor, E.; Candotti, F.; Morris, J.C.; Ford, H., Jr.; Marquez, V.E.; Johns, D.G.; Agbaria, R. Antitumor activity and metabolic activation of (N)-methanocarbathymidine, a novel thymidine analog with a pseudosugar moiety rigidly fixed in the northern conformation, in murine colon cancer cells expressing herpes simplex virus thymidine kinase. *Mol. Cancer Therap.* **2002**, *1*, 585–593.
 6. Boyer, J.L.; Adams, M.; Ravi, R.G.; Jacobson, K.A.; Harden, T.K. 2-Chloro N6-methyl-(N)-methanocarpa-2'-deoxyadenosine-3',5'-bisphosphate is a selective high affinity P2Y1 receptor antagonist. *Brit. J. Pharmacol.* **2002**, *135*, 2004–2010.
 7. Siddiqui, M.A.; Ford, H., Jr.; George, C.; Marquez, V.E. Synthesis, conformational analysis, and biological activity of a rigid carbocyclic analogue of 2'-deoxyaristeromycin built on a bicyclo[3.1.0]hexane template. *Nucleosides Nucleotides* **1996**, *15*, 235–250.
 8. Ezzitouni, A.; Russ, P.; Marquez, V.E. (1S,2R)-[(Benzyloxy)methyl]cyclopentenol. A versatile synthon for the preparation of 4',1'a-methano and 1',1'a-methanocarbocyclic nucleosides. *J. Org. Chem.* **1997**, *62*, 4870–4873.
 9. Marquez, V.E.; Russ, P.; Alonso, R.; Siddiqui, M.A.; Hernandez, S.; George, C.; Nicklaus, M.C.; Dai, F.; Ford, H., Jr. Synthesis of conformationally restricted carbocyclic nucleosides. The role of the O4'-oxygen in the key hydration step of adenosine deaminase. *Helv. Chim. Acta* **1999**, *82*, 2119–2129.
 10. Cohen, N.; Banner, B.L.; Laurenzano, A.J.; Carozza, L. 2,3-O-Isopropylidene-D-erythrionolactone. *Org. Syn.* **1985**, *63*, 127–135.
 11. Rao, B.V.; Lahiri, S. Stereoselective transformations leading to pentono-1,4-lactones. *J. Carbohydr. Chem.* **1996**, *15*, 975.
 12. Moon, H.R.; Kim, H.O.; Chun, M.W.; Jeong, L.S.; Marquez, V.E. Synthesis of cyclopropyl-fused carbocyclic nucleosides via the regioselective opening of cyclic sulfites. *J. Org. Chem.* **1999**, *64*, 4733–4741.
 13. Moon, H.R.; Ford, H., Jr.; Marquez, V.E. A remarkably simple chemico enzymatic approach to structurally complex bicyclo[3.1.0]hexane carbocyclic nucleosides. *Org. Lett.* **2000**, *2*, 3793–3796.
 14. Yoshimura, Y.; Moon, H.R.; Choi, Y.S.; Marquez, V.E. Enantioselective synthesis of bicyclo [3.1.0]hexane carbocyclic nucleosides via a lipase-catalyzed asymmetric acetylation. Characterization of an unusual acetal byproduct. *J. Org. Chem.* (in press).
 15. Rodriguez, J.B.; Marquez, V.E.; Nicklaus, M.C.; Mitsuya, H.; Barchi, J.J. Conformationally locked nucleoside analogues. Synthesis of dideoxycarbocyclic nucleoside analogues structurally related to neplanocin C. *J. Med. Chem.* **1994**, *37*, 3389–3399.
 16. Marquez, V.E.; Ezzitouni, A.; Russ, P.; Siddiqui, M.A.; Ford, H., Jr.; Feldman, R.J.; Mitsuya, H.; George, C.; Barchi, J.J., Jr. HIV-1 reverse transcriptase



- can discriminate between two conformationally locked carbocyclic AZT-triphosphate analogues. *J. Amer. Chem. Soc.* **1998**, *120*, 2780–2789.
17. Wang, P.; Nicklaus, M.C.; Marquez, V.E.; Brank, A.S.; Christman, J.K.; Banavali, N.K.; MacKerell, A.D., Jr. Use of oligodeoxyribonucleotides with conformationally constrained abasic sugar targets to probe the mechanism of base flipping by HhaI DNA (cytosine C5)-methyltransferase. *J. Amer. Chem. Soc.* **2000**, *122*, 12,422–12,434.



