

This article was downloaded by:

On: 25 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Toward a Catalytic Site in DNA: Polyaza Crown Ether as Non-Nucleosidic Building Blocks in DNA Conjugates

Ulla Jakobsen^a; Katja Rohr^a; Stefan Vogel^a

^a Nucleic Acid Center, Department of Chemistry, University of Southern Denmark, Odense M, Denmark

To cite this Article Jakobsen, Ulla , Rohr, Katja and Vogel, Stefan(2007) 'Toward a Catalytic Site in DNA: Polyaza Crown Ether as Non-Nucleosidic Building Blocks in DNA Conjugates', *Nucleosides, Nucleotides and Nucleic Acids*, 26: 10, 1419 – 1422

To link to this Article: DOI: 10.1080/15257770701539260

URL: <http://dx.doi.org/10.1080/15257770701539260>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

TOWARD A CATALYTIC SITE IN DNA: POLYAZA CROWN ETHER AS NON-NUCLEOSIDIC BUILDING BLOCKS IN DNA CONJUGATES

Ulla Jakobsen, Katja Rohr, and Stefan Vogel □ *Nucleic Acid Center, Department of Chemistry, University of Southern Denmark, Odense M, Denmark*

□ *A number of functionalized polyaza crown ether building blocks have been incorporated into DNA-conjugates as catalytic Cu²⁺ binding sites. The effect of the DNA-conjugate catalyst on the stereochemical outcome of a Cu²⁺-catalyzed Diels-Alder reaction will be presented.*

Keywords Polyaza crown ether; catalysis; DNA-conjugates

INTRODUCTION


Due to its architecture, DNA is an ideal scaffold for the construction of hybrid materials including catalytic systems (e.g., DNAzymes).^[1–7] Feringa et al. recently introduced a new modular assembled DNA-based catalyst.^[11] The catalytic Cu²⁺ binding site intercalates into dsDNA due to their polyaromatic nature. The Cu²⁺ binding site is thereby brought into proximity of the chiral environment of the dsDNA, which in turn leads to transfer of chirality from the dsDNA to the reaction products with an *ee* of up to 99% for the major (*endo*) isomer. We report in this communication another approach based on covalent incorporation of catalytic polyaza crown ether based Cu²⁺ binding sites into ssDNA.^[12]

RESULTS AND DISCUSSION

The synthetic route toward the desired building blocks (monomer **W**, **X**, and **Y**) is very flexible and can be easily adapted for a series of macrocyclic building blocks for automated DNA synthesis based on the phosphoramidite approach (Scheme 1).^[8,12] Macrocyclic **8** served as universal building block for the *N*-functionalization by acylation (**W**) or reductive amination (**X**, **Y**).

Funding of this work was provided by The Nucleic Acid Center (NAC).

Address correspondence to Stefan Vogel, Nucleic Acid Center, Department of Chemistry, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark. E-mail: snv@chem.sdu.dk



SCHEME 1 Synthesis of polyaza crown ether phosphoramidite building blocks and monomer substitution patterns: **W**: R = H, **X**: R = C₁₆H₃₃ (palmityl), **Y**: R = CH₂C₅H₄N (2-pyridyl).

Synthesis of ONs (oligonucleotides) was performed in 0.2 μ mol scale on an automated DNA synthesizer using the phosphoramidite approach. Incorporation of monomers **W**, **X**, and **Y** opposite to a complementary 17-mer ss-DNA sequence resulted in a destabilization (-10–12°C, Table 1, entries 2–4). This can be explained by the steric distortion of the corresponding dsDNA (**W**, **X**, and **Y** are $\sim 2 \times$ larger than A, G, C, and T). Juxtapositioned incorporations of **W**, **X**, and **Y** are well tolerated (+8°C, for **X**, Table 1, entry 5) despite the steric demand and non-nucleosidic structure of the monomers. The remarkable increase in thermal stability displayed by modification **X** is likely due to undisturbed interstrand base distances in the duplex and additional strong hydrophobic forces (C₁₆-palmityl chains). The chosen reaction (Scheme 2) has been used for comparison with the system reported by Feringa et al.^[11]

The 2,6-diamino-pyridine unit in the core of **8** is binding Cu²⁺ efficiently accompanied by the appearance of a characteristic UV band at 668 nm. UV-titration with increasing conc. of Cu²⁺ revealed a 1:1 stoichiometry of the complex.

The initially performed catalytic reactions have shown that all DNA-conjugates are able to catalyse the Diels-Alder reaction but only one of the DNA-conjugates displayed a very weak asymmetric induction (10% *ee*,

TABLE 1 Influence of monomer W-Y on thermal DNA duplex stability

Entry	Sequence	T_m [°C] ^a
1	5'-TGT-GGA-AGA-AGT-TGG-TG 3'-ACA-CCT-TCT-TCA-ACC-AC	56.0
2	5'-TGT-GGA-AGA-AGT-TGG-TG 3'-ACA-CCT-TC <u>W</u> -TCA-ACC-AC	44.0
3	5'-TGT-GGA-AGA-AGT-TGG-TG 3'-ACA-CCT-TC <u>X</u> -TCA-ACC-AC	44.5
4	5'-TGT-GGA-AGA-AGT-TGG-TG 3'-ACA-CCT-TC <u>Y</u> -TCA-ACC-AC	45.5
5	5'-TGT-GGA-AG <u>X</u> -AGT-TGG-TG 3'-ACA-CCT-TC <u>X</u> -TCA-ACC-AC	64.0

^a 10 mM sodium phosphate, 100 mM NaCl, 0.1 mM EDTA, adjusted to pH 7.0, concentration of 1 μ M for each DNA strand.


**SCHEME 2** Cu^{2+} catalyzed Diels-Alder reaction: DNA-conjugates **W**, **X**, and **Y** were used as catalysts.

Figure 1) under the conditions used by Feringa et al. (measured values of 9 and 11% have been averaged to 10%). We have not investigated the influence of the juxtapositioned base (e.g., T, G, C, instead of A) due to prohibitive amounts of ONs needed for further experiments. From our initial results and the putative structure of the catalytic center we assume that the reaction center is not close enough to the chiral environment to induce a stronger asymmetric induction, the use of a considerably smaller catalytic center is therefore the next step towards improved asymmetric induction.

A flexible synthetic strategy toward functionalized polyaza crown ether amidites as well as an efficient incorporation of the respective building




FIGURE 1 HPLC traces for a) racemic mixture, b) reaction catalyzed with dsDNA (Table 1, entry 3) with 10% *ee* (no effect on *ee* for **W** and **Y**), c) schematic model of the catalytic center embedded in a 17-mer dsDNA.

blocks into DNA sequences has been achieved. The results from the catalyzed Diels-Alder reactions have shown that a more rigid catalytic center in close proximity to the helical DNA regions is required to achieve significant asymmetric induction.

REFERENCES

1. Storhoff, J.J.; Mirkin, C.A. *Chem. Rev.* **1999**, 99, 1849–1862.
2. Seeman, N.C. *Trends in Biotechnology* **1999**, 17, 437–443.
3. Braun, E.; Eichen, Y.; Sivan, U.; Ben-Yoseph, G. *Nature* **1998**, 391, 775–778.
4. Yan, H.; Park, S.H.; Finkelstein, G.; Reif, J.H.; LaBean, T. H. *Science* **2003**, 301, 1882–1884.
5. Mirkin, C.A.; Letsinger, R.L.; Mucic, R.C.; Storhoff, J.J. *Nature* **1996**, 382, 607–609.
6. Mao, C.; LaBean, T.H.; Reif, J.H.; Seeman, N.C. *Nature* **2000**, 391, 493–496.
7. Seeman, N.C. *Trends Biochem. Sci.* **2005**, 30, 119–125.
8. Vogel, S.; Rohr, K.; Dahl, O.; Wengel, J. *Chem. Commun.* **2003**, 8, 1006–1007.
9. Li, X.; Liu, D.R. *Ang. Chem. Int. Ed.* **2004**, 43, 4848–4870.
10. Brunner, J.; Mokhir, A.; Kraemer, R. *J. Am. Chem. Soc.* **2003**, 125, 12410–12411.
11. Roelfs, G.; Feringa, B.L. *Ang. Chem. Int. Ed.* **2005**, 44, 3230–3232.
12. Rohr, K.; Vogel, S. *Chembiochem* **2006**, 7, 463–470.