

This article was downloaded by:

On: 25 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Fluorinated 7-Deazapurine 2'-Deoxyribonucleosides: Modification At The Nucleobase And The Sugar Moiety

Frank Seela^{ab}; Kuiying Xu^{ab}; Padmaja Chittepu^{ab}; Xin Ming^{ab}

^a Laboratory of Bioorganic Chemistry and Chemical Biology, Center for Nanotechnology, Münster, Germany ^b Laboratorium für Organische und Bioorganische Chemie, Universität Osnabrück, Osnabrück, Germany

To cite this Article Seela, Frank , Xu, Kuiying , Chittepu, Padmaja and Ming, Xin(2007) 'Fluorinated 7-Deazapurine 2'-Deoxyribonucleosides: Modification At The Nucleobase And The Sugar Moiety', *Nucleosides, Nucleotides and Nucleic Acids*, 26: 6, 607 – 610

To link to this Article: DOI: 10.1080/15257770701490357

URL: <http://dx.doi.org/10.1080/15257770701490357>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

FLUORINATED 7-DEAZAPURINE 2'-DEOXYRIBONUCLEOSIDES: MODIFICATION AT THE NUCLEOBASE AND THE SUGAR MOIETY

Frank Seela, Kuiying Xu, Padmaja Chittepu, and Xin Ming □ *Laboratory of Bioorganic Chemistry and Chemical Biology, Center for Nanotechnology, Münster, Germany, and Laboratorium für Organische und Bioorganische Chemie, Universität Osnabrück, Osnabrück, Germany*

□ *7-Deaza-7-fluoro-purine 2'-deoxynucleosides as well as 2'-deoxy-2'-fluoroarabinofuranosyl nucleosides 1–8 were synthesized. The fluorine atom was introduced on the base level with Selectfluor. Nucleobase-anion glycosylation was then employed to form the nucleosides. Properties of the fluorine compounds were studied in solution and in solid state. Compound 4a was incorporated into oligonucleotides where the stabilizing effect was observed.*


Keywords fluorination; pyrrolo[2,3-*d*]pyrimidine; glycosylation; sugar conformation; base pairing

INTRODUCTION

Replacement of hydrogen atoms or functional groups in an organic compound by fluorine atoms may cause a dramatic change in physical properties and biological activities, which is one of the reasons for the interest in the synthesis of fluoro compounds. Our group has expended effort to introduce the fluorine atom to nucleosides. We are interested in combining the favorable properties of pyrrolo[2,3-*d*]pyrimidine nucleosides with the well documented pharmaceutical behavior of fluorine substitution. Apart from the properties of monomeric nucleosides, 7-halogenated 7-deazapurine nucleosides can stabilize the DNA duplexes.^[1–4] Herein, we report on the synthesis of nucleosides **1–3** and **6–8**, as well as the phosphoramidite **9** and **10**. The nucleosides **4a**, **b** and **5** were described recently^[5] (Scheme 1). Oligonucleotides containing compound **4a** were prepared and their stability in DNA duplexes was determined.

We gratefully acknowledge financial support by Idenix Pharmaceuticals, Cambridge, USA.

Address correspondence to Frank Seela, Laboratory of Bioorganic Chemistry and Chemical Biology, Center for Nanotechnology, Heisenbergstraße 11, 48149 Münster, Germany, and Laboratorium für Organische und Bioorganische Chemie, Universität Osnabrück, Barbarastraße 7, 49069 Osnabrück, Germany. E-mail: frank.seela@uni-osnabrueck.de




SCHEME 1

RESULTS AND DISCUSSION


Earlier, a 7-fluoro residue was introduced in the nucleobase **11** employing Selectfluor.^[6] The same strategy was used to synthesize the fluorinated pyrrolo[2,3-*d*]pyrimidine derivatives **12** and **13** by utilizing lower temperature conditions than used for **11**. The nucleobase-anion glycosylation reaction^[7,8] of the base (**11-13**) afforded the 7-fluorinated inosine derivatives **4a,b**, as well as the 7-deaza-7-fluoro-2'-deoxyguanosine (**5**), 7-deaza-7-fluoro-2'-deoxyisoguanosine (**7**), and 7-deaza-7-fluoro-2'-deoxyxanthosine (**8**). Glycosylation of **11** with the halogenose **14a,b** gave the nucleosides **15a,b**, which were deprotected in NaOCH₃ to afford the 6-methoxy derivatives **1a,b**. Compounds **4a,b** were obtained by heating **1a,b** in aqueous NaOH (Scheme 2). Conformational analysis of the nucleosides **4a,b** showed that the fluorine substitution in the 2'-arabino position biased the conformation toward *N* (from 30% *N* for **4a** to 40% *N* for **4b**).^[5]

In a similar way, compounds **12** and **13** were glycosylated with **14a** to form the nucleosides **16** and **17**, which were deprotected in NaOMe/MeOH to give the intermediate methoxy derivatives **2** and **3**. Consequently, **2** and **3** were converted to 7-deaza-7-fluoro-2'-deoxyguanosine (**5**) and 7-deaza-7-fluoro-2'-deoxyxanthosine (**8**) in aqueous NaOH or with Me₃SiCl/NaI in MeCN. The isdG derivative **7** was obtained from **6** by photochemically induced displacement reaction performed in water (30% yield) (Scheme 3).



SCHEME 2

The 7-deazapurine 2'-deoxyribonucleosides **4a** and **5** were converted into the corresponding phosphoramidites **9** and **10** by standard procedures. Oligonucleotides containing compound **4a** were synthesized employing compound **9** in solid-phase oligonucleotide synthesis. As supposed to be an analog of the universal nucleoside 7-deaza-2'-deoxyinosine, compound **4a** was incorporated in the duplexes opposite to the canonical 2'-deoxynucleosides dA, dG, dC, and dT. According to Table 1, compound **4a** shows base pairing with duplex stabilities similar to 7-deaza-2'-deoxyinosine (**c⁷Id**).^[9] Thus, the 7-fluoro substituent has only a minor influence when only one 7-deaza-2'-deoxyinosine residue was replaced by the fluoro analog **4a**. However, a strong stabilizing effect was observed when twelve **4a** residues were incorporated in a self-complementary alternating 12-mer duplex **2323** (Table 1).



SCHEME 3

TABLE 1 T_m values and thermodynamic data of oligonucleotides containing **4a** and 7-deaza-2'-deoxyinosine (**c⁷I**)^{a,b}

Duplex	T_m	Duplex	T_m
5'-d(GGAAAA 4a AAAAGG)-3' (18)	43	5'-d(GGAAAA c⁷I AAAAGG)-3'	43 ^[9]
3'-d(CCTTTT A TTTTCC)-5' (19)		3'-d(CCTTTT A TTTTCC)-5'	
5'-d(GGAAAA 4a AAAAGG)-3' (18)	35	5'-d(GGAAAA c⁷I AAAAGG)-3'	37 ^[9]
3'-d(CCTTTT G TTTTCC)-5' (20)		3'-d(CCTTTT G TTTTCC)-5'	
5'-d(GGAAAA 4a AAAAGG)-3' (18)	46	5'-d(GGAAAA c⁷I AAAAGG)-3'	45 ^[9]
3'-d(CCTTTT C TTTTCC)-5' (21)		3'-d(CCTTTT C TTTTCC)-5'	
5'-d(GGAAAA 4a AAAAGG)-3' (18)	39	5'-d(GGAAAA c⁷I AAAAGG)-3'	40 ^[9]
3'-d(CCTTTT T TTTTCC)-5' (22)		3'-d(CCTTTT T TTTTCC)-5'	
5'-d(4aC4aC4aC4aC4aC4aC4a)-3' (23)	29 ^c	5'-d(c⁷ICc⁷ICc⁷ICc⁷ICc⁷ICc⁷IC)-3' (24)	13 ^c
3'-d(C4aC4aC4aC4aC4aC4aC4a)-5' (23)		3'-d(Cc⁷ICc⁷ICc⁷ICc⁷ICc⁷ICc⁷I)-5' (24)	

^aThermodynamic parameters are derived from the fitting of melting curves measured at 260 nm in 1 M NaCl, 100 mM MgCl₂, and 60 mM Na-cacodylate buffer, pH 7.0, with 5 μ M + 5 μ M single-strand concentration. ^b T_m values are given in °C. ^cMeasured in 100 mM NaCl, 10 mM MgCl₂, and 10 mM Na-cacodylate buffer, pH 7.0, with 10 μ M single-strand concentration.

CONCLUSION

7-deaza-7-fluoro-2'-deoxynucleosides were synthesized by nucleobase-anion glycosylation stereoselectively. A fluorine atom introduced in the 2'-arabino position biases the conformation of the sugar toward *N*. oligonucleotides containing 7-fluoro nucleosides increase the T_m value of duplex melting compared to the non-functionalized compounds (see duplex **23.23** versus **24.24**).

REFERENCES

- Seela, F.; Thomas, H. Duplex Stabilization of DNA: Oligo-nucleotides containing 7-substituted 7-deazaadenines. *Helv. Chim. Acta* **1995**, 78, 94–108.
- Ramzaeva, N.; Seela, F. Duplex-Stability of 7-deazapurine DNA: oligonucleotides containing 7-bromo or 7-iodo-7-deazaguanine. *Helv. Chim. Acta* **1996**, 79, 1549–1558.
- Seela, F.; Zulauf, M. 7-deazaadenine-DNA: Bulky 7-iodo substituents or hydrophobic 7-hexynyl chains are well accommodated in the major groove of oligonucleotide duplexes. *Chem. Eur. J.* **1998**, 4, 1781–1790.
- Balow, G.; Mohan, V.; Lesnik, E.A.; Johnston, J.F.; Monia, B.P.; Acevedo, O.L. Biophysical and antisense properties of oligodeoxynucleotides containing 7-propynyl-, 7-iodo- and 7-cyano-7-deaza-2-amino-2'-deoxyadenosines. *Nucleic Acids Res.* **1998**, 26, 3350–3357.
- Seela, F.; Xu, K.; Chittapu, P. Fluorinated pyrrolo[2,3-*d*]pyrimidine nucleosides: 7- Fluoro-7-deazapurine 2'-Deoxyribofuranosides and 2'-deoxy-2'-fluoroarabinofuranosyl derivatives. *Synthesis* **2006**, 2005–2012.
- Wang, X.; Seth, P.P.; Ranken, R.; Swayze, E.E.; Migawa, M.T. Synthesis and biological activity of 5-fluorotubercidin. *Nucleosides, Nucleotides Nucleic Acids* **2004**, 23, 161–170.
- Winkler, H.-D.; Seela, F. Synthesis of 2-amino-7-(2'-deoxy- β -D-erythro-pentofuranosyl)-3,7-dihydro-4H-pyrrolo[2,3-*d*]pyrimidin-4-one, a new isostere of 2'-deoxyguanosine. *J. Org. Chem.* **1983**, 48, 3119–3122.
- Rosemeyer, H.; Seela, F. Stereoselective synthesis of pyrrolo[2,3-*d*]pyrimidine α - and β -D-ribonucleosides from anomerically pure D-ribofuranosyl chlorides: Solid-liquid phase-transfer glycosylation and ¹⁵N-NMR spectra. *Helv. Chim. Acta* **1988**, 71, 1573–1585.
- Seela, F.; Mittelbach, C. 7-deaza-2'-deoxyinosine: A stable nucleoside with the ambiguous base pairing properties of 2'-deoxyinosine. *Nucleosides Nucleotides Nucleic Acids* **1999**, 18, 425–441.