

Total synthesis of paracentrone, C₃₁-allenic apo-carotenoid†

Yusuke Murakami, Masayuki Nakano, Takuya Shimofusa, Noriyuki Furuichi and Shigeo Katsumura*

*School of Science and Technology, Kwansai Gakuin University, Gakuen 2-1, Sanda, Hyogo, 669-1337, Japan. E-mail: katsumura@ksc.kwansei.ac.jp; Fax: +81-79-565-9077; Tel: +81-79-565-8314*Received 10th January 2005, Accepted 23rd February 2005
First published as an Advance Article on the web 11th March 2005

The stereocontrolled total synthesis of a C₃₁-allenic apo-carotenoid, paracentrone, was achieved by the convergent C₂₀ + C₁₁ = C₃₁ strategy. The key elements of our synthesis were the Pd-catalyzed cross-coupling to stereoselectively construct the conjugated polyene backbone skeleton and the designed geometrical isomerization at the central double bond of the conjugated polyene chain. In addition, the terminal oxygenated cyclohexane ring having the allenic moiety was prepared by the highly diastereoselective Sharpless epoxidation under our own reaction conditions.

Paracentrone (1) has been isolated from the sea urchin *Paracentrous lividus*¹ and the structure of this unique molecule was determined by Weedon and co-workers in 1969. This molecule is a C₃₁-apo-carotenoid, displaying a dihydroxycyclohexylidene allenic moiety at one terminal and a methyl ketone group at the other (Fig. 1).² It has been postulated that the naturally occurring paracentrone (1) is produced by the degradation of C₄₀-carotenoids, fucoxanthin or amarouciaxanthin A, in animals.³ The first synthesis of paracentrone (1) was achieved by Haugan in 1996 over 13 linear steps from the readily available (3*R*,5*R*)-actinol (7) with the C₁₅ + C₁₀ + C₅ + C₁ = C₃₁ strategy.⁴




Fig. 1 Paracentrone (1) and peridinin (2).

Our longstanding interest in carotenoids and related natural products^{5,6} led to the achievement of the stereocontrolled and convergent total synthesis of peridinin (2), a polyfunctional C₃₇-allenic nor-carotenoid.⁷ Based on this synthesis, we have started to develop a new stereocontrolled synthetic strategy for naturally occurring allenic carotenoids.⁸ In this communication, we disclose the stereocontrolled total synthesis of the C₃₁-allenic apo-carotenoid, paracentrone (1), using the convergent C₂₀ + C₁₁ = C₃₁ strategy.

The retrosynthetic disconnection at the C(15)–C(15') position, which is the central part of the conjugated olefin, led to two fragments; the C₂₀-allenic triol 3 and the C₁₁-trienol 4 (Fig. 2). We envisioned that the former would be provided from acetylene 5 and trienyl vinyl iodide 6 by employing the Sonogashira cross-coupling⁹ and S_N2' hydride reduction.¹⁰ The acetylene 5 was easily available from 7 via the Sharpless asymmetric epoxidation under the precise conditions established in our laboratory.^{5b,7}





Fig. 2 Retrosynthetic analysis.

The preparation of trienyl iodide 6 was started from the known vinyl stannane 8¹¹ using the modified de Lera's procedure (Scheme 1).¹² Vinyl stannane 8 was smoothly converted in high yield into the vinyl iodide 9 upon treatment with iodine in CH₂Cl₂. The addition of the carbanion derived from phosphonate 10¹³ to the corresponding aldehyde, prepared from 9, in the presence of DMPU¹⁴ gave the desired 6 in a highly stereoselective fashion (70% for two steps, 11*E* only, 13*E* : 13*Z* = 10 : 1 (carotenoid numbering)).




Scheme 1 Preparation of the trienyl iodide 6. *Reagents and conditions:* (a) I₂, Na₂CO₃, CH₂Cl₂, 0 °C, 10 min, quant.; (b) MnO₂, Na₂CO₃, ether, rt, 2 h; (c) 10, *n*BuLi, DMPU, THF, –20 °C, 5 min, 70% for two steps.

The Pd- and Cu-catalyzed Sonogashira reaction⁹ between acetylene 5^{5b,7} and trienyl iodide 6 proceeded smoothly to furnish the desired coupling product 11, with the complete retention of the stereochemistry, in 81% yield (Scheme 2). The reduction of 11 with DIBAL for the construction of the allenic moiety¹⁰ afforded the C₂₀-allenic triol 3,¹⁵ which was oxidized into the corresponding aldehyde 12 (13*E* : 13*Z* = 10 : 1 (carotenoid numbering)) with Dess–Martin periodinane.¹⁶ These C₂₀-allenic compounds 3 and 12 are not only the key intermediates for our paracentrone synthesis, but also the valuable half-fragments for the naturally occurring C₄₀-allenic carotenoids, for example fucoxanthin and neoxanthin (*vide infra*).


Meanwhile, the stereoselective preparation of the conjugated trienol 4 was also achieved by the consecutive Pd-catalyzed cross-coupling reactions. Carbon chain extension of the vinyl iodide 9 with (trimethylsilyl)acetylene under the Sonogashira conditions,⁹ followed by the sequential desilylation and stannylation of the resulting terminal acetylene moiety¹⁷ produced

† Electronic supplementary information (ESI) available: general experimental details. See <http://www.rsc.org/suppdata/ob/b5/b500316d/>



Scheme 2 Synthesis of the C_{20} -allenic triol **3** and aldehyde **12**. *Reagents and conditions:* (a) $\text{Pd}(\text{PPh}_3)_4$, CuI , $i\text{Pr}_2\text{NH}$, rt, 4 h, 81%; (b) DIBAL, CH_2Cl_2 , 0 °C, 30 min, 80%; (c) Dess–Martin periodinane, AcOEt , rt, 10 min.


the dienyl stannane **13** in 63% overall yield (Scheme 3). The vinyl bromide **15**, which is another component for **4**, was easily prepared from methyl bromomethacrylate **14**¹⁸ by the simple functional group manipulation. The Stille coupling¹⁹ between **13** and **15** in the presence of $\text{Pd}(\text{PPh}_3)_4$ and LiCl in DMF successfully afforded trienol **4**²⁰ bearing the desired all-*trans* stereochemistry in good yield.



Scheme 3 Synthesis of the trienol **4** and phosphonium salt **19**. *Reagents and conditions:* (a) (trimethylsilyl)acetylene, $\text{Pd}(\text{PPh}_3)_4$, CuI , Et_3N , THF, rt, 2 h; (b) K_2CO_3 , MeOH , rt, 2 h, 78% for two steps; (c) $n\text{Bu}_3\text{Sn}(n\text{Bu})\text{CuCNLi}_2$, THF, –30 °C, 1.5 h, 80%; (d) LiAlH_4 , ether, rt, 1 h, quant.; (e) MnO_2 , acetone, rt, 12 h; (f) MeMgBr , ether, rt, 30 min, 84% for two steps; (g) MnO_2 , ether, rt, 12 h, 83%; (h) $\text{Pd}(\text{PPh}_3)_4$, LiCl , DMF, 75 °C, 1 h, 84%; (i) PBr_3 , CH_2Cl_2 , 0 °C, 10 min; (j) PPh_3 , AcOEt , rt, 15 h, 74% for two steps; (k) 2-mercaptobenzothiazol, PPh_3 , DEAD, THF, rt, 30 min, 77%.

With the C_{20} -allenic aldehyde **12** and the C_{11} -trienol **4** in hand, our attention turned to the fragment union by the olefination reaction. First, the conversion of trienol **4** into the corresponding benzothiazoyl sulfone **18** for the modified Julia–Kocienski reaction²¹ was attempted, which was effectively utilized in our previous peridinin synthesis.⁷ The trienol **4** was transformed into the corresponding benzothiazolyl sulfide **16** by the Mitsunobu reaction with 2-mercaptobenzothiazole and the oxidation of **16** into the corresponding sulfone **18** was tried by keeping the geometry of all the *E*-olefins. However, this transformation unfortunately failed and caused a significant *E*–*Z* isomerization of the double bond under sulfur-atom oxidation conditions. After numerous attempts, we knew that the isomerization occurred from the corresponding sulfoxide **17** resulting from the [2,3]-sigmatropic rearrangement.²² On the other hand, the connection of the fragments was successfully achieved by the Wittig reaction, which is traditionally used in carotenoid synthesis. In this event, the trienol **4** was transformed into phosphonium salt **19** via the corresponding allylic bromide by the standard method. The Wittig condensation between **12** and **19** using NaH as a base proceeded at rt in the dark for three hours to afford a mixture of more than four isomers, including the major component in HPLC in 61% yield

(Scheme 4). This was probably caused by the *E/Z* mixtures at the C(13) and C(15) positions. Fortunately, we found that the mixture gradually converged to the major isomer at rt in CDCl_3 .²³ After one hour, the major isomer comprised nearly 70% of the products and was isolated by HPLC purification.²⁴ The spectral data of this isolated major isomer were in good agreement with those of the reported natural product.^{4,25} Thus, we achieved the stereocontrolled total synthesis of the C_{31} -allenic *apo*-carotenoid, paracentrone (**1**), by the convergent $C_{20} + C_{11} = C_{31}$ strategy.



Scheme 4 Total synthesis of paracentrone. *Reagents and conditions:* (a) NaH , DME, CH_2Cl_2 , rt, 3 h, 61%; (b) HPLC purification.²⁴

In our achievement of the total synthesis of the allenic carotenoids, peridinin (**2**)⁷ and paracentrone (**1**), we actually found that the central disubstituted double bond in the carotenoid molecules is easily isomerized from *cis* to *trans* in an acceptable ratio. These phenomena were also studied in detail in the field of spectroscopy, including the calculation of the electron distribution of the conjugated polyene systems.²⁶ Accordingly, the main subject for the stereocontrolled synthesis of the allenic carotenoids can be simplified based on how we synthesize the two fragments bisected at the central double bond in the polyene chain in a stereocontrolled manner. Based on these observations, we propose an effective $C_{20} + C_{20} = C_{40}$ strategy for the stereocontrolled synthesis of allenic carotenoids. This strategy would be generally applicable to the synthesis of widely occurring C_{40} -allenic carotenoids and is distinguishable from the previously employed $C_{15} + C_{10} + C_{15} = C_{40}$ one.²⁷

Acknowledgements

We thank Professor M. Ito and Dr Y. Yamano at Kobe Pharmaceutical University for their kind advice on the HPLC purification of **1**. We also thank Professor Y. Koyama and Dr R. Fujii at Kwansai Gakuin University for their valuable discussions. This work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan. N. F. is grateful for the research fellowships of the Japan Society for the Promotion of Science for Young Scientists and also for the Sasagawa science research grant from The Japan Science Society.

References

- G. Galasko, J. Hora, T. P. Toube, B. C. L. Weedon, D. Andre, M. Barbier, E. Lederer and V. R. Villanueva, *J. Chem. Soc. C*, 1969, 1264.
- (a) L. Bartlett, W. Klyne, W. P. Mose, P. M. Scopes, G. Galasko, A. K. Mallams, B. C. L. Weedon, J. Szabolcs and G. Toth, *J. Chem. Soc. C*, 1969, 2527; (b) T. E. DeVille, M. B. Hursthouse, S. W. Russell and B. C. L. Weedon, *Chem. Commun.*, 1969, 754; (c) J. R. Hlubucek, J. Hora, S. W. Russell, T. P. Toube and B. C. L. Weedon, *J. Chem. Soc., Perkin Trans. 1*, 1974, 848.
- (a) J. Hora, T. P. Toube and B. C. L. Weedon, *J. Chem. Soc. C*, 1970, 241; (b) T. Matsuno, M. Okubo and T. Komori, *J. Nat. Prod.*, 1985, 48, 606.
- (a) J. A. Haugan, *Tetrahedron Lett.*, 1996, 37, 3887; (b) J. A. Haugan, *J. Chem. Soc., Perkin Trans. 1*, 1997, 2731.
- (a) M. Nakano, N. Furuichi, H. Mori and S. Katsumura, *Tetrahedron Lett.*, 2001, 42, 7307; (b) M. Kuba, N. Furuichi and S. Katsumura, *Chem. Lett.*, 2002, 1248.
- For our early work in this area see: (a) S. Isoe, S. B. Hyeon, H. Ichikawa, S. Katsumura and T. Sakan, *Tetrahedron Lett.*, 1968, 5561; (b) S. Isoe, S. Katsumura, S. B. Hyeon and T. Sakan, *Tetrahedron*

- Let.*, 1971, 1089; (c) S. Isoe, S. B. Hyeon, S. Katsumura and T. Sakan, *Tetrahedron Lett.*, 1972, 2517; (d) S. Isoe, S. Katsumura and T. Sakan, *Helv. Chim. Acta*, 1973, **56**, 1514.
- 7 (a) N. Furuichi, H. Hara, T. Osaki, H. Mori and S. Katsumura, *Angew. Chem., Int. Ed.*, 2002, **41**, 1023; (b) N. Furuichi, H. Hara, T. Osaki, M. Nakano, H. Mori and S. Katsumura, *J. Org. Chem.*, 2004, **69**, 7949.
- 8 A. Hoffmann-Roder and N. Krause, *Angew. Chem., Int. Ed.*, 2004, **43**, 1196.
- 9 K. Sonogashira, in *Comprehensive Organic Synthesis*, ed. B. M. Trost and I. Fleming, Pergamon Press, Oxford, 1991, vol. 3, p. 521 and references cited therein.
- 10 A. Baumeler, W. Brade, A. Haag and C. H. Eugster, *Helv. Chim. Acta*, 1990, **73**, 700.
- 11 (a) H. Miyake and K. Yamamura, *Chem. Lett.*, 1989, 981; (b) J.-F. Betzer, F. Delalobe, B. Muller, A. Pancrazi and J. Prunet, *J. Org. Chem.*, 1997, **62**, 7768 and references cited therein.
- 12 B. Domingues, B. Iglesias and A. R. de Lera, *Tetrahedron*, 1999, **55**, 15071.
- 13 (a) I. Ahmed, R. N. Gedge and A. Nechwatel, *J. Chem. Soc. C*, 1968, 185; (b) G. Pattenden and B. C. L. Weedon, *J. Chem. Soc. C*, 1968, 1984.
- 14 Y. L. Behnani and M. F. Boehm, *J. Org. Chem.*, 1995, **60**, 1195.
- 15 Data for **3**: $[\alpha]_D^{25}$ -23.47 (c 0.58, CHCl₃); IR (KBr, cm⁻¹) 3356, 2961, 2928, 2860, 1930, 1724, 1641, 1454, 1373, 1253, 1157, 1070; ¹H NMR (400 MHz, CD₃OD) δ 6.54 (dd, 1H, *J* = 15.1, 11.2 Hz), 6.27 (d, 1H, *J* = 15.1 Hz), 6.07 (d, 1H, *J* = 10.5 Hz), 6.02 (s, 1H), 5.64 (t, 1H, *J* = 6.8 Hz), 4.22 (d, 2H, *J* = 6.8 Hz), 4.20 (m, 1H), 2.18 (ddd, 1H, *J* = 12.7, 3.9, 2.0 Hz), 1.89 (ddd, 1H, *J* = 14.9, 3.9, 2.0 Hz), 1.83 (s, 3H), 1.80 (s, 3H), 1.50–1.20 (m, 2H), 1.33 (s, 3H), 1.33 (s, 3H), 1.06 (s, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 203.4, 137.7, 137.2, 133.1, 131.8, 128.9, 125.8, 118.4, 103.6, 73.1, 64.7, 59.5, 50.6, 50.1, 36.6, 32.9, 31.4, 29.6, 14.2, 12.6; EI⁺ HRMS found *m/z* 318.2193, calcd for C₂₀H₃₀O₃, M⁺ 318.2194.
- 16 D. B. Dess and J. C. Martin, *J. Org. Chem.*, 1983, **48**, 4156.
- 17 (a) B. H. Lipshutz, J. A. Kozlowski and R. S. Wilhelm, *J. Org. Chem.*, 1984, **49**, 3943; (b) see also: ref. 11b.
- 18 P. A. Bieber, *Ann. Chem.*, 1954, **9**, 674.
- 19 J. K. Stille, *Angew. Chem., Int. Ed. Engl.*, 1980, **25**, 508 and references cited therein.
- 20 Data for **4**: IR (KBr, cm⁻¹) 3430, 3054, 3000, 2922, 1647, 1607, 1435, 1395, 1366, 1319, 1290, 1233, 1171, 1096, 1073, 1030; ¹H NMR (400 MHz, CDCl₃) δ 7.10 (d, 1H, *J* = 9.5 Hz), 6.62 (d, 1H, *J* = 15.1 Hz), 6.55 (dd, 1H, *J* = 15.1, 9.5 Hz), 5.87 (t, 1H, *J* = 6.6 Hz), 4.36 (d, 2H, *J* = 6.6 Hz), 2.36 (s, 3H), 1.92 (d, 3H, *J* = 1.2 Hz), 1.88 (d, 3H, *J* = 0.5 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 199.7, 143.9, 139.6, 136.1, 135.6, 134.9, 123.8, 59.4, 25.5, 12.5, 11.6.
- 21 (a) J. B. Baudin, G. Hareau, S. A. Julia and O. Ruel, *Tetrahedron Lett.*, 1991, **32**, 1175; (b) J. B. Baudin, G. Hareau, S. A. Julia and O. Ruel, *Bull. Soc. Chim. Fr.*, 1993, **130**, 336; (c) J. B. Baudin, G. Hareau, S. A. Julia, R. Lorne and O. Ruel, *Bull. Soc. Chim. Fr.*, 1993, **130**, 856; (d) P. R. Blakemore, W. J. Cole, P. J. Kocienski and A. Morley, *Synlett*, 1998, **1**, 26; (e) P. R. Blakemore, *J. Chem. Soc., Perkin Trans. 1*, 2002, 2563 and references cited therein.
- 22 The main pathway in the side-reactions would involve the [2,3]-sigmatropic rearrangement of the intermediate sulfoxide to the sulfenate ester, the details of which will be reported in a full account of this work. See: D. A. Evans and G. C. Andrew, *Acc. Chem. Res.*, 1974, **7**, 147.
- 23 The thermal isomerization from *cis* to *trans* at the central C(15)–C(15') disubstituted double bond was also observed in our peridinin synthesis.⁷ This feature was followed and detected by HPLC analysis, the details of which will be reported in a full account of this work.
- 24 Conditions: column: Develosil 60-5 (4.6 × 250 mm); mobile phase: THF–hexane–*i*PrOH = 15 : 67 : 3; flow rate: 1.0 ml min⁻¹; UV-detection: 450 nm; retention time: 22 min (all-*E*-isomer).
- 25 Data for paracentrone (**1**): IR (KBr, cm⁻¹) 3407, 2961, 2924, 2855, 1929, 1721, 1649, 1607, 1530, 1453, 1368, 1321, 1279, 1229, 1157, 1071, 1040, 992, 963; ¹H NMR (400 MHz, CDCl₃) δ 7.14 (dq, 1H, *J* = 10.5, 1.0 Hz), 6.73 (dd, 1H, *J* = 14.2, 11.5 Hz), 6.66 (d, 1H, *J* = 14.9 Hz), 6.63 (dd, 1H, *J* = 14.1, 11.5 Hz), 6.60 (dd, 1H, *J* = 14.6, 11.5 Hz), 6.59 (dd, 1H, *J* = 15.4, 10.7 Hz), 6.39 (d, 1H, *J* = 11.5 Hz), 6.34 (d, 1H, *J* = 15.1 Hz), 6.26 (d, 1H, *J* = 11.0 Hz), 6.12 (d, 1H, *J* = 11.3 Hz), 6.03 (s, 1H), 4.32 (m, 1H), 2.36 (s, 3H), 2.27 (ddd, 1H, *J* = 13.3, 4.2, 1.9 Hz), 1.991 (s, 3H), 1.985 (s, 3H), 1.96 (m, 1H), 1.94 (d, 3H, *J* = 1.0 Hz), 1.81 (s, 3H), 1.41 (m, 1H), 1.35 (s, 3H), 1.33 (s, 3H), 1.35 (m, 1H), 1.07 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 202.4, 199.4, 144.5, 139.9, 137.9, 137.0, 136.2, 135.5, 132.6, 132.3, 132.1, 129.4, 128.4, 125.6, 123.8, 117.7, 103.2, 73.0, 64.3, 49.5, 48.9, 35.8, 32.2, 31.4, 29.4, 25.6, 14.0, 12.9, 12.8, 11.7; ESI HRMS found *m/z* 461.3065, calcd for C₃₁H₄₂O₃ [M – H]⁻ 461.3056.
- 26 M. Kuki, Y. Koyama and H. Nagae, *J. Phys. Chem.*, 1991, **95**, 7171.
- 27 (a) A. Baumeler and C. H. Eugster, *Helv. Chim. Acta*, 1991, **74**, 469; (b) A. Baumeler and C. H. Eugster, *Helv. Chim. Acta*, 1992, **75**, 773; (c) Y. Yamano and M. Ito, *Chem. Pharm. Bull.*, 1994, **42**, 410; (d) Y. Yamano and M. Ito, *J. Chem. Soc., Perkin Trans. 1*, 1995, 1895; see also; (e) Y. Yamano and M. Ito, *J. Chem. Soc., Perkin Trans. 1*, 1993, 1599; (f) ref. 4.