

Preparation of 1,4-hydrobenzoquinones by the PCC/SiO₂-promoted double oxidation of 3-cyclohexene-1,2-diols

Hee Jin Kim and Sangho Koo*

Department of Chemistry, Myong Ji University, Yongin, Kyunggi-Do, 449-728, Korea.
E-mail: sangkoo@mju.ac.kr; Fax: 82 31 335 7248; Tel: 82 31 330 6185


Received 3rd August 2005, Accepted 26th August 2005
First published as an Advance Article on the web 2nd September 2005

The PCC/SiO₂-promoted double oxidation of 3-cyclohexene-1,2-diols, which were easily prepared by the two-step sequence of α -hydroxylation of various conjugated cyclohexenones and the subsequent nucleophilic carbonyl addition of alkyl anions, produced diversely substituted 1,4-hydrobenzoquinones.

Quinones or hydroquinones are useful building blocks for constructing polycyclic compounds in organic synthesis.¹ Quinoids containing polyphenyl side chains such as ubiquinones, menaquinones, and plastoquinones are biologically important natural products, which show physiological and clinical activities.² Most of the synthetic methods for these chemically and biologically important quinones or hydroquinones are based on the oxidation of phenol or methoxybenzene derivatives,³ which generally require strongly acidic or basic conditions.

We devised a totally different synthetic approach to 1,4-hydrobenzoquinones **2** by the double oxidation of 3-cyclohexene-1,2-diols **1** (Scheme 1). While the secondary homoallylic alcohol of **1** is oxidized to the ketone of **A**, the chromate ester of the sterically congested tertiary allylic alcohol of **B** ($R \neq H$) undergoes allylic migration to give the ester of the secondary allylic alcohol of **C**, which can be oxidized to the ketone of **D**. Enolization of the 1,4-diketone **D** gives rise to 1,4-hydrobenzoquinone **2**. This scenario has proven to be successful for 2-methyl-3-cyclohexene-1,2-diol (**1**) ($R = \text{Me}$ in Scheme 1), where 2-methyl-1,4-hydrobenzoquinone (**2**) was obtained after the double oxidation with the mild Cr(VI)-based reagents in CH₂Cl₂ at RT (Table 1). An excess of oxidant (4 equiv.) was required to give the optimum yield, even though two equiv. was theoretically necessary. Silica gel supported pyridinium chlorochromate⁴ (PCC) provided the best result (entry 4), which was prepared by mixing and pulverizing equal weight amounts of each material.

The scope of this synthetic method for 1,4-hydrobenzoquinones **2** depends on the availability of diversely substituted 3-cyclohexene-1,2-diols **1**. The preparation and the double oxidation of **1** was delineated in Scheme 2, and summarized in Table 2. We have reported a highly efficient synthetic method of methyl-substituted conjugated cyclohexenones by the reaction of β -ketoesters and α,β -unsaturated carbonyl compounds under



Scheme 1 A synthetic plan for 1,4-hydrobenzoquinones **2** by the double oxidation of 3-cyclohexene-1,2-diols **1** ($R = \text{alkyl}$).

t-BuOK–*t*-BuOH conditions.⁵ We prepared 2-cyclohexen-1-ones **3** with methyl or ethyl substituents at the 2-, 4-, and 5-positions, which would constitute the three substituents (R^1 , R^2 , and R^3 , respectively) of the 1,4-hydrobenzoquinones **2**. Pb(OAc)₄-mediated direct α -acetoxylation⁶ of the conjugated cyclohexenones **3** in refluxing toluene (conditions: A) produced 6-acetoxy-2-cyclohexen-1-ones **4** in reasonable yields except for *R*-(–)-carvone (**3f**), where no reaction was observed. α -Hydroxylation of the sterically demanding *R*-(–)-carvone (**3f**) should be carried out by the peroxycarboxylic acid oxidation of the corresponding trimethylsilyl (TMS) enol ether,⁷ which was prepared by the reaction with TMSCl in the presence of sodium hexamethyldisilazide (NaHMDS, conditions: B). Desilylated 6-hydroxy-2-cyclohexen-1-one **4f** was obtained in 90% yield in this case (entry 16). The fourth substituent R^4 was introduced by the nucleophilic addition of alkyl metal species to the carbonyl group of **4**. MeLi, EtMgBr, and *n*-BuLi were used to add Me, Et, and *n*-Bu groups, respectively to the six different cyclohexenones **4a–f** to produce eighteen different 3-cyclohexene-1,2-diols **1a–r** in reasonable yields.⁸

The key double oxidation of the 3-cyclohexene-1,2-diols **1a–r** proceeded well under the standard conditions of 4 equiv. of silica supported PCC in CH₂Cl₂ at RT for 1 h to give rise to the diversely substituted 1,4-hydrobenzoquinones **2a–r** in decent yields.⁹ 1,4-Hydrobenzoquinones **2** were formed exclusively under an argon atmosphere even with an excess (4 equiv.) of PCC oxidant. However, it was not possible to avoid obtaining the oxidized 1,4-benzoquinones during the purification process of SiO₂ column chromatography. The actual yield of 1,4-hydrobenzoquinones **2** must be higher by *ca.* 10% than the reported yields in Table 2.


We then studied the possibility of synthesizing 1,2-hydrobenzoquinones **5** by the PCC/SiO₂ oxidation of the 3-cyclohexene-1,2-diols **1s**, **1t**, and **1u**, which were easily prepared by hydride addition to the carbonyl groups of **4a**, **4b**, and **4f**, respectively (Scheme 3 and Table 3). We expected that the double oxidation of the secondary allylic and homoallylic

Table 1 The optimization of the double oxidation of 2-methyl-3-cyclohexene-1,2-diol (**1**) ($R = \text{Me}$ in Scheme 1)


Entry	Oxidant/equiv.	Reaction condition	Yield 2 (%) ^a
1	PCC (2)	RT for 3 h in CH ₂ Cl ₂	30
2	PDC (4)	RT for 1 h in CH ₂ Cl ₂	33
3	PCC (4)	RT for 2 h in CH ₂ Cl ₂	56
4	PCC (4)–SiO ₂	RT for 1 h in CH ₂ Cl ₂	78


^a Isolated yield of 2-methyl-1,4-hydrobenzoquinone (**2**) after SiO₂ chromatographic separation. 1,4-Benzoquinone was also obtained during the purification process, the yield of which was about 10% of the yield **2**.

Table 2 The yields of each reaction in Scheme 2 for diverse substrates


Entry	3	R ¹	R ²	R ³	Conditions ^a	4	R	Yield 4 (%) ^b	1	R ⁴ ^c	Yield 1 (%) ^b	2	Yield 2 (%) ^{b,d}
1	3a	H	H	Me	A	4a	Ac	81	1a	Me	76	2a	71
2									1b	Et	55	2b	59
3									1c	Bu	52	2c	62
4	3b	H	Me	H	A	4b	Ac	65	1d	Me	80	2d	70
5									1e	Et	69	2e	66
6									1f	Bu	54	2f	78
7	3c	Me	H	H	A	4c	Ac	68	1g	Me	71	2g	67
8									1h	Et	67	2h	57
9									1i	Bu	66	2i	52
10	3d	Me	H	Me	A	4d	Ac	70	1j	Me	75	2j	64
11									1k	Et	64	2k	55
12									1l	Bu	59	2l	69
13	3e	H	Me	Et	A	4e	Ac	77	1m	Me	91	2m	81
14									1n	Et	72	2n	63
15									1o	Bu	81	2o	69
16	3f	Me	H		B	4f	H	90	1p	Me	77	2p	77
17									1q	Et	56	2q	67
18									1r	Bu	70	2r	60

^a Reaction conditions for 4, A: 3 and Pb(OAc)₄ (2 equiv.) in toluene at reflux for 4 h; B: (i) 3, TMSCl, and NaHMDS in THF at 0 °C, (ii) urea-H₂O₂ (UHP) and phthalic anhydride in MeCN at RT for 5 h. ^b Isolated yields after SiO₂ chromatographic separation. ^c An excess amount (3 equiv.) of MeLi, EtMgBr, or *n*-BuLi was used to produce 3-cyclohexene-1,2-diols 1. ^d A small amount of 1,4-benzoquinone was also obtained during the purification process, which was about 10% of the yield of 2.

**Scheme 2** Synthesis of 1,4-hydrobenzoquinones 2 from 2-cyclohexen-1-ones 3.**Table 3** The yield 4 of the reaction in Scheme 3

Entry	1	R ¹	R ²	R ³	4	Yield of 4 (%) ^a
1	1s	H	H	Me	4s	78
2	1t	H	Me	H	4t	69
3	1u	Me	H		4f	83

^a Isolated yields after SiO₂ chromatographic separation.

**Scheme 3** The oxidation of 3-cyclohexene-1,2-diols 1s-u using the standard silica supported PCC (4 equiv.) conditions at RT (see Table 3).

alcohols of 1s-u would directly produce the corresponding 1,2-hydrobenzoquinones 5. However, only the mono oxidation of the allylic alcohol of 1s-u was observed to produce the 6-hydroxy-2-cyclohexen-1-ones 4s, 4t and 4f in 69–83% yields. No 1,2-hydrobenzoquinone 5 was obtained under the above PCC/SiO₂ oxidation conditions.

In conclusion, we have developed a novel and efficient synthetic method for diversely alkyl-substituted 1,4-hydrobenzoquinones 2 by the mild PCC/SiO₂-promoted double oxidation of 3-cyclohexene-1,2-diols 1, which were readily prepared from the corresponding 2-cyclohexen-1-ones 3. This double oxidation requires the secondary allylic alcohol and a tertiary allylic alcohol in the 1,2-position to produce 1,4-hydrobenzoquinones 2 after allylic migration of the tertiary allylic alcohol and the

subsequent oxidations. The 1,4-hydrobenzoquinones can be efficiently utilized in the total synthesis of various polycyclic natural products.

This work was supported by the Korea Research Foundation Grant (KRF-2003-015-C00342).

References

- (a) M. A. Ott and J. H. Noordik, *J. Chem. Inf. Comput. Sci.*, 1997, **37**, 98; (b) J. A. Valderrama, D. Pessoa-Mahana, R. A. Tapia, A. R. de Arias, H. Nakayama, S. Torres, J. Miret and M. E. Ferreira, *Tetrahedron*, 2001, **57**, 8653.
- (a) R. H. Thomson, *Naturally Occurring Quinones*, Academic Press, New York, 1971; (b) P. Mitchell and J. Moyle, *Coenzyme Q*, ed. G. Lenaz, Wiley, Chichester, UK, 1985; (c) G. Lenaz, *Coenzyme Q, Biochemistry, Bioenergetics and Clinical Applications of Ubiquinone*, Wiley-Interscience, New York, 1982.
- (a) L. Syper, K. Kloc and J. Młochowski, *Tetrahedron*, 1980, **36**, 123; (b) W. Adam, W. A. Herrmann, J. Lin, C. R. Saha-Möller, R. W. Fischer and J. D. G. Correia, *Angew. Chem., Int. Ed.*, 1994, **33**, 2475; (c) E. Keinan and D. Eren, *J. Org. Chem.*, 1987, **52**, 3872; (d) H. Tohma, H. Morioka, Y. Harayama, M. Hashizume and Y. Kita, *Tetrahedron Lett.*, 2001, **42**, 6899; (e) A. S. Chida, P. V. S. N. Vani, M. Chandrasekharan, R. Srinivasan and A. K. Singh, *Synth. Commun.*, 2001, **31**, 657; (f) O. Cherkaoui, P. Nebois and H. Fillion, *Tetrahedron*, 1996, **52**, 9499; (g) S.-I. Murahashi, T. Naota, N. Miyaguchi and S. Noda, *J. Am. Chem. Soc.*, 1996, **118**, 2509; (h) R. Salsdino, V. Neri, E. Mincione and P. Filippone, *Tetrahedron*, 2002, **58**, 8493.
- (a) F. A. Luzzio, R. W. Fitch, W. J. Moore and K. J. Mudd, *J. Chem. Educ.*, 1999, **76**, 974; (b) G. Piancatelli, A. Scettri and M. D'Auria, *Synthesis*, 1982, 245.
- B.-D. Chong, Y.-I. Ji, S.-S. Oh, J.-D. Yang, W. Baik and S. Koo, *J. Org. Chem.*, 1997, **62**, 9323.
- J. E. Yeo, X. Yang, H. J. Kim and S. Koo, *Chem. Commun.*, 2004, 236.
- (a) G. M. Rubottom and J. M. Gruber, *J. Org. Chem.*, 1978, **43**, 1599; (b) S. Choi and S. Koo, *J. Org. Chem.*, 2005, **70**, 3328.
- The representative experimental procedure for 1a: To a stirred solution of 4a (0.26 g, 1.5 mmol) in THF at –78 °C under an argon atmosphere was added a 1.6 M solution of MeLi in ether (2.9 mL, 4.6 mmol). The mixture was stirred at that temperature for 1 h and then at

RT for 2 h. The reaction was quenched with 10% aqueous NH_4Cl solution, and extracted with ethyl acetate. The organic mixture was washed with saturated aqueous NaCl solution, dried over anhydrous K_2CO_3 , filtered, and concentrated under reduced pressure. The crude product was purified by SiO_2 flash column chromatography to give **1a** (0.17 g, 1.2 mmol) in 76% yield. Data for **1a** (major isomer): ^1H NMR δ 1.09 (d, $J = 6.2$ Hz, 3H), 1.29 (s, 3H), 1.71 (s, 1H), 1.76–2.08 (m, 3H), 2.72 (s, 1H), 3.51 (d, $J = 5.0$ Hz, 1H), 5.49 (dt of A of ABq, $J_{\text{AB}} = 9.8$, $J_{\text{d}} = 1.8$, $J_{\text{t}} = 0.9$ Hz, 1H), 5.69 (dt of B of ABq, $J_{\text{AB}} = 9.8$, $J_{\text{d}} = 1.4$, $J_{\text{t}} = 0.9$ Hz, 1H) ppm; ^{13}C NMR δ 18.3, 28.4, 30.9, 31.9, 71.4, 76.9, 126.8, 131.7 ppm; IR (KBr) 3419, 1450, 1373, 1240 cm^{-1} .

9 The representative experimental procedure for **2a**: A mixture of PCC (1.20 g, 5.6 mmol) and silica gel (1.20 g, 70–230 mesh) was pulverized to give a light orange fine powder, which was added to a stirred solution of **1a** (0.17 g, 1.4 mmol) in CH_2Cl_2 at RT under an argon atmosphere. The mixture was stirred for 1 h, and the resulting mixture was filtered through a short pad of silica gel. The filtrate was concentrated under reduced pressure. The crude product was purified by SiO_2 flash column chromatography to give **2a** (0.13 g, 0.98 mmol) in 71% yield. The corresponding 1,4-benzoquinone (0.01, 0.08 mmol) was also obtained in 6% yield. Data for **2a**: ^1H NMR δ 2.18 (d, $J = 2.2$ Hz, 6H), 5.44 (br s, 1H), 5.47 (br s, 1H), 5.93 (br s, 2H) ppm; ^{13}C NMR δ 17.8, 128.2, 128.5, 144.1, 158.7 ppm; IR (KBr) 3450, 1377, 1235 cm^{-1} .