

# Concurrent primary and secondary deuterium kinetic isotope effects in anilinolysis of *O*-aryl methyl phosphonochloridothioates†

Md. Ehtesham Ul Hoque, Arun Kanti Guha, Chan Kyung Kim, Bon-Su Lee\* and Hai Whang Lee\*

Received 16th February 2009, Accepted 15th April 2009

First published as an Advance Article on the web 28th May 2009

DOI: 10.1039/b903148k

The nucleophilic substitution reactions of *Y*-*O*-aryl methyl phosphonochloridothioates with substituted anilines (XC<sub>6</sub>H<sub>4</sub>NH<sub>2</sub>) and deuterated anilines (XC<sub>6</sub>H<sub>4</sub>ND<sub>2</sub>) are investigated kinetically in acetonitrile at 55.0 °C. The Hammett and Brønsted plots for substituent (X) variations in the nucleophiles are biphasic concave downwards with a break region between X = H and 4-Cl. The deuterium kinetic isotope effects (DKIEs) are primary normal ( $k_{\text{H}}/k_{\text{D}} = 1.03\text{--}1.30$ ) for stronger nucleophiles (X = 4-MeO, 4-Me and H), and *extremely large* secondary inverse ( $k_{\text{H}}/k_{\text{D}} = 0.367\text{--}0.567$ ) for weaker nucleophiles (X = 4-Cl, 3-Cl and 3-NO<sub>2</sub>). The cross-interaction constants are negative ( $\rho_{\text{XY(H)}} = -0.95$  and  $\rho_{\text{XY(D)}} = -1.11$ ) for stronger nucleophiles, while positive ( $\rho_{\text{XY(H)}} = +0.77$  and  $\rho_{\text{XY(D)}} = +0.21$ ) for weaker nucleophiles. These kinetic results indicate that the mechanism changes from a concerted process involving frontside nucleophilic attack for stronger nucleophiles to a stepwise process with a rate-limiting leaving group expulsion from the intermediate involving backside attack for weaker nucleophiles. A hydrogen-bonded, four-center-type transition state (TS) is suggested for a frontside attack, while a trigonal bipyramidal pentacoordinate TS is suggested for a backside attack. The *unusually small* DKIEs, as small as or equal to 0.4, for weaker nucleophiles seem to be ascribed to severe steric congestion in the TS.

## Introduction

Kinetic isotope effects (KIEs) provide one of the most powerful tools for the elucidation of reaction mechanisms.<sup>1</sup> KIEs can give information about the transition state (TS) structure and can indicate the rate-limiting step in a reaction.<sup>2</sup>

In this respect, deuterium kinetic isotope effects (DKIEs) involving deuterated aniline (XC<sub>6</sub>H<sub>4</sub>ND<sub>2</sub>) nucleophiles have provided a useful means to determine the TS structures in nucleophilic substitution reactions, and how the reactants, especially through changes in substituents, alter the TS structures. Incorporation of deuterium in the nucleophile has an advantage in that the  $\alpha$ -DKIEs reflect only the degree of bond formation. When partial deprotonation of the aniline occurs in a rate-limiting step by hydrogen bonding, the  $k_{\text{H}}/k_{\text{D}}$  values are greater than unity. In contrast, DKIEs can only be inverse ( $k_{\text{H}}/k_{\text{D}} < 1.0$ ) in a normal S<sub>N</sub>2 reaction, since the N–H(D) vibrational frequencies invariably increase upon going to the TS because of an increase in steric congestion in the bond-making process.<sup>3</sup> Thus, the greater the extent of bond formation, the smaller the  $k_{\text{H}}/k_{\text{D}}$  value becomes.

We have reported the DKIEs for the reactions of various substrates<sup>4</sup> with deuterated amines to clarify the reaction mechanism. Some special results we have obtained are as follows: (i) the max. value of primary normal DKIE,  $k_{\text{H}}/k_{\text{D}} = 2.58$ , for the reactions of benzyl benzenesulfonates with deuterated anilines in MeOH.<sup>5</sup> (ii) The min. value of secondary inverse

DKIE,  $k_{\text{H}}/k_{\text{D}} = 0.739$ , for the reactions of phenylchloroformates with deuterated anilines in MeCN.<sup>6</sup> (iii) The max. values of primary normal DKIEs,  $k_{\text{H}}/k_{\text{D}} \approx 2.7$ , for the nucleophilic addition reactions of deuterated benzylamines to benzylidenemalonitriles,<sup>7</sup>  $\beta$ -nitrostilbene<sup>8</sup> and ethyl- $\alpha$ -cyanocinnamates<sup>9</sup> in MeCN. (iv) The min. value of secondary inverse DKIE,  $k_{\text{H}}/k_{\text{D}} = 0.803$ , for the reactions of *N*-methyl-*N*-phenylcarbamoyl chlorides with deuterated benzylamines in MeCN.<sup>10</sup> (v) The secondary normal  $\beta$ -type-DKIEs,  $k_{\text{H}}/k_{\text{D}} > 1$ , when the rate-determining step is the breakdown of the intermediate:  $k_{\text{H}}/k_{\text{D}} = 1.03\text{--}1.11$  for the reactions of phenylacetyl chlorides with deuterated anilines in MeCN,<sup>11</sup>  $k_{\text{H}}/k_{\text{D}} = 1.04\text{--}1.12$  for the reactions of 4-nitrophenyl *N*-phenylcarbamates with deuterated benzylamines in MeCN<sup>12</sup> and  $k_{\text{H}}/k_{\text{D}} = 1.02\text{--}1.11$  for the reactions of benzhydryl chlorides with deuterated pyrrolidine in MeCN.<sup>13</sup> The obtained order of 1.1 is consistent with the typical value of secondary normal  $\beta$ -DKIEs.<sup>14</sup>

Cross-interaction constants (CICs),  $\rho_{\text{XY}}$ , can be also a strong tool to provide the mechanistic criteria for the nucleophilic substitution reactions.<sup>15</sup>

$$\log(k_{\text{XY}}/k_{\text{HH}}) = \rho_{\text{X}}\sigma_{\text{X}} + \rho_{\text{Y}}\sigma_{\text{Y}} + \rho_{\text{XY}}\sigma_{\text{X}}\sigma_{\text{Y}} \quad (1a)$$

$$\rho_{\text{XY}} = \partial\rho_{\text{X}}/\partial\sigma_{\text{Y}} = \partial\rho_{\text{Y}}/\partial\sigma_{\text{X}} \quad (1b)$$

Here, X and Y are the substituents of the nucleophiles and substrates, respectively. The sign and magnitude of the CICs have made it possible to correctly interpret the reaction mechanism and the degree of tightness of the TS, respectively. The sign of  $\rho_{\text{XY}}$  is normally negative in a concerted S<sub>N</sub>2 reaction (or in a stepwise reaction with rate-limiting bond formation), whereas it is positive for a stepwise reaction with a rate-limiting leaving group departure from the intermediate. The magnitude of  $\rho_{\text{XY}}$  in a concerted S<sub>N</sub>2

Department of Chemistry, Inha University, Incheon, 402-751, Korea. E-mail: hwlee@inha.ac.kr; Fax: +82-32-873-9333; Tel: +82-32-860-7681

† Electronic supplementary information (ESI) available: Synthesis of substrates, product analysis with analytical and spectroscopic data and Hammett plots. See DOI: 10.1039/b903148k

**Table 1** Summary of selectivity parameters ( $\rho_X$ ,  $\beta_X$ ,  $\rho_Y$  and  $\rho_{XY}$ ), deuterium kinetic isotope effects ( $k_H/k_D$ ) and proposed mechanisms for the reactions of  $R_1R_2P(=A)$ ; A = O or S)Cl with  $XC_6H_4NH_2(D_2)$  in acetonitrile at 55.0 °C

$A(R_1, R_2)^a$	$-\rho_X$	$\beta_X$	$\rho_Y$	$\rho_{XY}$	$k_H/k_D$	Proposed mechanism	ref.
O(YPhO,PhO)	3.42–4.63	1.24–1.68	0.22–0.87	–1.31	0.61–0.87	Concerted $S_N2$ ; late TS; mainly backside attack <sup>d</sup>	18a
S(YPhO,PhO)	3.81–4.01	1.34–1.41	0.67–0.89	–0.22	1.11–1.33	Concerted $S_N2$ ; backside and frontside attack <sup>e</sup>	18c
O(YPhO,EtO)	3.09–3.40	1.09–1.20	0.41–0.90	–0.60	1.07–1.80	Concerted $S_N2$ ; backside and frontside attack	18f
S(YPhO,EtO)	3.14–3.40	1.10–1.19	0.74–0.95	–0.28	1.06–1.27	Concerted $S_N2$ ; backside and frontside attack	18f
O(EtO,EtO)	2.99/2.80 <sup>b</sup>	1.06/0.993	—	—	0.714–0.919	Concerted $S_N2$ ; mainly backside attack	18g
S(EtO,EtO)	2.75/2.70	0.977/0.957	—	—	1.01–1.10	Concerted $S_N2$ ; backside and frontside attack	18g
O(MeO,MeO)	2.72/2.56	0.962/0.907	—	—	0.798–0.979	Concerted $S_N2$ ; backside and frontside attack	18g
S(MeO,MeO)	2.81/2.73	0.993/0.963	—	—	0.945–1.06	Concerted $S_N2$ ; backside and frontside attack	18g
S(YPhS,Ph)	3.35–3.47	1.21–1.25	0.29–0.50	–0.31	1.15–1.59	Concerted $S_N2$ ; mainly frontside attack	18h
O(Ph,Ph)	4.78/4.56	1.69/1.62	—	—	1.42–1.82	Concerted $S_N2$ ; mainly frontside attack	18d
S(Ph,Ph)	3.97/3.94	1.40/1.40	—	—	1.00–1.10	Concerted $S_N2$ ; backside and frontside attack	18e
O(Ph,Me)	2.49/2.28	0.88/0.81	—	—	1.62–2.10	Concerted $S_N2$ ; mainly frontside attack	18i
O(Me,Me) <sup>c</sup>	4.59/4.42	1.62/1.56	—	—	0.703–0.899	Concerted $S_N2$ ; mainly backside attack	18i

<sup>a</sup> The substrates,  $R_1R_2P(=O)Cl$  and  $R_1R_2P(=S)Cl$ , are denoted as  $O(R_1R_2)$  and  $S(R_1R_2)$ , respectively. <sup>b</sup> Calculated from  $k_H/k_D$  / calculated from  $k_D$ . <sup>c</sup> Selectivity parameters and DKIEs are at 15.0 °C. <sup>d</sup> Backside attack indicates that the reaction proceeds through in-line type TS I. <sup>e</sup> Frontside attack indicates that the reaction proceeds through a hydrogen-bonded, four-center-type TS II.


reaction is inversely proportional to the distance between X and Y in the TS.

There is extensive literature on phosphoryl transfer reactions and two main types of mechanisms are known to occur, stepwise ( $A_N+D_N$ ) and concerted ( $A_ND_N$ ).<sup>16</sup> The attacking direction of the nucleophile can be backside and/or frontside depending upon substrate, nucleophile, leaving group and reaction conditions.<sup>17</sup> Two possible TS structures of frontside nucleophilic attack would be apical(Nu)-equatorial(Lg) or eq(Nu)-ap(Lg) in the trigonal bipyramidal pentacoordinate (TBP-5C), while that of backside attack is ap(Nu)-ap(Lg).

In our preceding papers, we reported the anilinolysis of phosphate derivatives in MeCN, and proposed a reaction mechanism based mainly on the DKIEs and CICs.<sup>18</sup> In Y-aryl phenyl chlorophosphates [(YPhO)(PhO)P(=O)Cl],<sup>18a</sup> a concerted mechanism with a late TS (in-line type TS I in Scheme 1) was proposed on the basis of large secondary inverse DKIEs,  $k_H/k_D = 0.61–0.87$  (*vide infra*), and a large negative CIC,  $\rho_{XY} = -1.31$ . A concerted mechanism with the participation of a hydrogen bonded four-center-type TS II (Scheme 1) was proposed in Y-aryl phenyl chlorothiophosphates [(YPhO)(PhO)P(=S)Cl],<sup>18c</sup> Y-aryl ethyl chlorophosphates [(YPhO)(EtO)P(=O)Cl]<sup>18f</sup> and Y-aryl ethyl chlorothiophosphates [(YPhO)(EtO)P(=S)Cl]<sup>18f</sup> on the basis of primary DKIEs and negative CICs ( $k_H/k_D = 1.11–1.33$  and  $\rho_{XY} = -0.22$ ,  $k_H/k_D = 1.07–1.80$  and  $\rho_{XY} = -0.60$ , and  $k_H/k_D = 1.06–1.27$  and  $\rho_{XY} = -0.28$ , respectively). In dimethyl chlorothiophosphate [(MeO)<sub>2</sub>P(=S)Cl],<sup>18g</sup> a linear free energy correlation and the DKIEs,  $k_H/k_D = 0.945–1.06$  (greater value for stronger nucleophile), were rationalized by a gradual TS variation

from a backside nucleophilic attack (TS I) for weakly basic anilines to a frontside attack involving a hydrogen bonded four-center-type TS II for strongly basic anilines. In dimethyl chlorophosphate [(MeO)<sub>2</sub>P(=O)Cl],<sup>18g</sup> diethyl chlorophosphate [(EtO)<sub>2</sub>P(=O)Cl]<sup>18g</sup> and dimethyl phosphinic chloride [Me<sub>2</sub>P(=O)Cl],<sup>18i</sup> the DKIEs are secondary inverse,  $k_H/k_D = 0.798–0.979$ , 0.714–0.919 and 0.703–0.899, respectively, implying a concerted mechanism dominantly through a TS I. In diethyl chlorothiophosphate [(EtO)<sub>2</sub>P(=S)Cl]<sup>18g</sup> and diphenyl thiophosphinic chloride [Ph<sub>2</sub>P(=S)Cl],<sup>18e</sup> the DKIEs are relatively small primary normal,  $k_H/k_D = 1.01–1.10$  and 1.00–1.10, respectively, implying a concerted mechanism through TS I and II. In diphenyl phosphinic chloride [Ph<sub>2</sub>P(=O)Cl],<sup>18d</sup> methyl phenyl phosphinic chloride [MePhP(=O)Cl]<sup>18i</sup> and Y-S-aryl phenyl phosphonochloridothioates [Ph(YPhS)P(=S)Cl],<sup>b</sup> the DKIEs are relatively large primary normal,  $k_H/k_D = 1.42–1.82$ , 1.62–2.10 and 1.15–1.59, respectively, implying a concerted mechanism predominantly through TS II. These results are summarized in Table 1 together with selectivity parameters ( $\rho_X$ ,  $\beta_X$ ,  $\rho_Y$  and  $\rho_{XY}$ ), DKIEs involving deuterated anilines and proposed mechanisms.


We have also reported on the pyridinolysis of phosphate derivatives in MeCN.<sup>19</sup> In Y-aryl phenyl chlorophosphates,<sup>19a</sup> a concerted mechanism with an early TS (in-line type) was proposed on the basis of a linear free energy correlation and small negative CICs,  $\rho_{XY} = -0.15$  and  $\beta_{XY} = -0.01$ .<sup>20</sup> However, in Z-aryl bis(4-methoxyphenyl) phosphates [(4-MeOPhO)<sub>2</sub>P(=O)OPhZ],<sup>19b</sup> free energy correlations show biphasic concave downwards in the variation of the basicities of both nucleophiles and leaving groups. The CICs,  $\rho_{XZ}$ , and Brønsted coefficients,  $\beta_X (= \beta_{nuc})$  and  $\beta_Z (= \beta_{lg})$ , showed that the mechanism (concerted or stepwise with rate-limiting formation of a TBP-5C intermediate) and attacking direction (backside or frontside) of pyridines are depending on both substituents in the nucleophiles (X) and leaving groups (Z). Furthermore, in Y-aryl phenyl isothiocyanophosphates [(YPhO)(PhO)P(=O)NCS],<sup>19c</sup> the Hammett plots show biphasic concave downwards in the variation of the substituents in the substrates (Y), while concave upwards in the variation of the basicities of the nucleophiles. We proposed the mechanistic change at  $\sigma_Y = 0$  from concerted to stepwise with rate-limiting expulsion



**Scheme 1** Proposed TS structures (A = O or S; L = H or D).

of the leaving group from a TBP-5C intermediate. A frontside attack TS for more basic nucleophiles was proposed.

Herein, we report an examination of the DKIEs on the aminolysis of *Y-O*-aryl methyl phosphonochloridothioates with *X*-anilines in MeCN at 55.0 °C (Scheme 2) to clarify the phosphoryl transfer mechanism and to rationalize the substituent effects of the nucleophiles and substrates on DKIEs.



**Scheme 2** Studied reaction systems.

## Results and discussion

The observed pseudo-first-order rate constants ( $k_{\text{obsd}}$ ) were found to follow eqn (2) for all of the reactions under pseudo-first-order conditions with a large excess of aniline nucleophiles.

$$k_{\text{obsd}} = k_0 + k_{\text{H(D)}}[\text{XC}_6\text{H}_4\text{NH}_2(\text{D}_2)] \quad (2)$$

The  $k_0$  values were negligible ( $k_0 = 0$ ) in acetonitrile. The second-order rate constants ( $k_{\text{H}}$  and  $k_{\text{D}}$ ) were obtained for at least five aniline concentrations ( $[\text{XC}_6\text{H}_4\text{NH}_2(\text{D}_2)]$ ). The linear plots of eqn (2) suggest that there are no base-catalysis or noticeable side reactions and that the overall reaction is described by Scheme 2.

Second-order rate constants and selectivity parameters ( $\rho_{\text{X}}$ ,  $\beta_{\text{X}}$  and  $\rho_{\text{Y}}$ ) are summarized in Tables 2 and 3, respectively. The  $\text{p}K_{\text{a}}$  values of anilines in water are used to obtain the Brønsted  $\beta_{\text{X}}$  values and are justified experimentally and theoretically.<sup>23</sup> The  $\text{p}K_{\text{a}}$  and  $\sigma$  values of deuterated anilines are assumed to be the same as those of anilines. The rate increases with a more electron-withdrawing substituent (Y) in the substrate and with a more electron-donating substituent (X) in the nucleophile, consistent with a typical nucleophilic substitution reaction with negative charge development at the reaction center P atom and with positive charge development at the nucleophilic N atom in the TS.

It needs to be stressed that Hammett (Fig. S1†:  $\log k_{\text{H}}$  vs.  $\sigma_{\text{X}}$  and  $\log k_{\text{D}}$  vs.  $\sigma_{\text{X}}$ ) and Brønsted (Fig. 1:  $\log k_{\text{H}}$  vs.  $\text{p}K_{\text{aX}}$  and  $\log k_{\text{D}}$  vs.  $\text{p}K_{\text{aX}}$ ) plots are biphasic concave downwards with a “discrete” break region between X = H and 4-Cl in the variation of the nucleophiles. The biphasic concave downwards nonlinear

**Table 2** Second-order rate constants ( $k_{\text{H(D)}} \times 10^4/\text{M}^{-1} \text{s}^{-1}$ ) for the reactions of *Y-O*-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2(\text{D}_2)$  in acetonitrile at 55.0 °C


X \ Y		4-MeO	4-Me	H	3-Cl	4-CN
4-MeO	$k_{\text{H}}$	81.2 ± 1.8 <sup>a</sup>	94.1 ± 1.6	101 ± 2	152 ± 2	244 ± 4
	$k_{\text{D}}$	68.8 ± 1.1	78.9 ± 1.8	81.9 ± 1.4	122 ± 2	188 ± 2
4-Me	$k_{\text{H}}$	77.2 ± 1.4	80.2 ± 1.8	95.4 ± 2.8	118 ± 3	163 ± 2
	$k_{\text{D}}$	67.3 ± 0.9	68.6 ± 1.1	79.7 ± 1.3	97.5 ± 2.2	126 ± 2
H	$k_{\text{H}}$	64.4 ± 1.3	67.2 ± 1.5	72.5 ± 2.1	83.0 ± 1.5	108 ± 2
	$k_{\text{D}}$	62.4 ± 1.3	63.0 ± 1.5	64.9 ± 1.9	71.7 ± 1.4	85.9 ± 1.5
4-Cl	$k_{\text{H}}$	51.5 ± 0.3	55.0 ± 1.0	58.0 ± 1.2	65.2 ± 1.6	69.0 ± 1.3
	$k_{\text{D}}$	105 ± 2	111 ± 3	115 ± 1	141 ± 3	188 ± 3
3-Cl	$k_{\text{H}}$	10.3 ± 0.2	12.2 ± 0.3	15.0 ± 0.4	20.4 ± 0.4	25.0 ± 0.3
	$k_{\text{D}}$	23.2 ± 0.3	24.9 ± 0.4	29.8 ± 0.9	40.1 ± 0.6	49.0 ± 0.6
3-NO <sub>2</sub>	$k_{\text{H}}$	0.242 ± 0.005	0.275 ± 0.007	0.322 ± 0.009	0.493 ± 0.009	0.800 ± 0.016
	$k_{\text{D}}$	0.605 ± 0.014	0.653 ± 0.014	0.750 ± 0.010	0.958 ± 0.024	1.41 ± 0.03

<sup>a</sup> Standard deviation.

**Table 3** Selectivity parameters<sup>a</sup> ( $\rho_{\text{X}}$ ,  $\beta_{\text{X}}$  and  $\rho_{\text{Y}}$ ) for the reactions of *Y-O*-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2(\text{D}_2)$  in acetonitrile at 55.0 °C

X	4-MeO	4-Me	H	4-Cl	3-Cl	3-NO <sub>2</sub>
$-\rho_{\text{Y(H)}}^{b,c}$	0.49 ± 0.04 <sup>f</sup>	0.34 ± 0.02	0.23 ± 0.03	0.13 ± 0.01	0.40 ± 0.03	0.54 ± 0.03
$-\rho_{\text{Y(D)}}^{d,e}$	0.45 ± 0.04	0.29 ± 0.02	0.14 ± 0.02	0.26 ± 0.03	0.35 ± 0.01	0.38 ± 0.03
X = 4-MeO, 4-Me and H						
Y	4-MeO	4-Me	H	3-Cl	4-CN	
$-\rho_{\text{X(H)}}^g$	0.38 ± 0.06	0.52 ± 0.07	0.55 ± 0.12	0.97 ± 0.05	1.28 ± 0.18	
$-\rho_{\text{X(D)}}^h$	0.16 ± 0.03	0.35 ± 0.10	0.39 ± 0.11	0.85 ± 0.05	1.23 ± 0.20	
$\beta_{\text{X(H)}}^i$	0.14 ± 0.02	0.19 ± 0.03	0.20 ± 0.03	0.35 ± 0.04	0.46 ± 0.09	
$\beta_{\text{X(D)}}^j$	0.06 ± 0.01	0.12 ± 0.04	0.14 ± 0.03	0.31 ± 0.03	0.44 ± 0.10	
X = 4-Cl, 3-Cl and 3-NO <sub>2</sub>						
Y	4-MeO	4-Me	H	3-Cl	4-CN	
$-\rho_{\text{X(H)}}^k$	4.84 ± 0.05	4.80 ± 0.04	4.74 ± 0.16	4.48 ± 0.26	4.10 ± 0.28	
$-\rho_{\text{X(D)}}^l$	4.66 ± 0.01	4.65 ± 0.01	4.58 ± 0.12	4.56 ± 0.20	4.45 ± 0.08	
$\beta_{\text{X(H)}}^m$	1.54 ± 0.01	1.53 ± 0.03	1.51 ± 0.07	1.43 ± 0.10	1.31 ± 0.11	
$\beta_{\text{X(D)}}^n$	1.49 ± 0.02	1.48 ± 0.02	1.46 ± 0.06	1.45 ± 0.08	1.42 ± 0.05	

<sup>a</sup> The  $\sigma$  and  $\text{p}K_{\text{a}}$  values in water were taken from ref. 21 and 22, respectively. <sup>b</sup> Subscript (H) indicates that the value is calculated from  $k_{\text{H}}$ . <sup>c</sup> Correlation coefficients ( $r$ ) were better than 0.980. <sup>d</sup> Subscript (D) indicates that the value is calculated from  $k_{\text{D}}$ . <sup>e</sup>  $r \geq 0.965$ . <sup>f</sup> Standard deviation. <sup>g</sup>  $r \geq 0.977$ . <sup>h</sup>  $r \geq 0.959$ . <sup>i</sup>  $r \geq 0.981$ . <sup>j</sup>  $r \geq 0.943$ . <sup>k</sup>  $r \geq 0.998$ . <sup>l</sup>  $r \geq 0.999$ . <sup>m</sup>  $r \geq 0.997$ . <sup>n</sup>  $r \geq 0.998$ .



**Fig. 1** The Brønsted plots of  $\log k_{\text{H(D)}}$  vs.  $\text{p}K_{\text{a}}(\text{X})$  for the reactions of *Y-O*-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2$  (a) and  $\text{XC}_6\text{H}_4\text{ND}_2$  (b) in acetonitrile at  $55.0^\circ\text{C}$ .

Hammett and Brønsted plots with a “continuous” break region (curvature or break point) are thought to result from a change in the rate-limiting step from the breakdown to the formation of an intermediate, as the basicity of the nucleophile increases.<sup>24</sup> At low  $\text{p}K_{\text{a}}$  values of nucleophiles, the breakdown step ( $k_{\text{b}}$ ) is rate-limiting with a steeper Brønsted slope. On the other hand, at high  $\text{p}K_{\text{a}}$  values of nucleophiles, the formation step ( $k_{\text{a}}$ ) is rate-limiting with a smaller Brønsted slope. At the center of the Brønsted curvature, a nucleophile with  $\text{p}K_{\text{a}} = \text{p}K_{\text{a}}^\circ$  has the same leaving ability from the intermediate as that of the leaving group ( $k_{-\text{a}} = k_{\text{b}}$ ). As a result, the Brønsted plot should be *continuous* with a break region or point. The *discrete* Brønsted plots are reported because of a desolvation step prior to the rate-limiting nucleophilic attack with a smaller value of  $\beta_{\text{X}}$  when the nucleophile is an anion and the solvent is polar protic, e.g., water.<sup>25</sup> However, in the present work, the much smaller values of  $\rho_{\text{X}}$  and  $\beta_{\text{X}}$  for the more basic anilines are not ascribed to a desolvation step prior to the rate-limiting nucleophilic attack, since the aniline nucleophile is neutral and the MeCN solvent is dipolar aprotic. Thus, the “discrete” break region in the present work is not ascribed to a change in the rate-limiting step in a stepwise mechanism or to a desolvation step prior to a rate-limiting nucleophilic attack.

Table 4 reveals that the DKIEs are distinctly divided into two parts, secondary inverse ( $k_{\text{H}}/k_{\text{D}} < 1$ ) for weakly basic anilines ( $\text{X} = 4\text{-Cl}$ ,  $3\text{-Cl}$  and  $3\text{-NO}_2$ ) and primary normal ( $k_{\text{H}}/k_{\text{D}} > 1$ ) for strongly basic anilines ( $\text{X} = 4\text{-MeO}$ ,  $4\text{-Me}$  and  $\text{H}$ ). The primary normal DKIE becomes systematically greater with a stronger nucleophile and with a more electron-withdrawing substituent in the substrate. As a result, the max. value of  $k_{\text{H}}/k_{\text{D}}$  is 1.30 when  $\text{X} = 4\text{-MeO}$  and  $\text{Y} = 4\text{-CN}$  while the min. value of  $k_{\text{H}}/k_{\text{D}}$  is 1.03 when  $\text{X} = \text{H}$  and  $\text{Y} = 4\text{-MeO}$ .


However, the tendencies of secondary inverse DKIEs are somewhat complicated and can be seen in two aspects: (a) substituent  $\text{X}$  in the nucleophile; (b) substituent  $\text{Y}$  in the substrate, respectively. Regarding (a), for  $\text{X} = 4\text{-Cl}$ , the  $k_{\text{H}}/k_{\text{D}}$  value becomes systematically greater with an electron-withdrawing substituent in the substrate, from  $\text{Y} = 4\text{-MeO}$  via  $4\text{-Me}$  to  $\text{H}$ , and then systematically smaller with an electron-withdrawing substituent in the substrate, from  $\text{Y} = \text{H}$  via  $3\text{-Cl}$  to  $4\text{-CN}$ . For  $\text{X} = (3\text{-Cl}$  and  $3\text{-NO}_2)$ , the  $k_{\text{H}}/k_{\text{D}}$  value becomes systematically greater with a stronger nucleophile and with an electron-withdrawing substituent in the substrate. Regarding (b), for  $\text{Y} = (4\text{-MeO}$ ,  $4\text{-Me}$  and  $\text{H})$ , the  $k_{\text{H}}/k_{\text{D}}$  value becomes systematically smaller with a weaker nucleophile and with an electron-donating substituent in the substrate, resulting in the min. value of  $k_{\text{H}}/k_{\text{D}} = 0.400$  when  $\text{X} = 3\text{-NO}_2$  and  $\text{Y} = 4\text{-MeO}$ , and the max. value of  $k_{\text{H}}/k_{\text{D}} = 0.504$  when  $\text{X} = 4\text{-Cl}$  and  $\text{Y} = \text{H}$ . For  $\text{Y} = (3\text{-Cl}$  and  $4\text{-CN})$ , the  $k_{\text{H}}/k_{\text{D}}$  value becomes systematically smaller with a stronger nucleophile and with an electron-withdrawing substituent in the substrate, resulting in the min. value of  $k_{\text{H}}/k_{\text{D}} = 0.367$  when  $\text{X} = 4\text{-Cl}$  and  $\text{Y} = 4\text{-CN}$ , and the max. value of  $k_{\text{H}}/k_{\text{D}} = 0.567$  when  $\text{X} = 3\text{-NO}_2$  and  $\text{Y} = 4\text{-CN}$ .

We propose the present trends as a mechanistic change from a stepwise mechanism with a rate-limiting leaving group departure from a TBP-5C intermediate involving a type TS I for weakly basic anilines ( $\text{X} = 4\text{-Cl}$ ,  $3\text{-Cl}$  and  $3\text{-NO}_2$ ) to a concerted mechanism involving a four-center hydrogen-bonded type TS II for strongly basic anilines ( $\text{X} = 4\text{-MeO}$ ,  $4\text{-Me}$  and  $\text{H}$ ), based on the following grounds: (i) The magnitudes of  $\rho_{\text{X}}$  and  $\beta_{\text{X}}$  are much greater (even 30 times greater for  $\rho_{\text{X}}$ ) for weaker nucleophiles ( $\rho_{\text{X}} = -4.84$  to  $-4.10$  and  $\beta_{\text{X}} = 1.54\text{--}1.31$ ) than for stronger nucleophiles ( $\rho_{\text{X}} = -1.28$  to  $-0.16$  and  $\beta_{\text{X}} = 0.46\text{--}0.14$ ) as shown in Table 2. It is well known that the magnitudes of  $\rho_{\text{X}}$  and  $\beta_{\text{X}}$  values for a stepwise reaction with a rate-limiting leaving group expulsion from the intermediate are greater than those of concerted mechanism. These are consistent with the proposed mechanism. (ii) The CICs for weaker nucleophiles are *positive* ( $\rho_{\text{XY(H)}} = +0.77$  and  $\rho_{\text{XY(D)}} = +0.21$ ; Fig. 2),<sup>26</sup> suggesting the stepwise mechanism with a rate-limiting leaving group expulsion from the intermediate. While those for stronger nucleophiles are relatively large *negative* ( $\rho_{\text{XY(H)}} = -0.95$  and  $\rho_{\text{XY(D)}} = -1.11$ ; Fig. 3), suggesting a concerted mechanism.<sup>15</sup> (iii) The DKIEs for weaker nucleophiles are *extremely large* secondary inverse ( $k_{\text{H}}/k_{\text{D}} = 0.367\text{--}0.567$ ; *unprecedentedly small* values), suggesting very severe steric crowding in the TS due to backside nucleophilic attack through in-line type TS I. While those for stronger nucleophiles are primary normal ( $k_{\text{H}}/k_{\text{D}} = 1.03\text{--}1.30$ ), implying frontside nucleophilic attack through a hydrogen bonded


**Table 4** Deuterium kinetic isotope effects ( $k_{\text{H}}/k_{\text{D}}$ ) for the reactions of *Y-O*-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2(\text{D}_2)$  in acetonitrile at  $55.0^\circ\text{C}$

X/Y	4-MeO	4-Me	H	3-Cl	4-CN
4-MeO	$1.18 \pm 0.03^a$	$1.19 \pm 0.03$	$1.23 \pm 0.04$	$1.25 \pm 0.03$	$1.30 \pm 0.03$
4-Me	$1.15 \pm 0.03$	$1.17 \pm 0.03$	$1.20 \pm 0.04$	$1.21 \pm 0.04$	$1.29 \pm 0.03$
H	$1.03 \pm 0.03$	$1.08 \pm 0.03$	$1.12 \pm 0.05$	$1.16 \pm 0.03$	$1.26 \pm 0.03$
4-Cl	$0.490 \pm 0.011$	$0.495 \pm 0.016$	$0.504 \pm 0.011$	$0.462 \pm 0.015$	$0.367 \pm 0.009$
3-Cl	$0.444 \pm 0.009$	$0.490 \pm 0.014$	$0.503 \pm 0.020$	$0.509 \pm 0.013$	$0.510 \pm 0.009$
3-NO <sub>2</sub>	$0.400 \pm 0.013$	$0.421 \pm 0.014$	$0.429 \pm 0.014$	$0.515 \pm 0.016$	$0.567 \pm 0.015$

<sup>a</sup> Standard error.



**Fig. 2** The  $\rho_{XY}$  ( $= \partial\rho_X/\partial\sigma_Y = \partial\rho_Y/\partial\sigma_X$ ) plots of  $\rho_X$  vs.  $\sigma_Y$  and  $\rho_Y$  vs.  $\sigma_X$  for the reactions of Y-O-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2$  (a) and  $\text{XC}_6\text{H}_4\text{ND}_2$  (b), when X = 4-Cl, 3-Cl and 3- $\text{NO}_2$  in acetonitrile at 55.0 °C. The obtained values by multiple regressions are  $\rho_{\text{XY(H)}} = +0.77 \pm 0.17$  ( $r = 0.998$ ) and  $\rho_{\text{XY(D)}} = +0.21 \pm 0.11$  ( $r = 0.999$ ).



**Fig. 3** The  $\rho_{XY}$  plots of  $\rho_X$  vs.  $\sigma_Y$  and  $\rho_Y$  vs.  $\sigma_X$  for the reactions of Y-O-aryl methyl phosphonochloridothioates with  $\text{XC}_6\text{H}_4\text{NH}_2$  (a) and  $\text{XC}_6\text{H}_4\text{ND}_2$  (b), when X = 4-MeO, 4-Me and H in acetonitrile at 55.0 °C. The obtained values by multiple regressions are  $\rho_{\text{XY(H)}} = -0.95 \pm 0.17$  ( $r = 0.980$ ) and  $\rho_{\text{XY(D)}} = -1.11 \pm 0.16$  ( $r = 0.976$ ).

four-center type TS II. (iv) As shown in Fig. 1b, the fall-off rate, from X = H (weaker aniline nucleophile) to X = 4-Cl (stronger aniline nucleophile), strongly suggests that a frontside attack is more favorable than backside for stronger aniline nucleophiles.

Let us examine the proposed mechanism on the basis of the steric effects and DKIEs, and the substituent effects of the nucleophiles and substrates on DKIEs in the cases of weaker and stronger nucleophiles, separately.

### Secondary inverse DKIEs for weaker nucleophiles (X = 4-Cl, 3-Cl and 3- $\text{NO}_2$ )

We reported that the second-order rate constants ( $k_{\text{H}}$  with  $\text{C}_6\text{H}_5\text{NH}_2$ ) for the anilinolysis of three phosphinates,  $\text{Me}_2\text{P}(=\text{O})\text{Cl}$ ,<sup>18i</sup>  $\text{MePhP}(=\text{O})\text{Cl}$ <sup>18i</sup> and  $\text{Ph}_2\text{P}(=\text{O})\text{Cl}$ <sup>18d</sup> gave relative rates of 4,520 ( $k_{\text{H}} = 7.82$ )<sup>27</sup>: 80 ( $k_{\text{H}} = 0.138$ ): 1 ( $k_{\text{H}} = 0.00173 \text{ M}^{-1} \text{ s}^{-1}$ ) in MeCN at 55.0 °C. Buncel and coworkers reported that the second-order rate constants for the ethanolysis of three phosphinates,  $\text{Me}_2\text{P}(=\text{O})(\text{OPh-4-NO}_2)$ ,  $\text{MePhP}(=\text{O})(\text{OPh-4-NO}_2)$  and  $\text{Ph}_2\text{P}(=\text{O})(\text{OPh-4-NO}_2)$  gave relative rates of 235 ( $k_{\text{EtO}^-} = 230$ ): 69 ( $k_{\text{EtO}^-} = 67.6$ ): 1 ( $k_{\text{EtO}^-} = 0.980 \text{ M}^{-1} \text{ s}^{-1}$ ) in anhydrous ethanol at 25.0 °C.<sup>28</sup> Williams and coworkers reported that the second-order rate constants for the phosphate catalyzed


hydrolysis of two phosphinates,  $\text{Me}_2\text{P}(=\text{O})(\text{OPh-4-NO}_2)$  and  $\text{Ph}_2\text{P}(=\text{O})(\text{OPh-4-NO}_2)$  gave the relative rate of 52 ( $k_{\text{HPO}_4^{2-}} = 1.06 \times 10^{-2}$ ): 1 ( $k_{\text{HPO}_4^{2-}} = 2.05 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ ) in 10% dioxane-aqueous 0.1M NaCl at 25.0 °C.<sup>29</sup> These results are contrary to expectations for the electronic influence of the ligands: Ph ( $\sigma_1 = 0.12$ ) and Me ( $\sigma_1 = -0.01$ ) groups;  $\text{Ph}_2$  ( $\sum\sigma_1 = 0.24$ );  $\text{MePh}$  ( $\sum\sigma_1 = 0.11$ );  $\text{Me}_2$  ( $\sum\sigma_1 = -0.02$ ).<sup>30</sup> A plot of  $\log k_{\text{H}}$  (for the unsubstituted aniline) against the Taft's steric constants according to eqn (3) for the anilinolysis of three phosphinates gives the sensitivity coefficient of the steric effects  $\delta = 0.737$  ( $r = 0.999$ ), where  $\sum E_s$  is the sum of the Taft's steric constants [ $\sum E_s = 0.00$  ( $\text{Me}_2$ ),  $-2.48$  ( $\text{MePh}$ ) and  $-4.96$  ( $\text{Ph}_2$ ), with  $E_s = 0.00$  ( $\text{Me}$ ) and  $-2.48$  ( $\text{Ph}$ )].<sup>31</sup> When the same values of  $\sum E_s$  as in the anilinolysis of three phosphinates are used,  $\delta = 0.478$  ( $r = 0.953$ : roughly linear) and  $\delta = 0.345$  are obtained for the ethanolysis of three phosphinates and for the hydrolysis of two phosphinates, respectively. These results suggest that the phosphoryl transfer reaction rates are predominantly dependent on steric effects over the inductive effects of the ligands in the reaction series. The greater influence of steric effects is observed for anilinolysis than for ethanolysis (and hydrolysis), due to the larger size of the aniline nucleophile compared to the ethanol (and water) nucleophile.

$$\log k = \delta \sum E_s + C \quad (3)$$

As mentioned earlier, the smallest value of the secondary inverse DKIE that we have ever obtained was  $k_{\text{H}}/k_{\text{D}} = 0.61$  (*vide supra*) for the reaction of 4-chlorophenyl phenyl chlorophosphate with 4-chloroaniline.<sup>18a</sup> In Y-aryl 4-chlorophenyl chlorophosphates [(YPhO)(4-ClPhO)P(=O)Cl],<sup>18b</sup> a large secondary inverse DKIE,  $k_{\text{H}}/k_{\text{D}} = 0.64$ , was observed when X = 4-Cl and Y = 4-Me. These small values of secondary inverse DKIEs imply that steric hindrance plays an important role in the TS.

Comparing the DKIEs of P=S systems,  $\text{R}_1\text{R}_2\text{P}(=\text{S})\text{Cl}$ , the values of  $k_{\text{H}}/k_{\text{D}}(\text{R}_1, \text{R}_2)$  are as follows: 1.15–1.59 (YPhS, Ph)<sup>18b</sup> > 1.10–1.46 (YPhO, 4-ClPhO)<sup>18c</sup> > 1.11–1.33 (YPhO, PhO)<sup>18c</sup> > 1.06–1.27 (YPhO, EtO)<sup>18f</sup> > 1.00–1.10 (Ph, Ph)<sup>18e</sup>  $\approx$  1.01–1.10 (EtO, EtO)<sup>18g</sup> > 0.945–1.06 (MeO, MeO).<sup>18g</sup> The sequence of the  $k_{\text{H}}/k_{\text{D}}$  values shows that the DKIEs become greater as the sizes of the ligands become larger. Thus, backside nucleophilic attack becomes less favorable while frontside attack becomes more favorable as the sizes of the ligands become larger because of the increment of steric hindrance. In the present work, the two ligands of YPhO and Me, a subtle combination of large and small, resulted in two distinct pathways, frontside and backside attack for strongly basic and weakly basic anilines, respectively.


In the present work, the overall second-order rate constant can be given by  $k_{\text{H(D)}} = (k_{\text{a}}/k_{-\text{a}})k_{\text{b}} = Kk_{\text{b}}$  in which the breakdown of the intermediate is the rate-limiting.



Since the equilibrium step,  $K$ , results in an almost negligible DKIE due to the cancellation of the  $k_{\text{a}}$  and  $k_{-\text{a}}$  terms, the observed  $k_{\text{H}}/k_{\text{D}}$  reflects the leaving group expulsion rate constant,  $k_{\text{b}}$ , from the intermediate.<sup>2f</sup> As a result, the secondary normal  $\beta$ -type-DKIEs,  $k_{\text{H}}/k_{\text{D}} > 1$ , can be expected.<sup>11–14, 32, 33</sup> However, the observed DKIEs are secondary inverse, completely contrary to our

expectations, and furthermore *extremely large secondary inverse*,  $k_{\text{H}}/k_{\text{D}} = 0.367\text{--}0.567 \ll 1$ .

The obtained secondary inverse DKIEs are mainly ascribed to a rate-limiting leaving group departure step,  $k_{\text{b}}$ . The observed *unprecedentedly small*  $k_{\text{H}}/k_{\text{D}}$  values should be accompanied with great extensive bond formation and severe steric congestion in the breakdown of the intermediate. Thus, the intermediate should be reactant-like and the TS2 should be very late and product-like (as shown in Scheme 3), resulting in  $k_{\text{H}}/k_{\text{D}} = 0.367\text{--}0.567 \ll 1$  over the secondary normal  $\beta$ -type-DKIEs. The steric congestion is thus, very severe.



**Scheme 3** Intermediate and TS 2 for the reactions of Y-O-aryl methyl phosphonochloridothioates with weakly basic anilines.

The sequence of  $k_{\text{H}}/k_{\text{D}}$  values depending on Y is almost in agreement with the expectations,  $Y = 4\text{-MeO} < 3\text{-Cl} < \text{H} < 4\text{-Me} < 4\text{-CN}$  (except  $X = 4\text{-Cl}$ ), and  $\partial\sigma_{\text{Y}} > 0$  leads to  $\partial\rho_{\text{X}} > 0$  or  $|\partial\rho_{\text{X}}| < 0$ , *i.e.*,  $\rho_{\text{XY}} > 0$ . However, the observed sequence of  $k_{\text{H}}/k_{\text{D}}$  values depending on X is divided into two parts:  $X = 3\text{-NO}_2 < 3\text{-Cl} < 4\text{-Cl}$  for  $Y = (4\text{-MeO}, 4\text{-Me}, \text{H})$ , whereas  $X = 4\text{-Cl} < 3\text{-Cl} < 3\text{-NO}_2$  for  $Y = (3\text{-Cl}, 4\text{-CN})$ . This implies that a weaker nucleophile approaches to a closer proximity to the reaction center for an electron-donating Y (min.  $k_{\text{H}}/k_{\text{D}} = 0.400$  when  $X = 3\text{-NO}_2$  and  $Y = 4\text{-MeO}$ ), while a stronger nucleophile approaches a closer proximity to the reaction center for an electron-withdrawing Y (min.  $k_{\text{H}}/k_{\text{D}} = 0.367$  when  $X = 4\text{-Cl}$  and  $Y = 4\text{-CN}$ ).

#### Primary normal DKIEs for stronger nucleophiles (X = 4-MeO, 4-Me and H)

The observed primary normal DKIE may be proportional to the extent of hydrogen bond formation. A greater degree of deprotonation would occur with a greater extent of bond formation. Considering only the electrostatic interactions, greater deprotonation would occur when the substituent of the aniline has greater electron-withdrawing abilities and the substituent of the substrate has greater electron-donating abilities, resulting in the reverse of the obtained sequence. Thus, both the nucleophilicity of aniline and the electrophilicity of the substrate exceed the electrostatic term of the substituents to determine the degree of deprotonation in the present work. Thus, the stronger nucleophile ( $\partial\sigma_{\text{X}} < 0$ ) leads to a greater extent of hydrogen bond formation ( $\partial\rho_{\text{Y}} > 0$ ), *i.e.*,  $\rho_{\text{XY}} < 0$ , and the expected sequence of the  $k_{\text{H}}/k_{\text{D}}$  values is  $X = 4\text{-MeO} > 4\text{-Me} > \text{H}$ , consistent with the observed sequence. In the case of the substituent Y of the substrate, the extent of hydrogen bond formation would be greater ( $\partial\rho_{\text{X}} < 0$  or  $|\partial\rho_{\text{X}}| > 0$ ) when the electron-withdrawing ability of Y is greater ( $\partial\sigma_{\text{Y}} > 0$ ), *i.e.*,  $\rho_{\text{XY}} < 0$ . The obtained sequence of  $k_{\text{H}}/k_{\text{D}}$  values,  $Y = 4\text{-CN} > 3\text{-Cl} > \text{H} > 4\text{-Me} > 4\text{-MeO}$ , is in line with the extent of bond formation. The reaction exhibited a large negative CICs,  $\rho_{\text{XY(H)}} = -0.95$  and  $\rho_{\text{XY(D)}} = -1.11$ , suggesting that the degree of

bond formation is extensive in the TS. The TS is believed to involve a four-center hydrogen-bonded type TS II, the result of frontside attack. The four-center TS is seen to be more stable by forming a stronger hydrogen bond for a stronger nucleophile.

It should be noted that the real primary normal DKIE due to the hydrogen bond between the hydrogen of the N-H(D) moiety and the Cl leaving group is greater than the observed value, since the other hydrogen of the N-H(D) moiety yields the secondary inverse DKIE.

## Experimental section

### Materials

HPLC-grade MeCN was used for kinetic studies without further purification. Anilines were redistilled or recrystallized before use as previously described.<sup>18</sup> Deuterated anilines were prepared by heating anilines with  $\text{D}_2\text{O}$  at  $85^\circ\text{C}$  for 72 h, with one drop of HCl added as a catalyst. After numerous attempts, the anilines were deuterated more than 98%, as confirmed by  $^1\text{H}$  NMR.

### Kinetic procedure

Pseudo-first-order rate constants ( $k_{\text{obsd}}$ ) were measured conductometrically at  $55.0 \pm 0.1^\circ\text{C}$  and determined by curve fitting analysis in an origin program. The conductivity bridge was a self-made computer automated A/D converter one. [Y-O-aryl methyl phosphonochloridothioates]  $\cong 3 \times 10^{-3}$  M and [X-C<sub>6</sub>H<sub>4</sub>NH<sub>2</sub>(D<sub>2</sub>)] = 0.1–0.5 M were employed. The  $k_{\text{H(D)}}$  values are the averages of more than three runs and reproducible to within  $\pm 3\%$ .

## Conclusions

The kinetics and mechanism of the reactions of Y-O-aryl methyl phosphonochloridothioates with substituted X-anilines and deuterated X-anilines are investigated in MeCN at  $55.0^\circ\text{C}$ . In the case of weakly basic anilines, DKIEs are unusually large secondary inverse and the sign of CICs is positive. On the contrary, in the case of strongly basic anilines, DKIEs are primary normal and the sign of CICs is negative. A stepwise mechanism with a rate-limiting leaving group departure from the intermediate, through backside nucleophilic attack in-line-type TS I, is proposed for weakly basic anilines, while a concerted mechanism through a hydrogen bonded, four-center-type TS II is proposed for strongly basic anilines. We know that free energy correlation of biphasic concave downwards could be ascribed to the change of the reaction mechanism from stepwise to concerted as the basicity of the nucleophile increases. We have found that the secondary inverse DKIE could be as small as or equal to, 0.4.

## Acknowledgements

This work was financially supported by the Korea Research Foundation (KRF-2008-314-C00207).

## References and notes

- 1 L. Melander, Jr. and W. H. Saunders, *Reaction Rates of Isotopic Molecules*, Wiley-Interscience, New York, 1980.
- 2 (a) I. Lee, *Chem. Soc. Rev.*, 1995, **24**, 223; (b) J. F. Marlier, *Acc. Chem. Res.*, 2001, **34**, 283; (c) K. C. Westaway, *Adv. Phys. Org. Chem.*, 2006, **41**,

- 217; (d) S. M. Villano, S. Kato and V. M. Bierbaum, *J. Am. Chem. Soc.*, 2006, **128**, 736; (e) S. Gronert, A. E. Fajin and L. Wong, *J. Am. Chem. Soc.*, 2007, **129**, 5330; (f) I. Lee, and D. D. Sung, in *The Chemistry of Anilines*, ed. Z. Rappoport, John Wiley & Sons Ltd, Chichester, 2007, ch. 10.
- 3 (a) R. A. Poirier, W. Youliang and K. C. Westaway, *J. Am. Chem. Soc.*, 1994, **116**, 2526; (b) H. Yamata, T. Ando, S. Nagase, M. Hanamusa and K. Morokuma, *J. Org. Chem.*, 1984, **49**, 631; (c) X. G. Xhao, S. C. Tucker and D. G. Truhlar, *J. Am. Chem. Soc.*, 1991, **113**, 826.
- 4 Here, the DKIEs of various substrates involving deuterated amines (aniline, benzylamine and pyrrolidine) are introduced, and those of phosphoryl transfer reactions involving deuterated anilines are briefly introduced.
- 5 I. Lee, H. J. Koh, B. S. Lee and H. W. Lee, *J. Chem. Soc., Chem. Commun.*, 1990, 335.
- 6 K. H. Yew, H. J. Koh, H. W. Lee and I. Lee, *J. Chem. Soc., Perkin Trans. 2*, 1995, 2263.
- 7 H. K. Oh, J. H. Yang, H. W. Lee and I. Lee, *J. Org. Chem.*, 2000, **65**, 2188.
- 8 H. K. Oh, T. S. Kim, H. W. Lee and I. Lee, *J. Chem. Soc., Perkin Trans. 2*, 2002, 282.
- 9 H. K. Oh, J. H. Yang, Y. H. Hwang, H. W. Lee and I. Lee, *Bull. Korean Chem. Soc.*, 2002, **23**, 221.
- 10 H. J. Koh, H. W. Lee and I. Lee, *Bull. Korean Chem. Soc.*, 1996, **17**, 712.
- 11 H. W. Lee, J. W. Lee, H. J. Koh and I. Lee, *Bull. Korean Chem. Soc.*, 1998, **19**, 642.
- 12 H. J. Koh, O. S. Kim, H. W. Lee and I. Lee, *J. Phys. Org. Chem.*, 1997, **10**, 725.
- 13 S. Chang, H. J. Koh, B. S. Lee and I. Lee, *J. Org. Chem.*, 1995, **60**, 7760.
- 14 The typical order of 1.1 is for C–H(D), but our result is for N–H(D): T. H. Lowry and K. S. Richardson, *Mechanism and Theory in Organic Chemistry*, 3rd edn, Harper and Row, New York, p. 239.
- 15 (a) I. Lee, *Chem. Soc. Rev.*, 1990, **19**, 317; (b) I. Lee, *Adv. Phys. Org. Chem.*, 1992, **27**, 57; (c) I. Lee and H. W. Lee, *Collect. Czech. Chem. Commun.*, 1999, **64**, 1529.
- 16 (a) R. F. Hudson, in *Structure and Mechanism in Organophosphorus Chemistry*, Academic Press, London, 1965, ch. 3; (b) A. J. Kirby and A. G. Varvoglis, *J. Am. Chem. Soc.*, 1967, **89**, 415; (c) F. H. Westheimer, *Acc. Chem. Res.*, 1968, **1**, 70; (d) G. R. J. Thatcher and R. Kluger, *Adv. Phys. Org. Chem.*, 1989, **25**, 99; (e) A. Williams, in *Concerted Organic and Bio-organic Mechanisms*, CRC Press, Boca Raton, 2000, ch. 6; (f) A. C. Hengge, *Adv. Phys. Org. Chem.*, 2005, **40**, 49; (g) A. C. Hengge and I. Onyido, *Curr. Org. Chem.*, 2005, **9**, 61; (h) K. C. Kumara Swamy and N. Satish Kumar, *Acc. Chem. Res.*, 2006, **39**, 324; (i) I. H. Um, J. Y. Hong and E. Buncel, *Chem. Commun.*, 2001, 27; (j) F. Terrier, A. P. Guevel, A. P. Chatrousse, G. Moutiers and E. Buncel, *Chem. Commun.*, 2003, 600.
- 17 (a) C. R. Hall and T. D. Inch, *Tetrahedron*, 1980, **36**, 2059; (b) R. Rowell and D. G. Gorenstein, *J. Am. Chem. Soc.*, 1981, **103**, 5894; (c) R. J. P. Corriu, J. P. Dutheil and G. F. Lanneau, *J. Am. Chem. Soc.*, 1984, **106**, 1060.
- 18 (a) A. K. Guha, H. W. Lee and I. Lee, *J. Chem. Soc., Perkin Trans. 2*, 1999, 765; (b) H. W. Lee, A. K. Guha and I. Lee, *Int. J. Chem. Kinet.*, 2002, **34**, 632; (c) M. E. U. Hoque, S. Dey, A. K. Guha, C. K. Kim, B. S. Lee and H. W. Lee, *J. Org. Chem.*, 2007, **72**, 5493; (d) M. E. U. Hoque and H. W. Lee, *Bull. Korean Chem. Soc.*, 2007, **28**, 936; (e) N. K. Dey, I. S. Han and H. W. Lee, *Bull. Korean Chem. Soc.*, 2007, **28**, 2003; (f) M. E. U. Hoque, N. K. Dey, C. K. Kim, B. S. Lee and H. W. Lee, *Org. Biomol. Chem.*, 2007, **5**, 3944; (g) N. K. Dey, M. E. U. Hoque, C. K. Kim, B. S. Lee and H. W. Lee, *J. Phys. Org. Chem.*, 2008, **21**, 544; (h) B. J. Lumbiny and H. W. Lee, *Bull. Korean Chem. Soc.*, 2008, **29**, 2065; (i) N. K. Dey, M. E. U. Hoque, C. K. Kim, B. S. Lee and H. W. Lee, *J. Phys. Org. Chem.*, 2009, **22**, 425.
- 19 (a) A. K. Guha, H. W. Lee and I. Lee, *J. Org. Chem.*, 2000, **65**, 12; (b) H. W. Lee, A. K. Guha, C. K. Kim and I. Lee, *J. Org. Chem.*, 2002, **67**, 2215; (c) K. K. Adhikary, H. W. Lee and I. Lee, *Bull. Korean Chem. Soc.*, 2003, **24**, 1135; (d) M. E. U. Hoque, N. K. Dey, A. K. Guha, C. K. Kim, B. S. Lee and H. W. Lee, *Bull. Korean Chem. Soc.*, 2007, **28**, 1797; (e) K. K. Adhikary, B. J. Lumbiny, C. K. Kim and H. W. Lee, *Bull. Korean Chem. Soc.*, 2008, **29**, 851; (f) B. J. Lumbiny, K. K. Adhikary, B. S. Lee and H. W. Lee, *Bull. Korean Chem. Soc.*, 2008, **29**, 1769.
- 20  $\beta_{XY}$  can be one of the CICs.  $\beta_{XY}$  is the same as  $p_{XY}$  defined by Jencks:  $\log(k_{XY}/k_{HH}) = \beta_X pK_{aX} + \beta_Y pK_{aY} + \beta_{XY} pK_{aX} pK_{aY}$  and  $\beta_{XY} = \partial \beta_X / \partial pK_{aY} = \partial \beta_Y / \partial pK_{aX}$ : (a) D. A. Jencks and W. P. Jencks, *J. Am. Chem. Soc.*, 1977, **99**, 7948; (b) W. P. Jencks, *Chem. Rev.*, 1985, **85**, 511.
- 21 C. Hansch, A. Leo and R. W. Taft, *Chem. Rev.*, 1991, **91**, 165.
- 22 A. Streitwieser, Jr., C. H. Heathcock and E. M. Kosower, in *Introduction to Organic Chemistry*, 4th edn, Macmillan Publishing Co., New York, 1992, p. 735. The  $pK_a$ s of X-anilinium ions in water are as follows: X = 4-MeO ( $pK_a = 5.34$ ); 4-Me (5.10); H (4.60); 4-Cl (3.98); 3-Cl (3.52); 3-NO<sub>2</sub> (2.47).
- 23 (a) C. D. Ritchie, in *Solute–Solvent Interactions*, ed. J. F. Coetzee and C. D. Ritchie, Marcel Dekker, New York, 1969, ch. 4; (b) J. F. Coetzee, *Prog. Phys. Org. Chem.*, 1967, **4**, 54; (c) W. J. Spillane, G. Hogan, P. McGrath, J. King and C. Brack, *J. Chem. Soc., Perkin Trans. 2*, 1996, 2099; (d) H. K. Oh, S. Y. Woo, C. H. Shin, Y. S. Park and I. Lee, *J. Org. Chem.*, 1997, **62**, 5780.
- 24 (a) A. Williams, *Free Energy Relationships in Organic and Bio-Organic Chemistry*, RSC, Cambridge, 2003, pp. 169–172; (b) H. J. Koh, K. L. Han, H. W. Lee and I. Lee, *J. Org. Chem.*, 2000, **65**, 4706; (c) H. K. Oh, M. H. Ku, H. W. Lee and I. Lee, *J. Org. Chem.*, 2002, **67**, 3874; (d) H. K. Oh, M. H. Ku, H. W. Lee and I. Lee, *J. Org. Chem.*, 2002, **67**, 8995; (e) H. K. Oh, J. M. Lee, H. W. Lee and I. Lee, *Int. J. Chem. Kinet.*, 2004, **36**, 434; (f) H. K. Oh, J. E. Park and H. W. Lee, *Bull. Korean Chem. Soc.*, 2004, **25**, 1041; (g) E. A. Castro, P. Pavez, D. Arellano and J. G. Santos, *J. Org. Chem.*, 2001, **66**, 6571; (h) E. A. Castro, P. Pavez and J. G. Santos, *J. Org. Chem.*, 2002, **67**, 3129; (i) E. A. Castro, P. Pavez and J. G. Santos, *J. Org. Chem.*, 2002, **67**, 4494; (j) E. A. Castro, M. Angel, P. Campodonico and J. G. Santos, *J. Org. Chem.*, 2002, **67**, 8911; (k) E. Humeres, N. A. Debacher, M. M. D. Sierra, J. D. Franco and A. Shutz, *J. Org. Chem.*, 1998, **63**, 1598; (l) W. J. Spillane, P. McGrath, C. Brack and A. B. O’Byrne, *J. Org. Chem.*, 2001, **66**, 6316; (m) A. S. Baynham, F. Hibbert and M. A. Malana, *J. Chem. Soc., Perkin Trans. 2*, 1993, 1711.
- 25 (a) W. P. Jencks, S. R. Brant, J. R. Gandler, G. Fendrich and C. Nakamura, *J. Am. Chem. Soc.*, 1982, **104**, 7045; (b) I. Onyido, K. Swierczek, J. Purcell and A. C. Hengge, *J. Am. Chem. Soc.*, 2005, **127**, 7703.
- 26 (a) Some examples involving rate-limiting leaving group expulsion from the intermediate with positive  $p_{XY}$ : I. Lee, C. S. Shim and H. W. Lee, *J. Chem. Soc., Perkin Trans. 2*, 1988, 975; (b) I. Lee, C. S. Shim, S. Y. Chung, H. Y. Kim and H. W. Lee, *J. Chem. Soc., Perkin Trans. 2*, 1988, 1919; (c) I. Lee, C. S. Shim and H. W. Lee, *J. Phys. Org. Chem.*, 1989, **2**, 484; (d) I. Lee, C. S. Shim and H. W. Lee, *J. Chem. Res.*, 1992, (S)90, (M)0769; (e) H. J. Koh, C. H. Lee, H. W. Lee and I. Lee, *Bull. Korean Chem. Soc.*, 1995, **16**, 839; (f) H. J. Koh, O. S. Lee, H. W. Lee and I. Lee, *J. Phys. Org. Chem.*, 1997, **10**, 725; (g) K. S. Lee, K. K. Adhikary, H. W. Lee, B. S. Lee and I. Lee, *Org. Biomol. Chem.*, 2003, **1**, 1989; (h) ref. 24b.
- 27 The  $k_H$  value of  $7.82 \text{ M}^{-1} \text{ s}^{-1}$  at  $55.0^\circ \text{C}$  was calculated by extrapolation in the Arrhenius plot ( $r = 0.999$ ) with experimental kinetic data:  $k_H = 0.776 (0.0^\circ \text{C})$ ,  $1.01 (5.0^\circ \text{C})$  and  $1.61 \text{ M}^{-1} \text{ s}^{-1} (15.0^\circ \text{C})$  from ref. 18i.
- 28 (a) I. Onyido, K. Albright and E. Buncel, *Org. Biomol. Chem.*, 2005, **3**, 1468; (b) E. J. Dunn and E. Buncel, *Can. J. Chem.*, 1989, **67**, 1440; (c) E. J. Dunn, R. T. Moir, E. Buncel, J. G. Purdon and R. A. B. Bannard, *Can. J. Chem.*, 1990, **68**, 1837; (d) E. Buncel, K. G. Albright and I. Onyido, *Org. Biomol. Chem.*, 2004, **2**, 601.
- 29 (a) K. T. Douglas and A. Williams, *J. Chem. Soc., Perkin Trans. 2*, 1976, 515; (b) A. Williams and R. A. Naylor, *J. Chem. Soc. B*, 1971, 1967.
- 30 M. Charton, *Prog. Phys. Org. Chem.*, 1987, **16**, 287.
- 31 R. W. Taft, in *Steric Effect in Organic Chemistry*, ed. M. S. Newman, Wiley, New York, 1956, ch. 3.
- 32 Ref. 1, ch. 6.
- 33 F. M. Menger and J. H. Smith, *J. Am. Chem. Soc.*, 1972, **94**, 3824.