

Intramolecular Michael addition reaction for the synthesis of benzylbutyrolactones†

Hu He, Li-Xin Dai and Shu-Li You*

Received 25th November 2009, Accepted 28th April 2010


First published as an Advance Article on the web 20th May 2010

DOI: 10.1039/b924770j


A convenient synthesis of benzyl- γ -butyrolactone derivatives *via* intramolecular Michael addition reaction of nitro-substituted aryl allyl β -ketocarboxylates is reported. The method features simple operation, mild reaction conditions and high efficiency.

Introduction

The lignan¹ lactones have attracted considerable interest over the years because of their broad biological activities.^{2,3} Dibenzylbutyrolactones, such as enterolactone **1** and matairesinol **2**, are such examples of bioactive lignans. Recent investigations have shown that enterolactone **1** inhibits breast and colon cancer and suppresses the growth of prostate cancer cells.⁴ Matairesinol **2** and its analogs have also been demonstrated to be potent inhibitors of HIV-type 1 integrase.⁵ In addition to antitumor and antiviral activities, these compounds display phytoestrogenic,⁶ immunoregulatory,⁷ and neuroprotective properties.⁸ Despite their enormous biological applications, there have been only a few synthetic studies documented in the literature.⁹



Recently, we reported the Ir-catalyzed decarboxylative alkylation of γ -substituted allyl β -ketocarboxylates.¹⁰ Highly regio- and enantioselective allylation of ketone enolates has been achieved through Ir-catalyzed decarboxylative allylic alkylation of allyl β -ketocarboxylate (eqn (1)). Surprisingly, when 4-nitrocinnamyl 3-oxo-3-phenylpropanoate (**3a**) was tested under the same reaction conditions, no decarboxylative alkylation product was obtained but an intramolecular Michael reaction proceeded, affording γ -butyrolactone **4a**.



State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Lu, Shanghai, 200032, P. R. China. E-mail: slyou@mail.sioc.ac.cn; Fax: (+86) 21-54925087

† Electronic supplementary information (ESI) available: Experimental procedures and analysis data for new compounds. CCDC reference numbers 747012 (**4a**) and 747013 (**4j**). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b924770j

The structure of the Michael addition product was further confirmed by an X-ray crystallographic analysis (Fig. 1).¹¹ Although various α,β -unsaturated compounds have been used in the Michael addition reaction, to our knowledge, this represents a rare example of the application of an electron-deficient phenyl alkene as the Michael acceptor. Notably, a similar cyclization reaction has also been observed by Craig and coworkers during their study on the decarboxylative Claisen rearrangement reactions of diallyl 2-sulfonylmalonates; however, only moderate yield was obtained.¹² An elegant study by Tan and Chen demonstrated that Mn(OAc)₃ could promote a similar cyclization efficiently *via* a radical process.¹³ The extensive synthetic efforts towards γ -butyrolactone derivatives¹⁴ emboldened us to carry out further studies into this Michael addition reaction.





Fig. 1 X-Ray crystal structure of **4a** (thermal ellipsoids are set at 30% probability).


Further investigation of the reaction disclosed that DBU alone could catalyze the intramolecular Michael addition to afford *trans*- γ -butyrolactones (eqn (2)). Herein, we report our detailed

Table 1 Optimization of the reaction conditions for the Michael addition of **3a**^a


Entry	Solvent	Base	Base (mol%)	T/°C	t/h	Yield (%) ^b
1	CH ₂ Cl ₂	DBU	100	Reflux	2	79
2	CH ₂ Cl ₂	Et ₃ N	100	25	12	Trace
3	CH ₂ Cl ₂	Et ₃ N	100	Reflux	36	82
4	CH ₂ Cl ₂	DBN	100	25	8	77
5	CH ₂ Cl ₂	KOAc	100	25	12	NR
6	CH ₂ Cl ₂	DABCO	100	25	12	NR
7	CH ₂ Cl ₂	Cs ₂ CO ₃	100	25	12	NR
8	CH ₂ Cl ₂	K ₂ CO ₃	100	25	12	NR
9	CH ₂ Cl ₂	DIEA	100	25	12	NR
10	CH ₂ Cl ₂	BSA	100	25	12	NR
11	CH ₂ Cl ₂	TMP	100	25	12	NR
12	CH ₂ Cl ₂	PS	100	25	12	NR
13	CH ₂ Cl ₂	Quinine	100	Reflux	12	NR
14	CH ₂ Cl ₂	DBU	100	25	4	86
15	CH ₂ Cl ₂	DBU	20	Reflux	36	94
16	THF	DBU	20	Reflux	2	85
17	THF	DBU	10	Reflux	12	93
18	THF	DBU	5	Reflux	18	89
19	toluene	DBU	10	60	43	62
20	EtOH	DBU	10	60	43	48
21	DME	DBU	10	60	8	99
22	DMF	DBU	10	60	1.5	99
23	DMF	DBU	10	25	10	98

^a Reaction conditions: 0.1 M of **3a**. ^b Isolated yields.

studies on this DBU-catalyzed intramolecular Michael addition reaction.







Results and Discussion

Initially, 4-nitrocinnamyl-3-oxo-3-phenylpropanoate (**3a**) was chosen as a model substrate to optimize the reaction conditions. The results are summarized in Table 1.

In the presence of 1 equiv. of DBU, refluxing **3a** in CH₂Cl₂ for 2 h afforded the cyclized product **4a** in 79% yield (entry 1, Table 1). We first examined different bases, as no reaction occurred without DBU. Most of the bases, such as KOAc, DABCO, Cs₂CO₃, K₂CO₃, DIEA, BSA, Proton Sponge and quinine were not suitable for this transformation (entries 5–13, Table 1), only DBN, Et₃N and DBU could promote this reaction (entries 1–4, Table 1). With Et₃N, the desired product **4a** was obtained only in trace amounts at 25 °C but in 82% yield in boiling dichloromethane (entries 2–3, Table 1). In the presence of 1 equiv. of DBN at 25 °C, **4a** was isolated in 77% yield (entry 4, Table 1). In general, a relatively strong base is needed for the Michael addition reaction.

After examining the loading of DBU, an excellent yield (93%) was obtained with 10 mol% of DBU (entries 14–18, Table 1). Different solvents such as toluene, THF, EtOH, DME and DMF,

Table 2 Substrate scope for the intramolecular Michael addition reaction^a


Entry	R ¹ , R ² , 4	t/h	Yield (%) ^b
1	4-NO ₂ , Ph, 4a	10	98
2	4-NO ₂ , 4-Me-C ₆ H ₄ , 4b	36	90
3	4-NO ₂ , 4-MeO-C ₆ H ₄ , 4c	3	96
4	4-NO ₂ , 4-Cl-C ₆ H ₄ , 4d	8	96
5	4-NO ₂ , 4-Br-C ₆ H ₄ , 4e	8	93
6	4-NO ₂ , 2-naphthyl, 4f	8	99
7	4-NO ₂ , 2-furyl, 4g	8	93
8	4-NO ₂ , <i>i</i> -Pr, 4h	24	75
9 ^c	4-NO ₂ , Me, 4i	30	71
10 ^c	 4j	14	83 (4:1)
11	2-NO ₂ , Ph, 4k	12	92
12	3-NO ₂ , Ph, 4l	n.r.	—
13	2-NO ₂ , 4,5-MeO,MeO Ph, 4m	12	72
14	4-NO ₂ , 4-NO ₂ , 4n	6	76
15	 4o	10	72
16	 4p	13	78


^a Reaction conditions: 10 mol% of DBU, 0.1 M of **3** in DMF, rt. ^b Isolated yields. ^c Reaction was run at 60 °C.

were all tolerated, and the reaction at 60 °C in DMF gave **4a** in almost quantitative yield (99%) in 1.5 h (entry 22, Table 1). Notably, the reaction in DMF with 10 mol% DBU at room temperature could also give **4a** in 98% yield (entry 23, Table 1). Given the mildness of these reaction conditions, they were chosen as the optimal reaction conditions.

In the presence of 10 mol% DBU in DMF at 25 °C, various substrates were tested and the results are summarized in Table 2. Substrates bearing either electron-donating groups (Me, MeO, entries 2–3, Table 2) or electron-withdrawing groups (Cl, Br, NO₂, entries 4, 5, and 14, Table 2) on the β-aryl keto esters were well tolerated and led to their corresponding products in excellent yields (76–96%). The reaction of 2-furyl and 2-naphthyl-substituted substrates **3f** and **3g** occurred also in excellent yields (93 and 99%, entries 6–7, Table 2). Under the same conditions, substrates **3h** and **3i**, derived from β-alkyl keto esters, could also be tolerated to afford the addition products in 71–75% yields (entries 8–9, Table 2). When substrate **3j** was used, a spiro product **4j** was obtained in 83% yield. The structure of the major diastereoisomer was confirmed by X-ray crystallographic analysis.¹⁵


Moreover, 2-nitro- and 3-nitro-substituted substrates **3k** and **3l** were tested in this reaction (entries 11–12, Table 2). Substrate **3k** bearing a 2-nitro group underwent cyclization smoothly to give the corresponding product **4k** in 92% yield. However, the reaction with 3-nitro-substituted substrate **3l** failed to give the desired Michael addition product. Only starting material was recovered, even at elevated reaction temperature (60 °C). To our delight, substrates **3m** and **3p** proceeded the Michael addition reaction smoothly to give butyrolactones **4m** and **4p** in 72 and 78% yield respectively (entries 13 and 16, Table 2). The toleration of various substituents in the substrates highly broadens the utility of the current methodology. In addition to the Michael reaction for C nucleophiles, the aza-Michael reaction was explored. When substrate **3o** was used, the corresponding aza-Michael product **4o** was isolated in 72% yield (entry 15, Table 2).

A straightforward catalytic cycle was proposed as depicted in Scheme 1. Upon treatment with DBU, substrate **3a** will lead to the formation of the carbanion intermediate **I**. The strong electron-withdrawing nature of nitro group facilitates the intramolecular Michael addition generating intermediate **II**, which leads to the formation of product **4a** after the protonation and generates DBU to complete the catalytic cycle.



Scheme 1 DBU-catalyzed intramolecular Michael addition reaction.

The current methodology is particularly attractive in organic synthesis due to the versatile transformation of the nitro group. The synthetic transformation of γ -butyrolactone has also been explored. For instance, 3-benzoyl-4-(4-iodobenzyl)dihydrofuran-2(3H)-one **4aa** was prepared in three steps from **4a**, where the nitro group was converted to an iodo group (eqn (3)).



Conclusions

In summary, we have developed an efficient method for the preparation of benzyl- γ -butyrolactone derivatives through a Michael addition reaction of nitro-substituted aryl allyl β -ketoesters. The reaction features simple operation, mild reaction conditions and atom economy. The versatile transformation

of the nitro group in the products makes the current methodology particularly attractive in organic synthesis.

Experimental

General procedure for the synthesis of benzylbutyrolactone derivatives **4**

A flame dried Schlenk tube was cooled to room temperature and filled with argon. To this flask, β -ketoesters **3** (0.2 mmol), DBU (0.02 mmol, 3 mg) and DMF (2 mL) were added. The reaction mixture was stirred at room temperature or 60 °C. After the reaction was complete (monitored by TLC), the reaction mixture was diluted with Et₂O and saturated NH₄Cl. The organic layer was separated, and the aqueous layer was extracted three times with Et₂O. The combined organic layers were dried over Na₂SO₄, and concentrated to afford the crude product. The residue was purified by silica gel column chromatography to afford the desired product **4**.

Full experimental details and characterization data are given in the ESI.†

Acknowledgements

We gratefully acknowledge the National Natural Science Foundation of China (20821002, 20872159, 20932008), National Basic Research Program of China (973 Program 2009CB825300), and the Chinese Academy of Sciences for generous financial support.

References

- For reviews: (a) R. S. Ward, *Nat. Prod. Rep.*, 1999, **16**, 75; (b) M. Saleem, H. J. Kim, M. S. Ali and Y. S. Lee, *Nat. Prod. Rep.*, 2005, **22**, 696.
- (a) E. Brown and A. Daugan, *Heterocycles*, 1987, **26**, 1169; (b) T. Umezawa, L. B. Davin, E. Yamamoto, D. G. Kinston and N. G. Lewis, *J. Chem. Soc., Chem. Commun.*, 1990, 1405; (c) R. S. Ward, *Tetrahedron*, 1990, **46**, 5029; (d) M. M. Murata, M. B. M. de Azevedo and A. E. Green, *J. Org. Chem.*, 1993, **58**, 7537; (e) M. Schottner and G. Spittler, *J. Nat. Prod.*, 1998, **61**, 119; (f) A. K. Tiwari, P. V. Srinivas, S. P. Kumar and I. M. Rao, *J. Agric. Food Chem.*, 2001, **49**, 4642; (g) T. Yokoyama, M. Okano, T. Noshita, S. Funayama and K. Ohtsuki, *Biol. Pharm. Bull.*, 2003, **26**, 371; (h) J. Fischer, A. J. Reynolds, L. A. Sharp and M. S. Sherburn, *Org. Lett.*, 2004, **6**, 1345; (i) J. J. Chen, H. Y. Fang, C. Y. Duh and I. S. Chen, *Planta Med.*, 2005, **71**, 470; and references cited therein.
- (a) T. Hirano, M. Gotoh and K. Oka, *Life Sci.*, 1994, **55**, 1061; (b) M. Takasaki, T. Konoshima, K. Komatsu, H. Tokuda and H. Nishino, *Cancer Lett.*, 2000, **158**, 53; (c) F. Boccardo, G. Lunardi, P. Guglielmini, M. Parodi, R. Murialdo, G. Schettini and A. Rubagotti, *Eur. J. Cancer*, 2004, **40**, 84.
- (a) H. Adlercreutz, *Baillière's Clinical Endocrinology and Metabolism*, 1998, **12**, 605; (b) A. Stark and Z. Madar, *J. Pediatr. Endocrinol. Metab.*, 2002, **15**, 561.
- (a) E. Eich, H. Pertz, M. Kaloga, J. Schulz, M. R. Fesen, A. Mazumder and Y. Pommier, *J. Med. Chem.*, 1996, **39**, 86; (b) L.-M. Yang, S.-J. Lin, T.-H. Yang and K.-H. Lee, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 941.
- L. P. Meagher, G. R. Beecher, V. P. Flanagan and B. W. Li, *J. Agric. Food Chem.*, 1999, **47**, 3173.
- J. Y. Cho, A. R. Kim, E. S. Yoo, K. U. Baik and M. H. Park, *J. Pharm. Pharmacol.*, 1999, **51**, 1267.
- (a) Y. P. Jang, S. R. Kim and Y. C. Kim, *Planta Med.*, 2001, **67**, 470; (b) Y. P. Jang, S. R. Kim, Y. H. Choi, J. Kim, S. G. Kim, G. J. Markelonis, T. H. Oh and Y. C. Kim, *J. Neurosci. Res.*, 2002, **68**, 233.
- (a) R. S. Ward, *Recent Advances in the Chemistry of Lignans in Studies in Natural Products Chemistry, Bioactive Natural Products, Part E*, Ed. A. U. Rahman, Elsevier, Amsterdam, 2000, vol. 1, pp. 739-798; (b) H.

- Zimmer, J. Rothe and J. M. Holbert, *J. Org. Chem.*, 1960, **25**, 1234; (c) Y. Takei, K. Mori and M. Matsui, *Agric. Biol. Chem.*, 1973, **37**, 637; (d) R. S. Ward, *Chem. Soc. Rev.*, 1982, **11**, 75; (e) P. K. Agrawal and R. P. Rastogi, *Phytochemistry*, 1982, **21**, 1459; (f) J. L. Belletire and D. F. Fry, *J. Org. Chem.*, 1987, **52**, 2549; (g) M. M. A. Rahman, P. M. Dewick, D. E. Jackson and J. A. Lucas, *Phytochemistry*, 1990, **29**, 1971; (h) A. Pelter, R. S. Ward, D. M. Jones and P. Maddocks, *Tetrahedron: Asymmetry*, 1990, **1**, 857; (i) J. L. Belletire and N. O. Mahmoodi, *J. Nat. Prod.*, 1992, **55**, 194; (j) A. Van Oeveren, J. F. G. A. Jansen and B. L. Feringa, *J. Org. Chem.*, 1994, **59**, 5999; (k) J. W. Bode, M. P. Doyle, M. M. Protopopova and Q.-L. Zhou, *J. Org. Chem.*, 1996, **61**, 9146; (l) J. Brinksma, H. Van Der Deen, A. Van Oeveren and B. L. Feringa, *J. Chem. Soc., Perkin Trans. 1*, 1998, 4159; (m) D. Enders, V. Lausberg, D. L. Signore and O. M. Berner, *Synthesis*, 2002, 515; (n) D. J. Bennett, P. L. Pickering and N. S. Simpkins, *Chem. Commun.*, 2004, 1392; (o) R. C. D. Brown and N. A. Swain, *Synthesis*, 2004, 811; (p) F. Zhu, W. Li, Q. Wang and Z. Hou, *Synlett*, 2006, 1780.
- 10 H He, X.-J. Zheng, Y. Li, L.-X. Dai and S.-L. You, *Org. Lett.*, 2007, **9**, 4339.
- 11 CCDC 747012 contains the supplementary crystallographic data for **4a**. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk /data request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), and are in the ESI.† Crystal data for compound **4a**: C₁₈H₁₅NO₅, *M* = 325.31, Monoclinic, *a* = 5.8692(11) Å, α = 90°, *b* = 7.6984(14) Å, β = 94.634(3)°, *c* = 34.643(7) Å, γ = 90°, *V* = 1560.2(5) Å³, *T* = 293(2) K, space group = *P*₂₁/*c*, *Z* = 4, Number of Reflections = 8662, Independent reflections = 3360 [R(int) = 0.1004], Final *R* indices [*I* > 2 δ (*I*)] *R*₁ = 0.0607, *wR*₂ = 0.1486, *R* indices (all data) *R*₁ = 0.0919, *wR*₂ = 0.1601.
- 12 D. Craig, M. I. Lansdell and S. E. Lewis, *Tetrahedron Lett.*, 2007, **48**, 7861.
- 13 X. Tan and C. Chen, *Angew. Chem., Int. Ed.*, 2006, **45**, 4345.
- 14 (a) A. Pelter, R. S. Ward and A. Abd-el-Ghani, *J. Chem. Soc., Perkin Trans. 1*, 1996, 1353; (b) N. Kise, T. Ueda, K. Kumada, Y. Terao and N. Ueda, *J. Org. Chem.*, 2000, **65**, 464; (c) R. S. Ward and D. D. Hughes, *Tetrahedron*, 2001, **57**, 4015; (d) R. S. Ward and D. D. Hughes, *Tetrahedron*, 2001, **57**, 5633; (e) P. Eklund, A. Lindholm, J.-P. Mikkola, A. Smeds, R. Lehtila and R. Sjöholm, *Org. Lett.*, 2003, **5**, 491; (f) P. C. Eklund, F. J. Sundell, A. I. Smeds and R. E. Sjöholm, *Org. Biomol. Chem.*, 2004, **2**, 2229; (g) S. G. A. Moinuddin, B. Youn, D. L. Bedgar, M. A. Costa, G. L. Helms, C. Kang, L. B. Davin and N. G. Lewis, *Org. Biomol. Chem.*, 2006, **4**, 808; (h) M. Ghosh, *Tetrahedron*, 2007, **63**, 11710; (i) R. Singh, G. C. Singh and S. K. Ghosh, *Eur. J. Org. Chem.*, 2007, 5376.
- 15 CCDC 747013 contains the supplementary crystallographic data for major diastereomer of **4j**. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk /data request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), and are in the ESI.† Crystal data for compound **4j**: C₁₅H₁₅NO₅, *M* = 289.28, Monoclinic, *a* = 6.8274(9) Å, α = 90°, *b* = 8.8266(11) Å, β = 96.661(2)°, *c* = 23.345(3) Å, γ = 90°, *V* = 1397.3(3) Å³, *T* = 293(2) K, space group = *P*₂₁/*c*, *Z* = 4, Number of Reflections = 7130, Independent reflections = 2597 [R(int) = 0.0997], Final *R* indices [*I* > 2 δ (*I*)] *R*₁ = 0.0632, *wR*₂ = 0.1538, *R* indices (all data) *R*₁ = 0.0808, *wR*₂ = 0.1656.