

Organic & Biomolecular Chemistry

www.rsc.org/obc

Volume 8 | Number 24 | 21 December 2010 | Pages 5461–5702



ISSN 1477-0520

RSC Publishing

COMMUNICATION

Robert H. Cichewicz *et al.*

Mutanobactin A from the human oral pathogen *Streptococcus mutans* is a cross-kingdom regulator of the yeast-mycelium transition

Mutanobactin A from the human oral pathogen *Streptococcus mutans* is a cross-kingdom regulator of the yeast-mycelium transition†

P. Matthew Joyner,^a Jinman Liu,^b Zhijun Zhang,^b Justin Merritt,^b Fengxia Qi^b and Robert H. Cichewicz^{*a,c}

Received 13th August 2010, Accepted 6th September 2010

DOI: 10.1039/c0ob00579g

The recent investigation of a gene cluster encoding for a hybrid PKS-NRPS metabolite in the oral pathogen *Streptococcus mutans* UA159 yielded evidence that this natural product might play an important role regulating a range of stress tolerance factors. We have now characterized the major compound generated from this gene cluster, mutanobactin A, and demonstrated that this secondary metabolite is also capable of influencing the yeast-mycelium transition of *Candida albicans*.

A complex suite of cross-species interactions are anticipated to occur among the wide range of microorganisms that constitute the human microbiome.^{1–3} While a variety of compounds such as peptides, lipids, and acyl-homoserine lactones have emerged as key players in the multifarious exchanges among microbes and their hosts,^{4,5} the important contributions of many other families of secondary metabolites have been largely overlooked. We reported that the deletion of a gene cluster encoding for a putative hybrid polyketide synthase-nonribosomal peptide synthetase (PKS-NRPS) derived metabolite in the human oral pathogenic bacteria *Streptococcus mutans* UA159 resulted in a loss of several resistance traits associated with oxygen and hydrogen peroxide tolerance, as well as biofilm formation.⁶ Given that a variety of secondary metabolites are excreted into the extracellular environment, we suspected that the biosynthetic products of this gene cluster (which we have dubbed the mutanobactins⁶) may have additional yet undefined functions related to the interaction of *S. mutans* with other members of the oral microbiome. This hypothesis recently gained support based on co-culture studies in which mutanobactin-competent *S. mutans* and a mutanobactin deletion mutant strain (Δ *mub*) were grown in the presence of *Candida albicans*. Whereas the mutanobactin producing strain of *S. mutans* was capable of maintaining *C. albicans* in a perpetual yeast morphological state (Fig. 1A), deletion of the mutanobactin cluster permitted *C. albicans* to shift into a mycelial growth pattern, which is believed to be the invasive form of the fungus (Fig. 1B). We now describe the distinctive structural features of the major hybrid PKS-NRPS-derived metabolite from *S. mutans*




Fig. 1 Phase contrast microscopy images showing co-cultures of *S. mutans* UA159 wild type (A) or Δ *mub* strain (B) with *C. albicans* ATCC 10231. Solid arrows indicate *S. mutans* cells, while open arrows denote *C. albicans* yeast (A) or mycelia (B). *S. mutans* UA159 strains were grown anaerobically overnight at 37 °C in a semi-defined medium consisting of 50% saliva, 10% artificial saliva solution, 1% glucose, 0.5% sucrose, and 0.2% peptone in water. Cultures were diluted (1 : 20) in fresh medium immediately before testing. *C. albicans* was prepared separately as overnight cultures under aerobic conditions at 37 °C in yeast extract-peptone medium with 2% glucose. For experiments, aliquots of the *C. albicans* culture were diluted (1 : 50) in the *S. mutans* cultures and the mixtures of cells grown anaerobically for 16 h at 37 °C.

UA159 and demonstrate that this biomolecule is capable of suppressing the morphological transition of *C. albicans* from yeast to mycelium.

Analytical-scale HPLC comparison of the crude extracts generated from wild-type *S. mutans* UA159 and Δ *mub* strains grown on brain-heart-infusion agar plates enabled us to identify two metabolites in the elution profile that were present only in the wild-type organism. Scale-up fermentation of the wild-type strain was performed in a bioreactor with 15 L of brain-heart-infusion broth under microaerobic conditions at 37 °C to enable the isolation and structure characterization of the major metabolite. After 48 h, the cells and broth were partitioned against ethyl acetate and the solvent removed *in vacuo*. The resulting extract was resuspended in methanol and the organic soluble material was defatted with hexane. The methanol layer was mixed with an equal volume of water and subjected to partitioning (3 \times) against dichloromethane. Solvent from the dichloromethane layer was removed under vacuum yielding 1.5 g of a dark brown crude extract. The extract was subjected to preparative-scale C₁₈ flash chromatography under gradient elution conditions (mobile phase: 20% to 100% methanol in water) to generate a single fraction (240 mg) containing the mutanobactins. The major mutanobactin metabolite was targeted for isolation by gradient preparative-scale C₁₈ HPLC (mobile phase: 50% to 100% methanol in water)

^aNatural Products Discovery Group, Department of Chemistry and Biochemistry, Stephenson Life Sciences Research Center, 101 Stephenson Parkway, University of Oklahoma, Norman, OK, 73019, U.S.A. E-mail: rhcichewicz@ou.edu; Tel: +1 (405) 325-6969

^bCollege of Dentistry, 975 NE 10th Street, The University of Oklahoma Health Sciences Center, Oklahoma City, OK, 73014, USA

^cGraduate Program in Ecology and Evolutionary Biology, University of Oklahoma, Norman, OK, USA, 73019, USA

† Electronic supplementary information (ESI) available: General methods, physical data, MS and NMR (¹H, ¹³C, ¹H-¹H COSY, ¹H-¹H TOCSY, ¹H-¹H NOESY, ¹H-¹³C HSQC, ¹H-¹³C HMBC, and ¹H-¹⁵N HMBC) for **1**. See DOI: 10.1039/c0ob00579g

followed by isocratic semi-preparative C_{18} HPLC (mobile phase 88% methanol in water), which yielded 8 mg of pure mutanobactin A (**1**) (Fig. 2 - refer to ESI for a summary of the physical and chemical data for **1**).[†]




Fig. 2 Structure of mutanobactin A (**1**), the major metabolite generated from the mutanobactin gene cluster in *S. mutans* UA159.

Inspection of the ^1H NMR data collected in $\text{DMSO}-d_6$ for **1** (Fig. S3[†]) confirmed our prior hypothesis⁶ concerning the partial peptidic nature of the mutanobactins (five exchangeable amide resonances were observed at δ_{H} 7.23, 7.77, 7.90, 8.05, and 8.59) (Table S1[†]). However, we were surprised by the presence of several overlapping methylene resonances centered at δ_{H} 1.23 that integrated for 12 protons since our previous inspection of the mutanobactin gene cluster did not provide evidence for this feature.⁶ Turning to the FT-ICR-MS data generated for **1**, we observed a pseudomolecular ion with a m/z of 719.41713 corresponding to a molecular formula of $\text{C}_{36}\text{H}_{59}\text{N}_6\text{O}_7\text{S}$ ($[\text{M} - \text{H}]^-$, calcd 719.41714, -0.01 mmu error) (Figure S1[†]). These data reinforced our suspicion that in addition to the predicted amino acid residues in **1**, the compound also contained a substantial number of non-NRPS-derived atoms.

Considering the disparity between the expected and observed structural data for **1**, we set about investigating the metabolite's structural features using a combination of ^1H and ^{13}C NMR, $^1\text{H}-^1\text{H}$ COSY, $^1\text{H}-^1\text{H}$ TOCSY, $^1\text{H}-^{13}\text{C}$ HSQC,

$^1\text{H}-^{13}\text{C}$ HMBC, and $^1\text{H}-^{15}\text{N}$ HMBC experiments (Fig. S3–10[†]). This approach yielded three distinct substructures (fragments A–C) that formed the backbone of **1** (Fig. 3). We quickly deduced that fragment A consisted of a tetrapeptide (Val-Pro-Ala-Leu) (Fig. 3) based on analysis of two-dimensional NMR correlation data (Fig. S5–10[†]). We found the $^1\text{H}-^1\text{H}$ TOCSY data were particularly useful for defining the spin systems originating from each of the alpha-protons to the hydrogens embedded in their respective amino acid side chains (Fig. 3). Marfey's analysis⁷ of the proposed amino acids was carried out by treating **1** with 6 M HCl at 120 °C for 18 h and derivatizing the hydrolysate with 1-fluoro-2,4-dinitrophenyl-5-L-alanineamide. Analytical HPLC analysis of the resultant mixture established the absolute configuration of four amino acids as L-valine (15*S*), L-proline (10*S*), D-alanine (7*R*), and L-leucine (1*S*). The other hydrolysis fragments generated from **1** were not examined due to their limited sample sizes.

Fragment B was established using a combination of NMR (*i.e.*, $^1\text{H}-^{13}\text{C}$ HMBC) (Fig. 3) and MS approaches. While many of the methylenes extending from both the ketone (C-27) and methyl (C-36) groups were readily assigned using $^1\text{H}-^{13}\text{C}$ HMBC correlation data, the significant overlap among a portion of the resonances required us to probe the metabolite using a different analytical method. Turning to ESI-MS/MS, we observed a prominent fragment ion at m/z 155 that was consistent with alpha-cleavage between C-27 and C-25 (Fig. S2[†]). This fragmentation accounted for a loss of $\text{C}_{10}\text{H}_{19}\text{O}$, which we interpreted as corresponding to a hydrocarbon chain consisting of eight methylenes and one methyl group.

The remaining atoms constituting fragment C consisted of $\text{C}_6\text{H}_9\text{N}_2\text{OS}$. Considering that this fragment must be covalently bonded with the two unassigned termini of fragment A and the single unassigned terminus of fragment B, we concluded that fragment C required two units of unsaturation. One unit of unsaturation was assigned as an amide carbonyl based on the appearance of a distinctive ^{13}C NMR resonance at δ_{C} 170.4. The second unit of unsaturation was attributed to a ring since no additional double bonds could be accounted for in the ^1H and




Fig. 3 Substructure fragments and selected two-dimensional NMR correlation data used to construct the planar structure of compound **1**. Fragment C candidates #1, #2, and #4 were ultimately rejected due to the inclusion of several incongruous $^4J_{\text{H-C}}$ couplings, as well as inconsistencies with $^1\text{H}-^1\text{H}$ TOCSY data.

^{13}C NMR data. The remaining non-hydrogen atoms in fragment C were rationalized to constitute a thiazepanone system, but five variations of this system could be envisioned. Fortunately, we could immediately eliminate 1,3-thiazepan-2-one and 1,2-thiazepan-3-one as possibilities since their predicted chemical shifts and/or questionable chemical stability were inconsistent with our data. In addition, 1,4-thiazepan-3-one and 1,3-thiazepan-4-one were also rejected since both ring systems necessitated the inclusion of isolated methylenes, which were incompatible with the observed ^1H - ^1H TOCSY data. Therefore, we deduced that fragment C was based on a 1,4-thiazepan-5-one ring system.

Noting that H-25 and C-20-NH were part of two separate spin systems (based on ^1H - ^1H TOCSY data), we restricted our investigation of fragment C to consider four possible substructure candidates (Fig. 3). Analysis of the ^1H - ^{13}C HMBC (optimized for $J_{\text{H-C}} = 8$ Hz) revealed that candidates #1 and #2 were unlikely possibilities since both of these substructures required at least three H-C couplings across four bonds. Similarly, candidate #4 also required at least one H-C coupling across four bonds and it too was removed from further consideration. In contrast, candidate #3 was strongly supported by the observed ^1H - ^{13}C HMBC data with all of the H-C couplings occurring within the target range of two to three bonds. Therefore, this fragment candidate was selected as the only plausible planar substructure for **1**. Unfortunately, extensive overlap among several critical ^1H NMR resonances within this fragment resulted in the generation of inconclusive ^1H - ^1H NOESY data, which prevented us from confidently assigning the relative configuration of fragment C. We are currently exploring whether this problem may be exacerbated by C-25 epimerization occurring as a consequence of tautomerization involving the C-27 ketone and C-26 amide carbonyl groups.

In light of the previously noted ^1H - ^{13}C HMBC correlations joining fragment C to fragments A and B (*vide supra*), we propose that **1** (Fig. 2) represents the structure of the major mutanobactin produced by *S. mutans* UA159. Further consideration of the mutanobactin biosynthetic gene cluster revealed that although most of the component building blocks predicted to appear in **1** are present, many unexpected features are also incorporated. One of the surprising components is the integration of an extended hydrocarbon chain. We speculate that regions of high homology to fatty-acid biosynthesis genes in the vicinity of the mutanobactin gene cluster may be involved in the generation of this structural component. In addition, we had predicted that the mutanobactins would contain an amino acid sequence consisting of Asp, Leu, Ser, Pro, Val, Cys, and Gly. Instead, we observed Leu, Ala, Pro, Val, Cys, and Gly (the latter two amino acids residues appear to have undergone further cyclization to generate the 1,4-thiazepan-5-one ring system). At this time, the fate of the Asp residue is unclear; however, we predict that it may have contributed to the generation of the C-26 amide carbonyl contained in the mutanobactin macrocycle.

Having already observed that a functional mutanobactin gene cluster was correlated with the ability of *S. mutans* to control the morphological transition of *C. albicans* from yeast to mycelium under co-culture conditions (Fig. 1), we next addressed whether purified **1** was capable of eliciting the same biological effect. Whereas control cultures of *C. albicans* readily formed tangled mats of intertwined mycelia *in vitro* (Fig. 4A), cells treated with **1** exhibited a remarkable preservation of yeast morphological




Fig. 4 Phase contrast microscopy images illustrating the morphology of *C. albicans* ATCC 10231 cultures grown with vehicle only (A) or treated with compound **1** (B). Whereas the mycelia morphology of the control *C. albicans* cells appeared as mats of tangled filaments, mycelia were not apparent in cells treated with **1**. For the experiment, overnight shake cultures of *C. albicans* were prepared under aerobic conditions in yeast extract-peptone medium with 2% glucose. The cells were diluted in fresh medium (1 : 100) and 100 μL aliquots added to the wells of a flat-bottom 96-well plate. Cells were treated with DMSO only (vehicle control) or compound **1** (2 μL of compound prepared in DMSO at 5 mg mL^{-1}).

characteristics (Fig. 4B), but did not show signs of decreased cell division. Based on our data, compound **1** joins a select group of two other *S. mutans*-derived biomolecules (*i.e.*, competence-stimulating peptide⁸ and *trans*-2-decenoic acid⁹) that are capable of re-directing the yeast-mycelium dimorphism of *C. albicans*.

The human microbiome is certain to yield scores of chemically unique and biologically intriguing secondary metabolites. For example, it was recently reported that *Staphylococcus aureus* biosynthesizes the NRPS-derived metabolites phevalin and aureusimine A as part of a virulence regulatory pathway.¹⁰ As new research tools emerge that further enhance the natural product research community's ability to deeply probe microbial secondary metabolite diversity,^{11,12} we expect that many new and complex intra- and inter-species relationships will come to light.¹³⁻¹⁵

Conclusion

The discovery of **1** is noteworthy since a very limited number of other signaling molecules are known to non-lethally influence the interactions between bacteria and fungi despite the expectation that these types of chemical exchanges are quite prevalent.^{16,17} Whereas the ability to generate compound **1** has been shown to be an important stress-resistance factor in *S. mutans*,⁶ we have now demonstrated that the mutanobactin metabolite is also capable of regulating cross-kingdom interactions with *C. albicans*. Furthermore, our data indicate that **1**, which contains a structurally rare 1,4-thiazepan-5-one ring system, represents a new and unusual addition to the emerging list of natural products biosynthesized by microbial species that are part of the human microbiome.

Acknowledgements

Financial support for this project was provided in part through a grant from the National Institutes of Health (1R01AI085161-01) and funds from the University of Oklahoma College of Arts and Sciences and the Department of Chemistry and Biochemistry.

References

- 1 A. Y. Peleg, D. A. Hogan and E. Mylonakis, *Nat. Rev. Microbiol.*, 2010, **8**, 340-349.

- 2 D. T. Hughes, D. A. Terekhova, L. Liou, C. J. Hovde, J. W. Sahl, A. V. Patankar, J. E. Gonzalez, T. S. Edrington, D. A. Rasko and V. Sperandio, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 9831–9836.
- 3 L. C. M. Antunes and R. B. R. Ferreira, *Crit. Rev. Microbiol.*, 2009, **35**, 69–80.
- 4 A. R. Pacheco and V. Sperandio, *Curr. Opin. Microbiol.*, 2009, **12**, 192–198.
- 5 W. W. L. Hsiao, C. Metz, D. P. Singh and J. Roth, *Endocrinol. Metab. Clin. North Am.*, 2008, **37**, 857–871.
- 6 C. Wu, R. Cichewicz, Y. Li, J. Liu, B. Roe, J. Ferretti, J. Merritt and F. Qi, *Appl. Environ. Microbiol.*, 2010, **76**, 5815–5826.
- 7 R. Bhushan and H. Brückner, *Amino Acids*, 2004, **27**, 231–247.
- 8 L. M. Jarosz, D. M. Deng, H. C. Van Der Mei, W. Crielaard and B. P. Krom, *Eukaryotic Cell*, 2009, **8**, 1658–1664.
- 9 R. Vilchez, A. Lemme, B. Ballhausen, V. Thiel, S. Schulz, R. Jansen, H. Sztajer and I. Wagner-Döbler, *ChemBioChem*, 2010, **11**, 1552–1562.
- 10 M. A. Wyatt, W. Wang, C. M. Roux, F. C. Beasley, D. E. Heinrichs, P. M. Dunman and N. A. Magarvey, *Science*, 2010, **329**, 294–296.
- 11 R. H. Cichewicz, J. C. Henrikson, X. Wang and K. M. Branscum, in *Manual of Industrial Microbiology and Biotechnology*, ed. R. H. Baltz, A. L. Demain and J. E. Davies, ASM Press, Washington, D. C., 3rd edn, 2010, ch. 7, pp. 78–95.
- 12 M. Zerikly and G. L. Challis, *ChemBioChem*, 2009, **10**, 625–633.
- 13 Y. L. Yang, Y. Xu, P. Straight and P. C. Dorrestein, *Nat. Chem. Biol.*, 2009, **5**, 885–887.
- 14 V. Schroeckh, K. Scherlach, H.-W. Nützmann, E. Shelest, W. Schmidt-Heck, J. Schuemann, K. Martin, C. Hertweck and A. A. Brakhage, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 14558–14563.
- 15 L. P. Partida-Martinez, S. Monajembashi, K.-O. Greulich and C. Hertweck, *Curr. Biol.*, 2007, **17**, 773–777.
- 16 J. Piel, *Nat. Prod. Rep.*, 2009, **26**, 338–362.
- 17 M. Tarkka, A. Sarniguet and P. Frey-Klett, *Curr. Genet.*, 2009, **55**, 233–243.