

Pd(II)-catalyzed decarboxylative allylation and Heck-coupling of arene carboxylates with allylic halides and esters†

Jiantao Wang,^a Zili Cui,^b Yuexia Zhang,^a Huajie Li,^a Long-Min Wu^a and Zhongquan Liu^{*a}

Received 8th September 2010, Accepted 12th November 2010

DOI: 10.1039/c0ob00696c

This work demonstrates an alternative method to prepare allylated arenes and aryl-substituted allylic esters *via* catalytic decarboxylative C–C bond formation using aromatic carboxylic acids and allylic halides and esters.


The development of more efficient protocols for C–C bond formation that use low-cost, safe and environmentally benign starting materials remains a critical challenge for organic chemists. Recently, metal-catalyzed decarboxylative couplings show that simple carboxylic acids seem to be efficient surrogates for both halides and organometallics.¹ Many efficient protocols to construct C–C bonds *via* palladium-catalyzed decarboxylative couplings have been developed by Myers,² Gooßen,³ Tunge,⁴ Miura,⁵ and others.⁶ Although most of these systems suffer from limited substrate scope, relatively high temperature and/or overloading additives, catalytic decarboxylative C–C bond formation remains very attractive since carboxylic acids are stable, easily available and relatively environmentally benign. We wish to report herein a palladium-catalyzed decarboxylative allylation of arenes by using arene carboxylates and allylic halides.

Allylic aromatic compounds show versatile biological activities in natural products.⁷ Additionally, they also represent an important class of organic intermediates in complex molecule synthesis due to the versatility of olefin transformations. In the past decades, two fundamental methods have been explored to construct the allylic arenes (Scheme 1).⁸ In path A, an aromatic nucleophile attacks an allyl electrophile to give the allylated arene. Traditional acid/Lewis acid catalyzed Friedel–Crafts aromatic

allylation could be also involved in this path. In path B, a polarity reversing attack gives the desired allylic arenes which might involve a transmetallation between an aryl metal intermediate formed by electrophilic aromatic substitution and an allylic partner. Although these methods are usually practical to prepare allylated arenes, most of them suffer from complex mixture of products (Friedel–Crafts aromatic allylation) and usage of organometallic reagents which might be relatively high-cost and low stability. Taking advantage of high stability, low-cost and commercial availability of arene carboxylates, we successfully accomplished a palladium-catalyzed decarboxylative arene allylation by using simple aryl carboxylic acids and allylic halides (Scheme 1, path C).⁴

As the initial research, we chose 2,4,5-trimethoxybenzoic acid and allylic bromide as standard substrates to optimize suitable conditions for this reaction (Table 1). It was found that the solvent, catalysts, and additives affect the reaction efficiency critically. The desired allylic arene was obtained in 62% yield under the conditions: 20 mol% Pd(OAc)₂, 3 eq. Ag₂CO₃, 5% DMSO/toluene (v/v), 110 °C, 2 h (entry 1). The investigation of solvent effect showed that toluene was a more effective solvent (entries 2–6). However, increase of the dosage of DMSO to 20% v/v of solvent gave only 49% yield of the product (entry 7). No products were detected without DMSO in this system (entry 8). Other silver salts as additives gave relatively low yields (entries 9–11). Addition of 20 mol% Cu₂O used as a co-catalyst gave a slightly higher yield of the product (entry 12). Reduction of the amount of Ag₂CO₃ gave relatively lower yields (entries 13). The yields of the desired product decreased to 45%, 55% and 55% by using 20 mol% of Pd(TFA)₂, 20 mol% of PdCl₂ and 10 mol% of Pd(OAc)₂ respectively (entries 14–16). A 46% yield of allylated arene was formed when the reaction was carried out at 80 °C (entry 17). Interestingly, the yields increased appreciably to 75% and 89% by reducing the dosage of Cu₂O to 5 mol% and 1 mol% respectively (entries 18 and 19). An 80% yield of the product was obtained by using 10 mol% Pd(OAc)₂ (entry 20). However, reducing the amount of Pd(OAc)₂ to 5 mol% resulted only 54% yield of the desired product (entry 21).

As depicted in Table 2, various arene carboxylates and allylic halides were tested as substrates for the coupling reaction under the typical conditions. The corresponding (*E*)- α -coupled allylic arenes were obtained as major products. Reaction of allylic iodide with 2,4,6-trimethoxybenzoic acid gave a very low yield of the product (entry 1). Electron-rich arene carboxylates gave high to excellent yields of the corresponding products except for




Scheme 1 Synthesis of allylic aromatic compounds.

^aState Key Laboratory of Applied Organic Chemistry and Department of Chemistry, Lanzhou University, Lanzhou 730000, P. R. China. E-mail: liuzhq@lzu.edu.cn; Fax: +86-931-8915557; Tel: +86-931-8912280

^bDepartment of Chemistry and Biology, Gannan Normal University, Ganzhou, Jiangxi 341000, P. R. China

† Electronic supplementary information (ESI) available: Experimental details, characterisation data and spectra. See DOI: 10.1039/c0ob00696c


Table 1 Optimization of the typical reaction conditions^a


Entry	Catalyst (mol%)	Additive (eq.)	Solvent	Yield (%) ^b
1	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	Toluene	62
2	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	Benzene	36
3	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	DCE	54
4	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	Dioxane	56
5	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	DMF	trace
6	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	DMSO	0
7	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	Toluene (20% DMSO)	49
8	Pd(OAc) ₂ (20)	Ag ₂ CO ₃ (3)	Toluene (0% DMSO)	15
9	Pd(OAc) ₂ (20)	Ag ₂ SO ₄ (3)	Toluene	31
10	Pd(OAc) ₂ (20)	Ag ₂ O (3)	Toluene	39
11	Pd(OAc) ₂ (20)	AgOAc (6)	Toluene	47
12	Pd(OAc) ₂ (20)/Cu ₂ O (20)	Ag ₂ CO ₃ (3)	Toluene	67
13	Pd(OAc) ₂ (20)/Cu ₂ O (20)	Ag ₂ CO ₃ (1)	Toluene	49
14	Pd(TFA) ₂ (20)/Cu ₂ O (20)	Ag ₂ CO ₃ (3)	Toluene	45
15	PdCl ₂ (20)/Cu ₂ O (20)	Ag ₂ CO ₃ (3)	Toluene	55
16	Pd(OAc) ₂ (10)/Cu ₂ O (20)	Ag ₂ CO ₃ (3)	Toluene	55
17 ^c	Pd(OAc) ₂ (20)/Cu ₂ O (20)	Ag ₂ CO ₃ (3)	Toluene	46
18	Pd(OAc) ₂ (20)/Cu ₂ O (5)	Ag ₂ CO ₃ (3)	Toluene	75
19	Pd(OAc)₂ (20)/Cu₂O (1)	Ag₂CO₃ (3)	Toluene	89
20	Pd(OAc)₂ (10)/Cu₂O (1)	Ag₂CO₃ (3)	Toluene	80
21	Pd(OAc) ₂ (5)/Cu ₂ O (1)	Ag ₂ CO ₃ (3)	Toluene	54

^a Reaction conditions: 2,4,5-trimethoxybenzoic acid (0.1 mmol), allylic bromide (0.2 mmol), 5% DMSO/solvent (v/v), 110 °C, 2 h, unless otherwise noted. ^b Yield determined by GC with nitrobenzene as internal standard. ^c The reaction was carried out at 80 °C.

2,4-dimethoxybenzoic acid (entries 2–5). The desired allylated arene was also obtained in good yield using halogen-substituted benzoic acid (entry 6). Although the yield was low, heterocyclic carboxylic acid was also an effective substrate (entry 7). Various alkyl-substituted allylic chlorides were proved to be effective coupling partners (entries 8–11). 1-Bromo-2-butene and aryl-substituted allylic halide gave low yields of the product (entries 12 and 13). A 33% yield of the desired product was obtained by using halide branched allylic halide such as 1, 2-dibromo-2-propene (entry 14). Although only electron-rich aromatic carboxylic acids bearing at least one *ortho* substituted group are effective substrates, the features of direct decarboxylative arene allylation using carboxylic acids and allylic halides, relatively few side products and short reaction time make this procedure very attractive.

In the Tsuji–Trost reaction, allylic halides and allylic esters act as effective coupling partners. However, the direct decarboxylative Heck-type coupling products were formed by using aromatic carboxylic acids and allylic carboxylic esters under similar conditions (Table 3). Various arene carboxylates and allylic carboxylic esters were investigated as substrates for the direct decarboxylative Heck-type coupling. Our experiments show that aromatic carboxylic acids with at least one *ortho* substituent are effective substrates (entries 1–6). It is noteworthy that polyfluorobenzoic acids give relatively low yields of the desired products (entries 7–9). Various allylic carboxylic esters have been proved to be effective coupling partners (entries 10–13). A competing Heck coupling would occur between the allylic and the acrylic double bonds (entries 12 and 13). However, the C–C bond formation only took place at the

Table 2 Palladium-catalyzed decarboxylative allylation of arenes^a


Entry	Allylic halide	Product	Yield (%) ^b
1			12
2			90
3			88
4			70
5 ^c			28
6			60
7			27
8			48
9 ^c			30
10			90 (<i>E/Z</i> = 1.8/1) ^d
11			82 (<i>E/Z</i> = 1.6/1) ^d
12 ^c			17
13 ^c			27
14 ^c			33

^a Reaction conditions: acid (0.3 mmol), allylic halide (0.6 mmol), 10 mol% Pd(OAc)₂, 1 mol% Cu₂O, 3 eq. Ag₂CO₃, 5% DMSO/toluene (v/v), 110 °C, 2 h. ^b Isolated yields. ^c 20 mol% Pd(OAc)₂. ^d The *E*- and *Z*-isomers are confirmed by NOE experiments.

carboxylic acids and allylic halides and esters. The mechanism for the decarboxylative allylation might involve a Heck-like addition of Ar-PdL₂(OAc) to the double bond followed by elimination of PdL₂(OAc)Cl or PdL₂(OAc)Br. Although one *ortho* substituent in the electron-rich arene carboxylic acid is necessary, the organometallic-reagent free, short reaction time, and relatively high chemoselectivity features make this system attractive. Further investigation of this procedure is underway in our laboratory.

Notes and references

- 1 For pioneering studies of decarboxylative cross-coupling reactions, see: M. Nilsson, *Acta Chem. Scand.*, 1958, **12**, 537; M. Nilsson, *Acta Chem. Scand.*, 1966, **20**, 423; for reviews, see: O. Baudoin, *Angew. Chem., Int. Ed.*, 2007, **46**, 1373; L. J. Gooßen, N. Rodríguez and K. Gooßen, *Angew. Chem., Int. Ed.*, 2008, **47**, 3100; L. J. Gooßen, K. Gooßen, N. Rodríguez, M. Blanchot, C. Linder and B. Zimmermann, *Pure Appl. Chem.*, 2008, **80**, 1725.
- 2 A. G. Myers, D. Tanaka and M. R. Mannion, *J. Am. Chem. Soc.*, 2002, **124**, 11250; D. Tanaka and A. G. Myers, *Org. Lett.*, 2004, **6**, 433; D. Tanaka, S. P. Romeril and A. G. Myers, *J. Am. Chem. Soc.*, 2005, **127**, 10323.
- 3 L. J. Gooßen, G. Deng and L. M. Levy, *Science*, 2006, **313**, 662; L. J. Gooßen, N. Rodríguez, B. Melzer, C. Linder, G. Deng and L. M. Levy, *J. Am. Chem. Soc.*, 2007, **129**, 4824; L. J. Gooßen, N. Rodríguez and C. Linder, *J. Am. Chem. Soc.*, 2008, **130**, 15248; L. J. Gooßen, C. Linder, N. Rodríguez, P. P. Lange and A. Fromm, *Chem. Commun.*, 2009, 7173; L. J. Gooßen, N. Rodríguez, P. P. Lange and C. Linder, *Angew. Chem., Int. Ed.*, 2010, **49**, 1111.
- 4 D. K. Rayabarapu and J. A. Tunge, *J. Am. Chem. Soc.*, 2005, **127**, 13510; C. Wang and J. A. Tunge, *Org. Lett.*, 2006, **8**, 3211; S. R. Waetzig and J. A. Tunge, *J. Am. Chem. Soc.*, 2007, **129**, 4138; J. D. Weaver and J. A. Tunge, *Org. Lett.*, 2008, **10**, 4657; R. Jana, R. Trivedi and J. A. Tunge, *Org. Lett.*, 2009, **11**, 3434; A. Recio III and J. A. Tunge, *Org. Lett.*, 2009, **11**, 5630; R. Trivedi and J. A. Tunge, *Org. Lett.*, 2009, **11**, 5650; A. J. Grenning and J. A. Tunge, *Org. Lett.*, 2010, **12**, 740; K. Chattopadhyay, R. Jana, V. W. Day, J. T. Douglas and J. A. Tunge, *Org. Lett.*, 2010, **12**, 3042; R. R. P. Torregrosa, Y. Ariyaratna, K. Chattopadhyay and J. A. Tunge, *J. Am. Chem. Soc.*, 2010, **132**, 9280; J. D. Weaver, B. J. Ka, D. K. Morris, W. Thompson and J. A. Tunge, *J. Am. Chem. Soc.*, 2010, **132**, 12179.
- 5 M. Nakano, H. Tsurugi, T. Satoh and M. Miura, *Org. Lett.*, 2008, **10**, 1851; M. Miyasaka, A. Fukushima, T. Satoh, K. Hirano and M. Miura, *Chem.–Eur. J.*, 2009, **15**, 3674; M. Yamashita, K. Hirano, T. Satoh and M. Miura, *Org. Lett.*, 2009, **11**, 2337; M. Shimizu, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2009, **74**, 3478; M. Yamashita, K. Hirano, T. Satoh and M. Miura, *Org. Lett.*, 2010, **12**, 592.
- 6 For selected examples, see: P. Forgione, M.-C. Brochu, M. St-Onge, K. H. Thesen, M. D. Bailey and F. Bilodeau, *J. Am. Chem. Soc.*, 2006, **128**, 11350; J.-M. Becht, C. Catala, C. Le Drian and A. Wagner, *Org. Lett.*, 2007, **9**, 1781; A. Voutchkova, A. Coplin, N. E. Leadbeater and R. H. Crabtree, *Chem. Commun.*, 2008, 6312; J. Cornella, C. Sanchez, D. Banawa and I. Larrosa, *Chem. Commun.*, 2009, 7176; P. Hu, J. Kan, W. P. Su and M. C. Hong, *Org. Lett.*, 2009, **11**, 2341; J. Cornella, P. Lu and I. Larrosa, *Org. Lett.*, 2009, **11**, 5506; Z. Y. Wang, Q. P. Ding, X. D. He and J. Wu, *Org. Biomol. Chem.*, 2009, **7**, 863; R. Shang, Y. Fu, Y. Wang, Q. Xu, H. Z. Yu and L. Liu, *Angew. Chem., Int. Ed.*, 2009, **48**, 9350; R. Shang, Y. Fu, J.-B. Li, S.-L. Zhang, Q.-X. Guo and L. Liu, *J. Am. Chem. Soc.*, 2009, **131**, 5738; Z.-M. Sun and P. Zhao, *Angew. Chem., Int. Ed.*, 2009, **48**, 6726; H.-P. Bi, L. Zhao, Y.-M. Liang and C.-J. Li, *Angew. Chem., Int. Ed.*, 2009, **48**, 792; S.-L. Zhang, Y. Fu, R. Shang, Q.-X. Guo and L. Liu, *J. Am. Chem. Soc.*, 2010, **132**, 638; R. Shang, Q. Xu, Y.-Y. Jiang, Y. Wang and L. Liu, *Org. Lett.*, 2010, **12**, 1000; K. Xie, Z. Yang, X. Zhou, X. Li, S. Wang, Z. Tan, X. An and C.-C. Guo, *Org. Lett.*, 2010, **12**, 1564; F. Zhang and M. F. Greaney, *Angew. Chem. Int. Ed.*, 2010, **49**, 2768; Y. Zhao, Y. Zhang, J. Wang, H. Li, L.-M. Wu and Z.-Q. Liu, *Synlett*, 2010, **15**, 2352.
- 7 E. J. Corey, L. O. Weigel, D. Floyd and M. G. Bock, *J. Am. Chem. Soc.*, 1978, **100**, 2916; E. Wenkert, J. B. Fernandes, E. L. Michelotti and C. S. Swindell, *Synthesis*, 1983, 701; B. M. Trost and F. D. Toste, *J. Am. Chem. Soc.*, 2000, **122**, 11262; B. M. Trost, O. R. Thiel and H.-C. Tsui, *J. Am. Chem. Soc.*, 2002, **124**, 11616.
- 8 M. Ochiai, M. Arimoto and E. Fujita, *Tetrahedron Lett.*, 1981, **22**, 4491; J. W. Dieter, Z. Li and K. M. Nicholas, *Tetrahedron Lett.*, 1987, **28**, 5415; M. Kodomari, S. Nawa and T. Miyoshi, *J. Chem. Soc., Chem. Commun.*, 1995, 1895; J. W. Faller and R. G. Kultyshev, *Organometallics*, 2002, **21**, 5911; N. Tsukada, Y. Yagura, T. Sato and Y. Inoue, *Synlett*, 2003, 1431; S. Hayashi, K. Hirano, H. Yorimitsu and K. Oshima, *J. Am. Chem. Soc.*, 2006, **128**, 2210; S. Sebelius, V. J. Olsson, O. A. Wallner and K. J. Szabó, *J. Am. Chem. Soc.*, 2006, **128**, 8150; M. Iwasaki, S. Hayashi, K. Hirano, H. Yorimitsu and K. Oshima, *J. Am. Chem. Soc.*, 2007, **129**, 4463; S. E. Denmark and N. S. Werner, *J. Am. Chem. Soc.*, 2008, **130**, 16382; V. Maslak, Z. T. Vujosevic and R. N. Saicic, *Tetrahedron Lett.*, 2009, **50**, 1858; J. Cvengroš, J. Schütte, N. Schlörner, J. Neudörfl and H.-G. Schmalz, *Angew. Chem., Int. Ed.*, 2009, **48**, 6148; S. E. Denmark and N. S. Werner, *J. Am. Chem. Soc.*, 2010, **132**, 3612; H. Ohmiya, Y. Makida, D. Li, M. Tanabe and M. Sawamura, *J. Am. Chem. Soc.*, 2010, **132**, 879.