

A non-metal catalysed oxidation of primary azides to nitriles at ambient temperature†

Manjunath Lamani, Pradeep Devadig and Kandikere Ramaiah Prabhu*

Received 19th November 2011, Accepted 10th January 2012

DOI: 10.1039/c2ob06949k

A novel non-metal catalyzed oxidation of organic azides to nitriles under solvent-free conditions is presented employing catalytic amounts of KI, and DABCO in aq. TBHP at room temperature. This non-metal catalyzed oxidation of azides provides good selectivity as double and triple bonds were not oxidized under the present reaction conditions.

Introduction

The nitrile functionality has a unique identity in organic synthesis as it can be easily transformed into valuable functional groups such as acids, amines, amides *etc.*¹ Traditionally, aliphatic nitriles are synthesized by employing inorganic cyanides through nucleophilic substitution methods, whereas aromatic nitriles are prepared by Sandmeyer's reaction.^{2–5} The search for convenient and useful methods for the synthesis of nitriles has led to novel chemistry as well as newer methods of synthesizing nitriles from a variety of easily available substrates without using inorganic cyanide. Most of the methods that transform alcohols or aldehydes to nitriles use NH₃ as the nitrogen source.^{6a,b} Additionally, dehydration of aldoximes or amides to form nitriles is also reported.^{6c,d}

Azides are generally considered hazardous. Therefore, the utility of azides, particularly organic azides, has been limited to a greater extent.⁷ Generally, azides are used in the Staudinger reaction, Schmidt reaction and Curtius rearrangement.^{1f} After the discovery of click chemistry, the utility triazole, which is generated by the reaction of azide and terminal alkynes, as a linker between biomolecules and ligands has led several research groups to re-evaluate the utility of azides in organic synthesis.⁸ The contemporary developments, and versatile applications of azides in chemistry, biology, medicine and materials science was reviewed recently by Brase *et al.*^{8b}

Efforts in this direction revealed that azides that contain α -hydrogens are useful precursors for the oxidation to the corresponding nitriles. Hence, these strategies turned out to be useful methods to synthesize benzonitriles from the corresponding benzylic azides. Recently, Jiao and co-workers⁹ reported copper


catalyzed transformation of methylarenes to aryl nitriles and conversion of allylarenes to alkenyl nitriles by employing excess Me₃SiN₃, FeCl₂, and DDQ (3 equiv.). These two protocols are specific to methylarenes and allylic systems respectively.⁹ Recently, copper catalysed oxidation of aliphatic azides to their corresponding nitriles was reported by our group.¹⁰ Apart from these protocols, ruthenium catalysed aerobic oxidation of aliphatic primary azides to nitriles¹¹ and BrF₃ promoted transformation of primary azides to the corresponding nitriles^{12a} were reported. Interestingly, Pd⁰ is known to catalyse decomposition of primary azides to the corresponding nitriles along with amines, and imines as by-products.^{12b} Most of these reports utilize either metal catalysts or harsh reaction conditions such as heating the reaction mixture to promote the reaction. Interestingly, iodine compounds^{13,14} are used in a variety of oxidative transformations. Hence, there is a surge in the use of iodine and its derivatives as they offer environmentally benign reactions.¹⁴ In this context, we report a selective oxidation of aliphatic primary azides to the corresponding nitriles by employing a catalytic amount of KI using TBHP as the oxidant.

Results and discussion

Initial optimization studies were performed by employing 4-methoxybenzyl azide (**1a**) as the precursor. As seen in Table 1, preliminary investigations were initiated using TBHP as an oxidant. The oxidation of **1a** with a catalytic amount of I₂ and TBHP (3 equiv.) furnished a mixture of corresponding nitrile (**2a**), acid (**3a**) and amide (**4a**) (entry 1, Table 1). To circumvent this problem, further reactions were performed using I₂ and TBHP along with a variety of organic bases. Utility of I₂ either in stoichiometric or catalytic amounts, under solvent-free conditions, furnished a mixture of nitrile (**2a**), acid (**3a**) and amide (**4a**) in the presence of a catalytic amount of pyridine (entries 2 and 3). Further optimization studies with a variety of bases such as triethylamine, DIPEA, DMAP, DABCO resulted in mixture of corresponding nitrile (**2a**), and acid (**3a**) (entries 4–8, Table 1).

Department of Organic Chemistry, Indian Institute of Science, Bangalore 560 012, Karnataka, India. E-mail: prabhu@orgchem.iisc.ernet.in; Fax: (+) 91-80-23600529; Tel: (+) 91-80-22932887

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c2ob06949k

Table 1 Optimization studies


Entry	Catalyst (equiv.)	Additive (equiv.)	Solvent	Yield ^a (%)		
				2a	3a ^b	4a ^b
1	I ₂ (1)	None	Neat	24	64	12
2	I ₂ (1)	Pyridine (0.2)	Neat	11	67	18
3	I ₂ (0.25)	Pyridine (0.25)	Neat	32	40	28
4	I ₂ (0.2)	Et ₃ N (0.2)	Neat	n.r.		
5	I ₂ (0.2)	DIPEA (0.2)	Toluene ^c	20	<5	—
6	I ₂ (0.2)	DMAP (0.2)	Toluene ^c	60	<5	—
7	I ₂ (0.2)	DABCO (0.2)	Toluene ^c	64	<5	—
8	I ₂ (0.2)	DABCO (1)	Neat	72	<5	—
9	KI (0.25)	None	Neat	70	<5	—
10	KI (0.25)	DABCO (1)	Neat	88	—	—
11	KI (0.25)	DABCO (0.25)	Neat	88	—	—
12	KI (0.2)	DABCO (0.25)	Neat	82	—	—
13	KI (0.1)	DABCO (0.1)	Neat	73	—	—
14	KI (0.1)	DABCO (1)	Neat	72	—	—
15 ^d	KI (0.25)	DABCO (0.25)	Neat	66	—	—
16 ^e	KI (0.25)	DABCO (0.25)	Neat	60	—	—

^a Isolated yield. ^b Yield determined by GC. ^c Reaction is carried out at 110 °C. ^d Used 2 equiv. of aq. TBHP. ^e Used 1 equiv. of aq. TBHP.

These results suggest that I₂–TBHP system, under the present reaction conditions, is not a suitable combination for the oxidation of azides. As KI is known to react with TBHP to produce I₂, we carried out a few experiments using KI as the iodine source (entries 9–16). As expected, reaction of **1a** with KI (25 mol%) and TBHP (1 equiv.) under solvent-free reaction conditions furnished the corresponding nitrile **2a** in 70% along with trace amount of **3a** (entry 9). Remarkably, the reaction of **1a** with KI (25 mol%), DABCO (1 equiv.) and TBHP (3 equiv.) under solvent-free reaction conditions produced **2a** as the only product (entry 10). Further, experiments using varied amounts of KI, TBHP and DABCO at room temperature (entries 11–16) indicated that the reaction of **1a** with KI (25 mol%), TBHP (3 equiv.) and DABCO (25 mol%) are the most suitable conditions to furnish **2a** in good yield (88%, entry 11).¹⁵ Further, employing a variety of oxidants such as molecular oxygen, H₂O₂, cumene hydroperoxide, oxone, or sodium perborate for the present oxidation did not furnish the nitrile.

Having established the optimal reaction conditions for the oxidation, we continued our investigation to generalize the results. As can be seen from Table 2, benzyl azide (**1b**) underwent a smooth reaction under the optimized conditions to afford benzonitrile (**2b**) in excellent yield (entry 1, Table 2). The azides bearing methoxy substituents were oxidized to the corresponding nitriles **2c** and **2d** in excellent yields (86% and 89% respectively, entries 2 and 3). Oxidation of 2,4-dichlorobenzyl azide (**1e**) and 4-benzyloxybenzyl azide (**1f**) furnished corresponding nitriles **2e** and **2f** in good yields (81% and 83%, entries 4 and 5). Allyloxy and propargyloxy substituted benzyl azides **1g** and **1h** underwent a facile oxidation to furnish the corresponding nitriles **2g** and **2h** in good yields (entries 6 and 7). Cinnamyl azide **1i**

was smoothly transformed to cinnamitrile **2i** under the standard reaction conditions (85%, entry 8). It is noteworthy that allyl, propargyl, and cinnamyl groups, which are easily amenable for oxidation, are well tolerated under the present reaction conditions. 1-(Azidomethyl)naphthalene **1j** provided the product 1-naphthoitrile **2j** in good yield (entry 9, 79%). Electron withdrawing substituents such as a nitro group at *p*-position of benzyl azide **1k** needed a longer time to produce the corresponding nitrile **2k** in good yield (91%). Similarly, the oxidation of methyl 2-(azidomethyl)benzoate (**1l**) proceeded effectively to furnish the corresponding nitrile **2l** in good yield. The oxidation of heliotropin derived benzyl azide **1m** under standard reaction conditions proceeded efficiently to furnish the corresponding nitrile (**2m**) (85%, entry 12). Despite the fact that 2-(azidomethyl)pyridine needed longer time for the oxidation, the reaction produced good yield of the corresponding nitrile **2n** in 80%, (entry 13). Oxidation of 1-azidoundecane (**1o**) under reflux conditions furnished the corresponding nitrile **2o** in moderate yield. Nevertheless, the sluggish reactivity of aliphatic azide is exploited in demonstrating a chemoselective oxidation of a benzylic azide in the presence of an aliphatic azide. Hence, the oxidation of diazide **1p** under the present reaction conditions resulted in the oxidation of the benzylic azide to nitrile, whereas the aliphatic azido-functionality was inert under the reaction conditions to yield the corresponding ω-azido nitrile **2p** in good yield (77%, entry 15). Similarly, secondary azides under the optimized reaction conditions were unaffected during the reaction.































A few experiments were performed to follow the reaction mechanism (Scheme 1). It is well known that the reaction of KI with TBHP produces either hypoiodite¹⁶ or I₂ and KOH.¹⁷ The reaction of I₂ and KOH with *p*-methoxybenzyl azide either at room temperature or at reflux conditions did not furnish the nitrile and the starting material was intact. However, the oxidation of azide **1a** with TBHP and KI under standard reaction conditions in the presence of a radical scavenger BHT (2,6-bis(1,1-dimethylethyl)-4-methylphenol) failed to produce **2a** and the starting material was recovered. This reaction indicates that the reaction is proceeding through a radical intermediate. Based on this preliminary information, we believe that the reaction might go through either *tert*-butoxy radical, or *tert*-butylperoxy radical, which further reacts with azide to form the corresponding nitrile. However, the role of DABCO is not very clear. We believe that DABCO may react with I₂ to generate DABCO salt, which may further react with TBHP to generate *tert*-butoxy radical, or *tert*-butylperoxy radical.¹⁸ This reaction mechanism may be similar to the one that is reported for the oxidation of benzylic amines to the corresponding nitrile.¹⁹ Further work is underway in our laboratory.

Conclusions


In conclusion we have demonstrated a novel non-metal catalyzed oxidation of organic azides to nitriles under solvent-free conditions. This reaction employs catalytic amounts of KI (25 mol %), DABCO (25 mol%) and aq. TBHP (3 equiv., 70% solution in water) as an oxidant. Besides this, the reaction provide a good selectivity, as double and triple bonds were not oxidized under

Table 2 Oxidation of primary azides to nitriles

$$\text{R-CH}_2\text{-N}_3 + \text{KI} \xrightarrow[\text{70\%}]{\text{Aq. TBHP}} + \text{NMP} \xrightarrow[\text{RT}]{\text{Neat}} \text{R-CN}$$

Entry	Substrate	Time (h)	Product	Yield (%) ^a
1 ^b		10		91
2		7		86
3		4		89
4		24		81
5		18		83
6		36		83
7		24		81
8		8		85
9		24		79
10		48		91
11		48		83
12		8		85
13		48		80
14		24		51 ^b
15		18		77

^a Isolated yield. ^b Yield determined by GC.



Scheme 1 Proposed mechanism for the oxidation of primary azides to nitriles.

the reaction conditions. To the best of our knowledge, this is the first report of metal-free oxidation of organic azides to the corresponding nitriles. Additionally, chemoselective oxidation of benzylic azides against aliphatic azides increases the potential application of the present method.

Experimental

Typical experimental procedure for the synthesis of azides^{10b}

Benzyl bromide or chloride derivatives (1 mmol) and sodium azide (2 mmol) in acetonitrile (4 mL) were refluxed until completion of the reaction (monitored by TLC). Evaporated the solvent, quenched with water, extracted with Et₂O (3 × 25 mL), combined organic extract was washed with water (3 × 25 mL), dried over Na₂SO₄, concentrated under vacuum. The residue was chromatographed on silica gel with EtOAc:Hexane, 2:98 to afford benzyl azide derivatives.

General experimental

NMR spectra were recorded on a JEOL LA-300, BRUKER-AV400 spectrometer in CDCl₃, tetramethylsilane (TMS; δ = 0.00 ppm) served as internal standards for ¹H NMR. The corresponding residual non-deuterated solvent signal (CDCl₃: δ = 77.00 ppm) was used as internal standards for ¹³C NMR. IR spectra were measured using a JASCO FT/IR-410 spectrometer, and Perkin-Elmer FT/IR Spectrum BX, GX. Mass spectra were measured with Micromass Q-ToF (ESI-HRMS). Column chromatography was conducted on Silica gel 230–400 mesh (Merck) and preparative thin-layer chromatography was carried out using SILICA GEL GF-254.

Typical experimental procedure for the synthesis of nitriles

Typical experimental procedure for 4-methoxybenzotrile (**2a**): Aqueous TBHP (70% solution in water, 3 mmol) was added dropwise over one minute to a well stirred mixture of 1-(azidomethyl)-4-methoxybenzene (**1a**, 1 mmol), KI (0.25 mmol) and DABCO (0.25 mmol) and stirred at room temperature for 8 h. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with the saturated aq. Na₂S₂O₃, and extracted with ethyl acetate (3 × 15 mL). The combined organic layer was washed with dil. HCl (10%, 5 mL), followed by water, dried over anhydrous Na₂SO₄, and the solvent was removed under vacuum to afford the crude product. The crude mixture was purified by column chromatography on silica gel (EtOAc–

pet ether 1:19 to 1:10) to furnish 88% of 4-methoxybenzotrile (**2a**) as a colorless solid.

Characterization data for azides

1-(Azidomethyl)-4-methoxybenzene^{10b} (**1a**). Colorless liquid; Yield – 90%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2098; ¹H NMR (400 MHz, CDCl₃): δ 7.24 (d, *J* = 8.28 Hz, 2H), 6.91 (d, *J* = 8.52 Hz, 2H), 4.26 (s, 2H), 3.81 (s, 3H); ¹³C NMR (100 MHz, CDCl₃): δ 159.6, 129.7, 127.3, 114.1, 55.2, 54.3, Anal. Calcd for C₈H₉N₃O C, 58.88; H, 5.56; N, 25.75; Found: 58.81; H, 5.82; N, 26.00.

(Azidomethyl)benzene^{10b} (**1b**). Colorless liquid; Yield – 85%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2097; ¹H NMR (400 MHz, CDCl₃): δ 7.40–7.26 (m, 5H), 4.32 (s, 2H); ¹³C NMR (100 MHz, CDCl₃): 135.3, 128.8, 128.3, 128.2, 54.8; Anal. Calcd for C₇H₇N₃ C, 63.14; H, 5.30; N, 31.56; Found: 62.77; H, 5.36; N, 31.16.

4-(Azidomethyl)-1,2-dimethoxybenzene^{10b} (**1c**). Pale yellow oil; Yield – 74%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2096; ¹H NMR (400 MHz, CDCl₃): δ 7.00–6.70 (m, 3H), 4.27 (s, 2H), 3.90 (s, 3H), 3.89 (s, 3H); ¹³C NMR (100 MHz, CDCl₃): δ:149.1, 149.0, 54.7, 127.7, 120.8, 111.3, 111.00, 55.8, 55.8; GC-MS (*m/z*): 193 (M⁺).

5-(Azidomethyl)-1,2,3-trimethoxybenzene²⁰ (**1d**). Colorless liquid; Yield – 75%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2101; ¹H NMR (400 MHz, CDCl₃): δ 6.53 (s, 2H), 4.28 (s, 2H), 3.87 (s, 6H), 3.85 (s, 3H); ¹³C NMR (100 MHz, CDCl₃): δ 153.4, 137.9, 130.9, 105.1, 60.8, 56.1, 55.0; Anal. Calcd for C₁₀H₁₃O₃N₃ C, 53.80; H, 5.87; N, 18.82; Found: 53.85; H, 6.25; N, 19.17.

1-(Azidomethyl)-2,4-dichlorobenzene (**1e**)²¹ Colorless liquid; Yield – 90%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2105; ¹H NMR (400 MHz, CDCl₃): 7.42 (d, *J* = 2 Hz, 1H), 7.34–7.32 (m, 1H), 7.29–7.26 (m, 1H), 4.46 (s, 2H); ¹³C NMR (100 MHz, CDCl₃): δ 134.8, 134.4, 131.9, 130.7, 129.6, 127.4, 51.7; Anal. Calcd for C₇H₅Cl₂N₃ C, 41.61; H, 2.49; N, 20.80; Found: C, 41.26; H, 2.83; N, 20.98.

1-(Azidomethyl)-4-(benzyloxy)benzene²² (**1f**). White Solid; Yield – 75%; *mp*: 35–37 °C (lit.²¹ 35 °C); *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2097; ¹H NMR (400 MHz, CDCl₃): δ 7.43–7.30 (m, 5H), 7.23 (d, *J* = 8.56 Hz, 2H), 6.97 (d, *J* = 8.60 Hz, 2H), 5.06 (s, 2H), 4.25 (s, 2H); ¹³C NMR (100 MHz, CDCl₃): δ 158.8, 136.8, 129.7, 128.6, 127.9, 127.7, 127.4, 115.1, 70.0, 54.3; Anal. Calcd for C₁₄H₁₃ON₃ C, 70.28; H, 5.48; N, 17.56; Found: C, 70.47; H, 5.68; N, 17.93.

1-(Allyloxy)-4-(azidomethyl)benzene (**1g**). Colorless liquid; Yield – 85%; *R_f* (25% EtOAc–hexane) 0.9; IR (Neat, cm⁻¹): 2098; ¹H NMR (400 MHz, CDCl₃): δ 7.23 (d, *J* = 8.60 Hz, 2H), 6.92 (d, *J* = 8.68 Hz, 2H), 6.10–6.00 (m, 1H), 5.41 (dd, *J*₁ = 1.6 Hz, *J*₂ = 17.28 Hz, 1H), 5.29 (dd, *J*₁ = 1.4 Hz, *J*₂ = 10.48 Hz, 1H), 4.54 (dt, *J*₁ = 1.52 Hz, *J*₂ = 5.28 Hz, 2H) 4.26 (s, 2H); ¹³C NMR (100 MHz, CDCl₃): δ 158.6, 133.0, 129.7, 127.5, 117.7, 114.9, 114.8, 68.8, 54.4; Anal. Calcd for C₁₀H₁₁N₃O C, 63.48; H, 5.86; N, 22.21; Found: C, 63.48; H, 5.90; N, 22.14.

1-(Azidomethyl)-4-(prop-2-yn-1-yloxy)benzene.²³ (1h). Pale yellow liquid; Yield – 81%; R_f (25% EtOAc–hexane) 0.8; IR (Neat, cm^{-1}) 2098 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 7.25 (d, $J = 8.6$ Hz, 2H), 6.98 (d, $J = 8.6$ Hz, 2H), 4.69 (d, $J = 2.28$ Hz, 2H), 4.27 (s, 2H), 2.53 (t, $J = 2.36$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 157.5, 129.6, 128.3, 115.1, 78.3, 77.3, 76.9, 76.7, 75.7, 55.7, 54.2; Anal. Calcd for $\text{C}_{10}\text{H}_9\text{N}_3\text{O}$ C, 64.16; H, 4.85; N, 22.45; Found: 63.80; H, 5.12; N, 22.21.

(3-Azidoprop-1-en-1-yl) benzene.^{10b} (1i). Pale yellow liquid; Yield – 75%; R_f (25% EtOAc–hexane) 0.9; IR (Neat, cm^{-1}) 2098; ^1H NMR (400 MHz, CDCl_3): δ 7.40–7.23 (m, 5H), 6.64 (d, $J = 15.8$ Hz, 1H), 6.23 (dt, $J_1 = 6.6$ Hz, $J_2 = 15.72$ Hz, 1H), 3.93 (d, $J = 6.56$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 135.9, 134.5, 128.6, 128.1, 126.6, 122.3, 52.9; Anal. Calcd for $\text{C}_9\text{H}_9\text{N}_3$ C, 67.90; H, 5.70; N, 26.40; Found: 67.73; H, 5.81; N, 26.04.

1-(Azidomethyl)naphthalene.^{10a} (1j). Colorless liquid Yield – 80%; R_f (25% EtOAc–hexane) 0.8; IR (Neat, cm^{-1}): 2099; ^1H NMR (400 MHz, CDCl_3): δ 8.0 (d, $J = 8$ Hz, 1H), 7.87–7.81 (m, 2H), 7.57–7.48 (m, 2H), 7.44–7.41 (m, 2H), 4.71 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 133.8, 131.3, 130.9, 129.3, 128.7, 127.2, 126.6, 126.1, 125.1, 123.4, 52.9; Anal. Calcd for $\text{C}_{11}\text{H}_9\text{N}_3$ C, 72.11; H, 4.95; N, 22.94; Found: C, 72.08; H, 4.99; N, 22.66.

1-(Azidomethyl)-4-nitrobenzene.²⁴ (1k). Pale yellow liquid; Yield – 92%; R_f (25% EtOAc–hexane) 0.8; IR (Neat, cm^{-1}): 2099; ^1H NMR (400 MHz, CDCl_3): 8.24 (d, $J = 8.52$ Hz, 2H), 7.50 (d, $J = 8.48$ Hz, 2H), 4.51 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 147.7, 142.7, 128.5, 123.9, 53.7; Anal. Calcd for $\text{C}_7\text{H}_6\text{N}_4\text{O}_2$, C, 47.19; H, 3.39; N, 31.45; Found: C, 47.35; H, 3.72; N, 30.86.

Methyl 2-(azidomethyl)benzoate.^{10b} (1l). Pale yellow liquid; Yield – 76%; R_f (25% EtOAc–hexane) 0.8; IR (Neat, cm^{-1}): 2103, 1719; ^1H NMR (400 MHz, CDCl_3): δ 8.02 (dd, $J_1 = 1.24$ Hz, $J_2 = 7.84$ Hz, 1H), 7.55 (dt, $J_1 = 1.44$ Hz, $J_2 = 7.64$ Hz, 1H), 7.49–7.47 (m, 1H), 7.40 (dt, $J_1 = 1.44$ Hz, $J_2 = 7.68$ Hz, 1H), 4.81 (s, 2H), 3.92 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 167.0, 137.3, 132.6, 131.1, 129.7, 128.7, 128.0, 53.1, 52.3; HRESI-MS (m/z): Calculated for $\text{C}_9\text{H}_9\text{N}_3\text{O}_2$ (M + Na): 214.0592, found (M + Na): 214.0554.

5-(Azidomethyl)benzo[d][1,3]dioxole.²⁰ (1m). Colorless liquid; Yield – 83%; R_f (25% EtOAc–hexane) 0.8; IR (Neat, cm^{-1}): 2097; ^1H NMR (400 MHz, CDCl_3): δ 6.80 – 6.76 (m, 3H), 5.97 (s, 2H), 4.22 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 148.0, 147.6, 129.0, 121.9, 108.7, 108.3, 101.2, 54.7; Anal. Calcd for $\text{C}_8\text{H}_7\text{O}_2\text{N}_3$ C, 54.24; H, 3.98; N, 23.72; Found: C, 54.79; H, 4.37; N, 23.83.

2-(Azidomethyl)pyridine.²⁵ (1n). Colorless oil; Yield – 85%; R_f (25% EtOAc–hexane) 0.5; IR (Neat, cm^{-1}): 2103; ^1H NMR (400 MHz, CDCl_3): δ 8.60–8.59 (m, 1H), 7.71 (td, $J_1 = 1.8$ Hz, $J_2 = 7.68$ Hz, 1H), 7.34 (d, $J = 7.76$ Hz, 1H), 7.25–7.22 (m, 1H), 4.49 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 155.6, 149.6, 136.9, 122.8, 121.9, 55.6; Anal. Calcd for $\text{C}_6\text{H}_6\text{N}_4$ C, 53.72; H, 4.51; N, 41.77; Found: C, 53.75; H, 4.95; N, 41.59.

2-(6-Azidomethyl)oxy-5-(azidomethyl)-1,3-dimethoxybenzene (1o)

Colorless liquid; Yield – 75%; R_f (25% EtOAc–hexane) 0.7; Prepared as shown in general Scheme 1. IR (Neat, cm^{-1}): 2097; ^1H NMR (400 MHz, CDCl_3): δ 6.52 (s, 2H), 4.28 (s, 2H), 3.96 (t, $J = 6.6$ Hz, 2H), 3.85 (s, 6H), 3.28 (t, $J = 6.92$ Hz, 2H), 1.79–1.72 (m, 2H), 1.65–1.60 (m, 2H), 1.53–1.43 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3): δ 153.4, 137.1, 130.7, 105.2, 73.1, 56.1, 55.1, 51.4, 29.9, 28.8, 26.5, 25.4; HRESI-MS (m/z): Calculated for $\text{C}_{15}\text{H}_{22}\text{N}_6\text{O}_3$ (M + Na): 357.1651, found (M + Na): 357.1653.

Characterisation data for nitriles

4-Methoxybenzonitrile.^{9a} (2a). Colorless solid; Yield – 88%; mp : 55–57 °C (lit.³ 56–57 °C); R_f (25% EtOAc–hexane) 0.50; Prepared as given in general experimental procedure. IR (Neat, cm^{-1}): 2219; ^1H NMR (400 MHz, CDCl_3): δ 7.59 (d, $J = 8.9$ Hz, 2H), 6.95 (d, $J = 8.9$ Hz, 2H), 3.86 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 162.8, 133.9, 119.2, 114.7, 103.9, 55.5; HRESI-MS (m/z): Calculated for $\text{C}_8\text{H}_7\text{NO}$ (M + Na): 156.0425, found (M + Na): 156.0427.

3,4-Dimethoxybenzonitrile.^{9a} (2c). Colorless solid; Yield – 86%; mp : 63–65 °C (lit.³ 65–66 °C); R_f (25% EtOAc–hexane) 0.50; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}) 2224.0; ^1H NMR (400 MHz, CDCl_3): δ 7.29 (d, $J = 8.12$ Hz, 1H), 7.08 (s, 1H), 6.91 (d, $J = 8.32$ Hz, 1H), 3.94 (s, 3H), 3.91 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 152.8, 149.1, 126.4, 119.2, 113.8, 111.1, 103.8, 56.1, 56.0; HRESI-MS (m/z): Calculated for $\text{C}_9\text{H}_9\text{NO}_2$ (M + Na): 186.0531, found (M + Na): 186.0532.

3,4,5-Trimethoxybenzonitrile.^{6a} (2d). Colorless solid; Yield – 89%; mp : 93–95 °C (lit.³ 92–94 °C); R_f (25% EtOAc–hexane) 0.50; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}) 2224.0; ^1H NMR (400 MHz, CDCl_3): δ 6.86 (s, 2H), 3.90 (s, 3H), 3.88 (s, 6H); ^{13}C NMR (100 MHz, CDCl_3): δ 153.5, 142.3, 118.9, 109.4, 106.7, 61.0, 56.3; HRESI-MS (m/z): Calculated for $\text{C}_{10}\text{H}_{11}\text{NO}_3$ (M + Na): 216.0637, found (M + Na): 216.0644.

2,4-Dichlorobenzonitrile.²⁶ (2e). Colorless solid; Yield – 81%; mp : 56–59 °C (lit.³ 58–61 °C); R_f (25% EtOAc–hexane) 0.50; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}) 2224.0; ^1H NMR (400 MHz, CDCl_3): δ 7.62 (d, $J = 8.32$ Hz, 1H), 7.55 (d, $J = 1.56$ Hz, 1H), 7.38 (dd, $J_1 = 1.6$ Hz, $J_2 = 8.36$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ 140.1, 137.8, 134.6, 130.2, 127.8, 115.2, 111.8; HRESI-MS (m/z): Calculated for $\text{C}_7\text{H}_3\text{Cl}_2\text{N}$ (M + H): 171.9721, found (M + H): 171.9730.

4-(Benzyloxy)benzonitrile.²⁷ (2f). Colorless solid; Yield – 83%; mp : 93–95 °C; R_f (25% EtOAc–hexane) 0.5; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2217; ^1H NMR (400 MHz, CDCl_3): δ 7.58 (d, $J = 8.80$ Hz, 2H), 7.41–7.35 (m, 5H), 7.02 (d, $J = 8.76$ Hz, 2H), 5.11 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 161.9, 135.6, 134.0, 128.7, 128.4, 127.4, 119.1, 115.5, 104.1, 70.2; HRESI-MS (m/z): Calculated for $\text{C}_{14}\text{H}_{11}\text{NO}$ (M + Na): 232.0738, found (M + Na): 232.0757.

4-(Allyloxy)benzotrile.²⁸ (2g). Colorless solid; Yield – 83%; *mp*: 43–46 °C (lit.³ 43–44 °C); *R_f* (25% EtOAc–hexane) 0.5; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2223.73; ¹H NMR (400 MHz, CDCl_3): δ 7.57 (d, $J = 8.76$ Hz, 2H), 6.96 (d, $J = 8.72$ Hz, 2H), 6.08–5.98 (m, 1H), 5.42 (d, $J = 17.24$ Hz, 1H), 5.33 (d, $J = 10.52$ Hz, 1H), 4.59 (dd, $J_1 = 0.92$ Hz, $J_2 = 5.36$ Hz, 2H); ¹³C NMR (100 MHz, CDCl_3): δ 161.8, 133.9, 132.0, 119.1, 118.4, 115.3, 104.0, 69.0; HRESI-MS (*m/z*): Calculated for $\text{C}_{10}\text{H}_9\text{NO}$ (M + Na): 182.0582, found (M + Na): 182.0597.

Cinnamotrile.^{6a} (2h). Colorless liquid; Yield – 85%; *R_f* (25% EtOAc–hexane) 0.50; Prepared as shown in general experimental procedure. IR (Neat): 2218 cm^{-1} , ¹H NMR (400 MHz, CDCl_3): δ 7.46–7.37 (m, 6H), 5.87 (d, $J = 16.68$ Hz, 1H); ¹³C NMR (100 MHz, CDCl_3): δ 150.5, 133.5, 131.2, 129.1, 127.3, 118.1, 96.3; HRESI-MS (*m/z*): Calculated for $\text{C}_9\text{H}_7\text{N}$ (M + H): 130.0657, found (M + H): 130.0644.

4-(Prop-2-yn-1-yloxy)benzotrile (2i). Colorless solid; Yield – 81%; *mp*: 111–113 °C; *R_f* (25% EtOAc–hexane) 0.5; Prepared as shown in general experimental procedure. IR (Neat, cm^{-1}): 2223; ¹H NMR (400 MHz, CDCl_3): δ 7.61 (d, $J = 8.8$ Hz, 2H), 7.04 (d, $J = 8.76$ Hz, 2H), 4.76 (d, $J = 2.12$ Hz, 2H), 2.57 (t, $J = 2$ Hz, 1H); ¹³C NMR (100 MHz, CDCl_3): δ 160.6, 133.9, 118.9, 115.6, 104.9, 76.7, 76.5, 55.9; HRESI-MS (*m/z*): Calculated for $\text{C}_{10}\text{H}_7\text{NO}$ (M + H): 158.0606, found (M + H): 158.0602.

1-Naphthotrile.^{6a} (2j). Colorless solid; Yield – 79%; *mp*: 55–57 °C (lit.³ 56–58 °C); *R_f* (25% EtOAc–hexane) 0.5; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2222; ¹H NMR (400 MHz, CDCl_3): δ 8.24 (d, $J = 8.28$ Hz, 1H), 8.08 (d, $J = 8.28$ Hz, 1H), 7.93–7.90 (m, 2H), 7.70 (t, $J = 7.68$ Hz, 1H), 7.62 (t, $J = 7.28$ Hz, 1H), 7.53 (t, $J = 7.72$ Hz, 1H); ¹³C NMR (100 MHz, CDCl_3): δ 133.3, 132.9, 132.6, 132.3, 128.6, 128.5, 127.5, 125.1, 124.9, 117.8, 110.1; HRESI-MS (*m/z*): Calculated for $\text{C}_{11}\text{H}_7\text{N}$ (M + H): 154.0657, found (M^+): 154.0648.

4-Nitrobenzotrile.^{6a} (2k). Colorless solid; Yield – 91%; *mp*: 147–149 °C (lit.³ 148–149 °C); *R_f* (25% EtOAc–hexane) 0.5; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2222; ¹H NMR (400 MHz, CDCl_3): δ 8.37 (d, $J = 8.96$ Hz, 2H), 7.90 (d, $J = 8.92$ Hz, 2H); ¹³C NMR (100 MHz, CDCl_3): δ 150.0, 133.4, 124.2, 118.3, 116.8; Anal. Calcd for $\text{C}_7\text{H}_4\text{N}_2\text{O}_2$ C, 56.76; H, 2.72; N, 18.91; Found: C, 56.67; H, 3.26; N, 19.14.

Methyl 2-cyanobenzoate.²⁹ (2l). Colorless solid; Yield – 83%; *mp*: 50–53 °C (lit.³ 50–52 °C); *R_f* (25% EtOAc–hexane) 0.4; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2229; ¹H NMR (400 MHz, CDCl_3): δ 8.16–8.14 (m, 1H), 7.83–7.81 (m, 1H), 7.71–7.65 (m, 2H), 4.01 (s, 3H); ¹³C NMR (100 MHz, CDCl_3): δ 164.5, 134.8, 132.7, 132.5, 132.4, 131.2, 117.5, 112.9, 52.8; HRESI-MS (*m/z*): Calculated for $\text{C}_9\text{H}_7\text{NO}_2$ (M + Na): 184.0374, found (M + Na): 184.0382.

Benzo[d][1,3]dioxole-5-carbonitrile.³⁰ (2m). Colorless solid; Yield – 82%; *mp*: 89–92 °C (lit.³ 90–93 °C); *R_f* (25% EtOAc–hexane) 0.5; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2226; ¹H NMR (400 MHz, CDCl_3): δ

7.21 (d, $J = 8.04$ Hz, 1H), 7.03 (s, 1H), 6.86 (d, $J = 8.08$ Hz, 1H), 6.07 (s, 2H); ¹³C NMR (100 MHz, CDCl_3): δ 151.5, 147.9, 128.2, 118.9, 111.3, 109.1, 104.9, 102.2; HRESI-MS (*m/z*): Calculated for $\text{C}_8\text{H}_5\text{NO}_2$ (M + H): 148.0399, found (M + H): 148.0394.

Pyridine-2-carbonitrile.³¹ (2n). Colorless solid; Yield – 80%; *mp*: 46–49 °C (lit.³ 45–48 °C); *R_f* (25% EtOAc–hexane) 0.4; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2221.5; ¹H NMR (400 MHz, CDCl_3): δ 8.74 (d, $J = 4.48$ Hz, 1H), 7.87 (t, $J = 7.76$ Hz, 1H), 7.72 (d, $J = 7.72$ Hz, 1H), 7.55 (t, $J = 5.48$ Hz, 1H); ¹³C NMR (100 MHz, CDCl_3): δ 151.1, 137.0, 134.0, 128.5, 126.9, 117.1069; HRESI-MS (*m/z*): Calculated for $\text{C}_6\text{H}_4\text{N}_2$ (M + H): 105.0453, found (M + H): 105.0461.

4-((6-Azidohexyl)oxy)-3,5-dimethoxybenzotrile (2p). Colorless Liquid; Yield – 77%; *R_f* (25% EtOAc–hexane) 0.6; Prepared as given in the general experimental procedure. IR (Neat, cm^{-1}): 2226.9, 2094.4; ¹H NMR (400 MHz, CDCl_3): δ 6.85 (s, 2H), 4.02 (t, $J = 6.56$ Hz, 2H), 3.86 (s, 6H), 3.28 (t, $J = 6.88$ Hz, 2H), 1.79–1.72 (m, 2H), 1.67–1.60 (m, 2H), 1.53–1.41 (m, 4H); ¹³C NMR (100 MHz, CDCl_3): δ 153.8, 141.7, 119.0, 109.5, 106.5, 73.4, 56.3, 51.4, 29.9, 28.8, 26.4, 25.3; HRESI-MS (*m/z*): Calculated for $\text{C}_{15}\text{H}_{20}\text{N}_4\text{O}_3$ (M + Na): 327.1433, found (M + H): 327.1432.

Acknowledgements

This work is funded by Indian Institute of Science, CSIR, New-Delhi (No. 01(2415)/10/EMR-II) and RL Fine Chem. Thanks are due to Dr A. R. Ramesha and Prof. S. Chandrasekhar for encouragement and useful discussions. M. L. thanks CSIR, New-Delhi for a senior research fellowship.

References

- (a) A. J. Fatiadi, in *Preparation and Synthetic Applications of Cyano Compounds*, ed. S. Patai and Z. Rappaport, Wiley, New York, 1983; (b) S. Arseniyadis, K. S. Kyler and D. S. Watt, in *Organic Reactions*, ed. W. G. Dauben, Wiley, New York, vol. 31, 1984, pp. 1–374; (c) R. C. Larock, *Comprehensive Organic Transformations*, VCH, New York, 1989; (d) A. Kleemann, J. Engel, B. Kutscher and D. Reichert, *Pharmaceutical Substance: Synthesis Patents, Applications*, Georg Thieme, Stuttgart, 4th edn, 2001; (e) J. S. Miller and J. L. Manson, *Acc. Chem. Res.*, 2001, **34**, 563; (f) M. B. Smith and J. March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, Wiley, Hoboken, NJ, 6th edn, 2007.
- (a) F. Hagedorn, H. P. Gelbke, in *Ullmanns Enzyklopadie der Technischen Chemie*, ed. E. Bartholome, E. Biekert, H. Hellmann, H. Ley, W. M. Weigert and E. Weise, Verlag Chemie, Weinheim, 4th edn, 1979, vol. 17, p. 333; (b) G. P. Ellis and T. M. Romney-Alexander, *Chem. Rev.*, 1987, **87**, 779; (c) V. V. Grushin and H. Alper, *Chem. Rev.*, 1994, **94**, 1047; (d) L. R. Subramanian, *Science of Synthesis*, Georg Thieme, Stuttgart, 2004, vol. 19, p. 163.
- (a) K. W. Rosenmund and E. Struck, *Ber. Dtsch. Chem. Ges.*, 1919, **2**, 1749; (b) D. F. Mowry, *Chem. Rev.*, 1948, **42**, 189; (c) K. Friedrich, and K. Wallenfels, in *The Chemistry of the Cyano Group*, ed. Z. Rappaport, Interscience, London, 1970, p. 67; (d) J. Lindley, *Tetrahedron*, 1984, **40**, 1433; (e) P. Kurtz in *Houben-Weyl, Methoden der Organischen Chemie*, ed. E. Muller, Georg Thieme, Stuttgart, 4th edn, 1952, Part III, vol. 8, p. 302.
- (a) M. Sundermeier, A. Zapf and M. Beller, *Chem. Commun.*, 2004, 1388; (b) D. Wang, L. Kuang, Z. Li and K. Ding, *Synlett*, 2008, 69; (c) H.-J. Cristau, A. Ouali, J.-F. Spindler and M. Taillefer, *Chem.–Eur. J.*

- 2005, **11**, 2483; (d) J. Zanon, A. Klapars and S. L. Buchwald, *J. Am. Chem. Soc.*, 2003, **125**, 2890.
- 5 (a) T. Sandmeyer, *Ber. Dtsch. Chem. Ges.*, 1885, **18**, 1946; (b) T. Sandmeyer, *Ber. Dtsch. Chem. Ges.*, 1885, **18**, 1492; (c) T. Sandmeyer, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 2650.
- 6 (a) S. Iida and H. Togo, *Tetrahedron*, 2007, **63**, 8274; (b) T. Oischi, K. Yamaguchi and N. Mizuno, *Angew. Chem., Int. Ed.*, 2009, **48**, 6286; (c) E. Choi, C. Lee, Y. Na and S. Chang, *Org. Lett.*, 2002, **4**, 2369; (d) K. Yamaguchi, H. Fujiwara, Y. Ogasawara, M. Kotani and N. Mizuno, *Angew. Chem., Int. Ed.*, 2007, **46**, 3922.
- 7 V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, **41**, 2596.
- 8 (a) M. D. Best, M. M. Rowland and H. E. Bostic, *Acc. Chem. Res.*, 2011, **44**, 686; (b) S. Brase, C. Gil, K. Knepper and V. Zimmermann, *Angew. Chem., Int. Ed.*, 2005, **44**, 5188; (c) P. F. van Swieten, M. A. Leeuwenburgh, B. M. Kessler and H. S. Overkleeft, *Org. Biomol. Chem.*, 2005, **3**, 20; (d) L. Ballell, K. J. Alink, M. Slijper, C. Versluis, R. M. J. Liskamp and R. J. Pieters, *Chem. Bio Chem.*, 2005, **6**, 291; (e) *Organic Azides—Syntheses and Applications*, ed. S. Braese and K. Banert, Wiley, Chichester, 2010.
- 9 (a) W. Zhou, L. Zhang and N. Jiao, *Angew. Chem., Int. Ed.*, 2009, **48**, 7094; (b) C. Qin and N. Jiao, *J. Am. Chem. Soc.*, 2010, **132**, 15893; (c) W. Zhou, J. Xu, L. Zhang and N. Jiao, *Org. Lett.*, 2010, **12**, 2888.
- 10 (a) M. Lamani and K. R. Prabhu, *Angew. Chem., Int. Ed.*, 2010, **49**, 6622; (b) M. Maddani and K. R. Prabhu, *Tetrahedron Lett.*, 2008, **49**, 4526.
- 11 J. He, K. Yamaguchi and N. Mizuno, *J. Org. Chem.*, 2011, **76**, 4606.
- 12 (a) R. Sasson and S. Rozen, *Org. Lett.*, 2005, **7**, 2177; (b) H. Hayashi, A. Ohno and S. Oka, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 506.
- 13 (a) M. J. Mphahlele, *Molecules*, 2009, **14**, 5308; (b) M. Jereb, D. Vrazic and M. Zupan, *Tetrahedron*, 2011, **67**, 1355.
- 14 (a) M. Uyanik, H. Okamoto, T. Yasui and K. Ishihara, *Science*, 2010, **328**, 1376; (b) M. Uyanik, D. Suzuki, T. Yasui and K. Ishihara, *Angew. Chem., Int. Ed.*, 2011, **50**, 5331; (c) L. Chen, E. Shi, Z. Liu, S. Chen, W. Wei, H. Li, K. Xu and X. Wan, *Chem.–Eur. J.*, 2011, **17**, 4085.
- 15 Although 2 equiv. of TBHP was enough for the oxidation of reactive substrates such as *p*-methoxybenzyl azide, moderately reactive substrates needed 3 equiv. of TBHP. Hence, 3 equiv. of TBHP was used for rest of the substrates.
- 16 S. Yamada, D. Morizono and K. Yamamoto, *Tetrahedron Lett.*, 1992, **33**, 4329.
- 17 R. A. Kumar, C. U. Maheswari, S. Ghantasala, C. Jyoti and K. R. Reddy, *Adv. Synth. Catal.*, 2011, **353**, 401.
- 18 J. Zhang, Z. Wang, Y. Wang, C. Wan, X. Zheng and Z. Wang, *Green Chem.*, 2009, **11**, 1973.
- 19 Unlike the transformation catalysed by CuI/TBHP system, the present oxidations of benzylic azides appear to proceed via a radical intermediate. However, both the oxidations proceed smoothly with TBHP, whereas other oxidants such as molecular oxygen, H₂O₂, cumene hydroperoxide, oxone, or sodium perborate were not useful for the present transformation.
- 20 D. Imperio, T. Pirali, U. Galli, F. Pagliari, L. Cafici, P. L. Canonico, G. Sorba, A. A. Genazzani and G. C. Tron, *Bioorg. Med. Chem.*, 2007, **15**, 6748.
- 21 X. L. Wang, K. Wan and C. H. Zhou, *Eur. J. Med. Chem.*, 2010, **45**, 4631.
- 22 V. Maslak, Z. Yan, S. Xia, J. Gallucci, C. M. Hadad and J. D. Badjic, *J. Am. Chem. Soc.*, 2006, **128**, 5887.
- 23 S. Binauld, D. Damiron, T. Hamaide, J.-P. Pascault, E. Fleury and E. Drockenmuller, *Chem. Commun.*, 2008, 4138.
- 24 C. Pardin, I. Roy, W. D. Lubell and J. W. Keillor, *Chem. Biol. Drug Des.*, 2008, **72**, 189.
- 25 L. P. Spencer, R. Altwer, P. Wei, L. Gelmini, J. Gauld and D. W. Stephan, *Organometallics*, 2003, **22**, 3841.
- 26 C. O. Kangani, B. W. Day and D. E. Kelley, *Tetrahedron Lett.*, 2008, **49**, 914.
- 27 A. K. Chakraborti and S. V. Chankeshwara, *J. Org. Chem.*, 2009, **74**, 1367.
- 28 D. S. Bhalerao, U. S. Mahajan, K. H. Chaudhari and K. G. Akamanchi, *J. Org. Chem.*, 2007, **72**, 662.
- 29 M. Sundermeier, A. Zapf, S. Mutyala, W. Baumann, J. Sans, S. Weiss and M. Beller, *Chem.–Eur. J.*, 2003, **9**, 1828.
- 30 P. Anbarasan, H. Neumann and M. Beller, *Chem.–Eur. J.*, 2011, **17**, 4217.
- 31 C. Yang and J. M. Williams, *Org. Lett.*, 2004, **6**, 2837.