

Cite this: *Org. Biomol. Chem.*, 2012, **10**, 1102

www.rsc.org/obc

PAPER

Revisiting the reaction of hydroxyl radicals with vicinal diols in water†

Dong Jiang,^a Sebastián Barata-Vallejo,^{‡b} Bernard T. Golding,^{*a} Carla Ferreri^b and Chrysostomos Chatgililoglu^{*b}

Received 7th September 2011, Accepted 26th October 2011

DOI: 10.1039/c1ob06538f

The carbonyl products of the reactions of hydroxyl radicals with three vicinal diols (ethane-1,2-diol, propane-1,2-diol and butane-2,3-diol) have been identified and quantified. Hydroxyl radicals were produced by γ -radiolysis of N_2O -saturated aqueous solutions. The reactions result in the formation of alkoxy radicals (~15%) followed by β -fragmentation, and α -hydroxyl alkyl radicals that undergo H_2O elimination. The latter process is part of a radical chain reaction at higher diol concentrations.

Introduction


The acid-catalyzed conversion of vicinal diols to aldehydes or ketones exemplified by the 'pinacol rearrangement' has been known since 1860¹ and exhibits well-established mechanisms initiated by protonation of a hydroxyl group. For diols containing tertiary alcohols (e.g. pinacol, 2,3-dihydroxy-2,3-dimethylbutane), the classical mechanism entails formation of a tertiary carbocation by loss of water from the protonated diol followed by migration of a methyl group from the adjacent carbon centre.² The discovery³ of enzyme-catalysed conversions of 1,2-diols to aldehydes in the 1960s was an apparent biochemical addition to the chemical armoury. However, it was soon established that the enzymatic mechanism^{4,5} was fundamentally different from that of the pinacol rearrangement. In particular, it was decisively shown that the key intermediates in the mechanism of the diol dehydratase-catalysed conversion of propane-1,2-diol to propionaldehyde are a substrate-derived radical (S^{*}) and a product-related radical (P^{*}) [see Scheme 1].⁶ These species arise by attack of the 5'-

deoxyadenosyl radical (Ado-CH₂^{*}), derived from coenzyme B₁₂ (5'-deoxyadenosylcobalamin, AdoCbl) on the diol leading to S^{*} and 5'-deoxyadenosine (Ado-CH₂-H).⁴ The conversion of S^{*} to P^{*} was shown by ¹⁸O-labelling to exhibit the migration of a hydroxyl group from C-2 to C-1.⁷ Independent of these discoveries, it was revealed that reactions of simple 1,2-diols with hydroxyl radicals led to the same radical (i.e. S^{*}) proposed as an intermediate in the enzyme-catalyzed reactions and the ensuing chemistry of this radical was similar.^{8,9} It was further shown that the S^{*} species underwent either acid- or base-catalyzed elimination of a diol hydroxyl group (as water or hydroxide) to afford a carbon radical stabilized by an adjacent carbonyl group¹⁰ (e.g. the formylmethyl radical from ethane-1,2-diol, see Scheme 2). The relevance of this radical chemistry to the catalytic mechanism of diol dehydratase was immediately pointed out.¹¹ In this paper, we describe product studies for the reactions of hydroxyl radicals with three diols (ethane-1,2-diol, propane-1,2-diol and butane-2,3-diol). This research serves to show the extraordinary selectivity of diol dehydratase in its action on 1,2-diols and contributes to the mechanistic understanding of both the chemistry and biochemistry of carbon radicals from vicinal diols.


Results and discussion

Radiolytic production of transients

Radiolysis of neutral water leads to the reactive species e_{aq}⁻, HO^{*} and H^{*}, together with H⁺ and H₂O₂ as shown in eqn (1). The values in parentheses represent the radiation chemical yields (*G*) in units





Scheme 1 Diol dehydratase-catalysed pathway for the conversion of propane-1,2-diol into propionaldehyde (mauve indicates isotopic labelling with deuterium or tritium; red indicates labelling with ¹⁸O).



Scheme 2 Conversion of ethane-1,2-diol to the 1,2-dihydroxyethyl radical induced by the hydroxyl radical.

of $\mu\text{mol J}^{-1}$.^{12,13} In a N_2O -saturated solution ($\sim 0.02 \text{ M}$ of N_2O), e_{aq}^- is transformed into a HO^\bullet radical with a rate constant $k_2 = 9.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (eqn (2)), affording $G(\text{HO}^\bullet) = 0.55 \mu\text{mol J}^{-1}$, *i.e.*, HO^\bullet radicals and H^\bullet atoms account for 90% and 10%, respectively, of the reactive species.



When a vicinal diol is added to the solution, this acts as a scavenger of HO^\bullet radicals and H^\bullet atoms, to generate mainly α -hydroxyalkyl radicals. In Table 1, the rate constants for the reactions of the vicinal diols used in this work with hydroxyl radicals and hydrogen atoms are shown.^{12,13} It is worth mentioning that hydroxyl radicals react with all three substrates with similar rate constants, whereas the rate constants for H^\bullet are expected to increase with increasing methyl substitution.

Ethane-1,2-diol

Sample solutions (0.2 or 2 M) of ethane-1,2-diol (**1**) in unbuffered H_2O were saturated by N_2O prior to irradiation. Irradiation doses were up to 300 Gy and to 3000 Gy for 0.2 and 2.0 M solutions, respectively, using a dose rate of *ca.* 5.9 Gy min^{-1} . 2,4-Dinitrophenylhydrazine (2,4-DNP) was added to convert the carbonyl-containing products to 2,4-DNP derivatives. HPLC analysis of the mixture of 2,4-DNPs indicated the formation of formaldehyde and acetaldehyde (Fig. 1). For recognition and quantification of the 2,4-DNPs of the carbonyl compounds, a published protocol was employed.¹⁴ The comparison was obtained from the same derivatisation procedure performed with the commercially available compounds followed by HPLC analysis.

From the concentration values of formaldehyde and acetaldehyde derived from Fig. 1, the radiation chemical yields (G) can be calculated by dividing the formation of the products (mol kg^{-1}) by the absorbed dose ($1 \text{ Gy} = 1 \text{ J kg}^{-1}$). Analysis of the data in terms of radiation chemical yield gave $G(\text{HCHO}) = 0.20$ and $G(\text{CH}_3\text{CHO}) =$

Table 1 Rate constants for the reactions of HO^\bullet radicals and H^\bullet atoms with vicinal diols

Diol	$k(\text{HO}^\bullet)^a / \text{M}^{-1} \text{ s}^{-1}$	$k(\text{H}^\bullet)^a / \text{M}^{-1} \text{ s}^{-1}$
Ethane-1,2-diol	1.8×10^9	1.4×10^7
Propane-1,2-diol	1.7×10^9	
Butane-2,3-diol	1.3×10^9	

^a Ref. 12 and 13.




Fig. 1 HPLC analyses of γ -irradiation of N_2O -saturated unbuffered solutions of 0.2 M (upper) and 2.0 M (lower) ethane-1,2-diol at various doses (dose rate = 5.9 Gy min^{-1}) after 2,4-DNP derivatisation of the carbonyl-containing compounds. The black and blue peaks correspond to formaldehyde and acetaldehyde, respectively.

$2.31 \mu\text{mol J}^{-1}$ when the lines are extrapolated to zero dose for 2 M solutions (Fig. 2, lower). The data obtained from the analogous experiment with 0.2 M solutions gave $G(\text{HCHO}) = 0.16$ and $G(\text{CH}_3\text{CHO}) = 0.43 \mu\text{mol J}^{-1}$ when the lines are extrapolated to zero dose (Fig. 2, upper).





Fig. 2 Radiation chemical yields (G) of HCHO (●) and CH_3CHO (□) vs. dose from the experiments with 0.2 M (upper) and 2.0 M (lower) ethane-1,2-diol (*cf.* Fig. 1).

The experiment with 2 M ethane-1,2-diol produced $G(\text{CH}_3\text{CHO}) = 2.31 \mu\text{mol J}^{-1}$, which is 4–5 times more than $G(\mathbf{2}) = 0.53$. This result indicated a chain reaction for the formation of CH_3CHO .¹⁰ Scheme 3 illustrates our proposal, which involves H_2O elimination from the nucleophilic radical $\mathbf{2}$ with formation of the electrophilic radical $\mathbf{4}$ and the subsequent hydrogen abstraction from the starting material that completes the chain.



Scheme 3 Proposed mechanism for the reaction of $\text{HO}^\bullet/\text{H}^\bullet$ with ethane-1,2-diol (**1**). The compounds quantified by the 2,4-DNP procedures (Fig. 1) are shown in red.

The above findings need further comments. Asmus and co-workers some time ago investigated the site of HO^\bullet attack on aliphatic alcohols in aqueous solution using pulse radiolysis.¹⁵ The formation of reducing and oxidising radicals were identified by their reaction with $\text{C}(\text{NO}_2)_4$ and I^- , respectively. Thus, it was estimated that the reaction of HO^\bullet with CH_3OH occurs 93% from the CH_3 group and 7% from the OH site. On the other hand, the reaction of HO^\bullet with ethane-1,2-diol was assigned 100% to occur from the CH_2 moieties, because no oxidising species were obtained in the presence of $5 \times 10^{-4} \text{ M I}^-$. The findings that the formation of HCHO occurs in substantial quantities, suggests another scenario. On the basis of $G(\text{HCHO}) = 0.16$ and assuming that $G(\text{HCHO}) = 2 \times G(\mathbf{3})$, we calculated a $G(\mathbf{3}) = 0.08 \mu\text{mol J}^{-1}$ and consequently a $G(\mathbf{2}) = 0.53 \mu\text{mol J}^{-1}$ (cf. eqn (1) and (2)). Since H-atoms are expected to produce only radical $\mathbf{2}$, a $G(\mathbf{3}) = 0.08$ indicates that 14–15% of generated HO^\bullet radicals react with one of the two HO-groups of ethane-1,2-diol. This is in good agreement with the site of attack on CH_3OH discussed above. We also suggest that the fragmentation of $\mathbf{3}$ (oxidising species) to give HCHO and HOCH_2^\bullet (reducing species) is fast enough in aqueous solution to compete with the bimolecular oxidation of iodide ions, so that the product radicals are only reducing species.¹⁵ It is well-known that alkoxy radicals undergo facile β -fragmentation and 1,2-H-shift. For example, the β -fragmentation of the *tert*-butoxyl radical is $1.2 \times 10^6 \text{ s}^{-1}$ in water¹⁶ and the 1,2-H-shift in $\text{CH}_3\text{CH}_2\text{O}^\bullet$ is $(5 \pm 2) \times 10^6 \text{ s}^{-1}$ in water.¹⁷ The β -fragmentation of radical $\mathbf{3}$ is expected to be much faster than *tert*-butoxyl radical due to the formation of the stabilized HOCH_2^\bullet radical *versus* the CH_3^\bullet radical. The high reactivity of HO^\bullet radicals towards the OH moiety of ethane-1,2-diol is unsurprising. We have shown that for $\text{X-H} + \text{Y}^\bullet$ to give $\text{X}^\bullet + \text{H-Y}$, a major factor affecting the energy of activation is X-Y antibonding at the TS in addition to the enthalpy of reaction.¹⁸ It was suggested that abstraction from a OH moiety will have a

Table 2 Radiation chemical yields (G , $\mu\text{mol J}^{-1}$) for the reaction of $\text{HO}^\bullet/\text{H}^\bullet$ with propane-1,2-diol (**5**)

Products ^a	G from 0.2 M 5 , ^b	G from 2 M 5 , ^b
HCHO	0.10	0.08
CH_3CHO	0.11	0.17
Acetone	1.55	3.05
$\text{CH}_3\text{CH}_2\text{CHO}$	0.05	0.09

^a All compounds quantified by the 2,4-DNP procedure. ^b Values are extrapolated to zero dose (see ESI†).

relative low energy of activation, because O–O antibonding is low as a result of the weak O–O bond.

Propane-1,2-diol

Sample solutions (0.2 or 2 M) of propane-1,2-diol in unbuffered H_2O were saturated by N_2O prior to irradiation. Various irradiation doses up to 300 Gy and 2500 Gy for 0.2 and 2.0 M solutions, respectively, were employed. Derivatisation to a 2,4-DNP mixture and HPLC analysis indicated the formation of formaldehyde, acetaldehyde, acetone and propionaldehyde (Fig. 3) on comparison with authentic compounds. Analysis of the data in terms of radiation chemical yields is reported in Table 2.





Fig. 3 HPLC analyses of γ -irradiation of N_2O -saturated unbuffered solutions of 0.2 M (upper) and 2.0 M (lower) propane-1,2-diol at various doses (dose rate = 5.9 Gy min^{-1}) after 2,4-DNP derivatisation of the carbonyl compounds. The black and blue peaks correspond to formaldehyde and acetaldehyde, whereas the red and green peaks correspond to acetone and propionaldehyde, respectively.

Scheme 4 shows the mechanism we conceived for the reaction of $\text{HO}^\bullet/\text{H}^\bullet$ with propane-1,2-diol, taking into consideration the mechanism for ethane-1,2-diol discussed above (Scheme 3). H-atoms are expected to produce only radicals **6** and **7**, whereas HO^\bullet radicals abstract hydrogen atoms from all possible sites to give radicals **6–10**. The fact that $G(\text{acetone})$ increases from 1.55



Scheme 4 Proposed mechanism for the reaction of HO·/H· with propane-1,2-diol (**5**). The compounds quantified by the 2,4-DNP procedures (Fig. 3) are shown in red.

to 3.05 by increasing the concentration of starting material from 0.2 to 2 M, clearly indicates a chain reaction. Both radicals **6** and **7** undergo H₂O elimination to give the electrophilic radicals **11** and **12**, respectively, and their reaction with **5** gives mainly the most stable radical **7**. The remaining radicals **8**, **9**, and **10** are expected to afford the same products by β -fragmentation. This unimolecular process for **8** should be much slower than that for **9** and **10**. Asmus and co-workers¹⁵ reported that the reaction of HO· with propane-1,2-diol occurs 79.2% by abstraction of an H atom from the α position of the HO-group, and the remaining 20.7% from the methyl site. By analogy with the ethane-1,2-diol case, we suggest that 79.2% of the reducing species contain also the hydrogen abstraction from the OH site, because the fragmentation of **9** and **10** (oxidizing species) is fast enough in aqueous solution to compete with the bimolecular oxidation of iodide ions.

Butane-2,3-diol

Sample solutions (0.2 or 2 M) of butane-2,3-diol in unbuffered H₂O were saturated by N₂O prior to irradiation. Various irradiation doses up to 300 Gy and 1500 Gy for the 0.2 and 2.0 M solutions, respectively, were used. Derivatisation to a 2,4-DNP mixture and HPLC analysis indicated the formation of acetaldehyde, 3-hydroxybutan-2-one and 2-butanone (Fig. 4) by comparison with authentic compounds. Analysis of the data in terms of radiation chemical yields is reported in Table 3.

Scheme 5 shows the mechanism we conceived for the reaction of HO·/H· with butane-2,3-diol, taking into consideration the mechanisms for ethane-1,2-diol and propane-1,2-diol reported above (Scheme 1). H-atoms are expected to produce only radicals **14** and **15**, whereas HO· radicals abstract hydrogen atoms from all possible sites to give radicals **14**–**16**. The fact that $G(\text{CH}_3\text{CHO})$ and




Fig. 4 HPLC analyses of γ -irradiation of N₂O-saturated unbuffered solutions of 0.2 M at 300 Gy (upper) and 2.0 M at various doses (lower) of butane-2,3-diol (dose rate = 5.9 Gy min⁻¹) after 2,4-DNP derivatisation of the carbonyl compounds. The yellow and blue peaks correspond to 3-hydroxybutan-2-one (**18**) and acetaldehyde, whereas the magenta and grey peaks correspond to 2-butanone and the unknown carbonyl product, respectively.

Table 3 Radiation chemical yields (G , $\mu\text{mol J}^{-1}$) for the reaction of HO·/H· with butane-2,3-diol (**13**)

Products ^a	G from 0.2 M 13 , ^b	G from 2 M 13 , ^b
CH ₃ CHO	0.13	0.27
Butan-2-one	0.16	0.25
3-Hydroxybutan-2-one	0.08	0.08
Unknown carbonyl	0.03	0.04

^a All compounds quantified by the 2,4-DNP procedure. ^b Values are extrapolated to zero dose (see ESI[†]).

$G(\text{butan-2-one})$ are nearly doubled by increasing the concentration of starting material from 0.2 to 2 M, suggests that radical **14** undergoes H₂O elimination to give the electrophilic radical **17**, and its reaction with 2,3-diol **13** gives butan-2-one and both radicals **14** and **15**. The radicals **15** and **16** are expected to afford the same products by β -fragmentation, whereas 3-hydroxybutan-2-one (**18**) should result from the disproportionation of radical **15** followed by tautomerism. However, the presence of an unknown carbonyl derivative suggests an even more complex reaction scheme. Asmus and co-workers¹⁵ reported that the reaction of HO· with butane-2,3-diol occurs 71% by abstraction of an H atom from the α

- 5 W. Buckel and B. T. Golding, Radical Enzymes, in: *Encyclopedia of Radicals in Chemistry, Biology and Materials*, ed. C. Chatgililoglu and A. Studer, Wiley, Chichester, 2012.
- 6 S. A. Cockle, H. A. O. Hill, R. J. P. Williams, S. P. Davies and M. A. Foster, *J. Am. Chem. Soc.*, 1972, **94**, 275.
- 7 J. Rétey, A. Umami-Ronchi and D. Arigoni, *Experientia*, 1966, **22**, 72.
- 8 A. L. Buley, R. O. C. Norman and J. Pritchett, *J. Chem. Soc. B*, 1966, 849.
- 9 C. Walling and R. A. Johnson, *J. Am. Chem. Soc.*, 1975, **97**, 2405.
- 10 C. von Sonntag, *Free-Radical-Induced DNA Damage and Its Repair: A Chemical Perspective*, Springer-Verlag, Berlin, 2006.
- 11 B. T. Golding and L. Radom, *J. Am. Chem. Soc.*, 1976, **98**, 6331.
- 12 G. V. Buxton, C. L. Greenstock, W. P. Helman and A. B. Ross, *J. Phys. Chem. Ref. Data*, 1988, **17**, 513.
- 13 A. B. Ross, W. G. Mallard, W. P. Helman, G. V. Buxton, R. E. Huie and P. Neta, *NDRL-NIST Solution Kinetic Database - Ver. 3*, Notre Dame Radiation Laboratory, Notre Dame, IN and NIST Standard Reference Data, Gaithersburg, MD, 1998.
- 14 S. Barata-Vallejo, C. Ferreri, A. Postigo and C. Chatgililoglu, *Chem. Res. Toxicol.*, 2010, **23**, 258.
- 15 K. D. Asmus, H. Möckel and A. Henglein, *J. Phys. Chem.*, 1973, **77**, 1218.
- 16 M. Erben-Russ, W. Bors and M. Saran, *Int. J. Radiat. Biol.*, 1987, **52**, 393.
- 17 M. Bonifacic, D. A. Armstrong, I. Stefanic and K.-D. Asmus, *J. Phys. Chem. B*, 2003, **107**, 7268.
- 18 A. A. Zavitsas and C. Chatgililoglu, *J. Am. Chem. Soc.*, 1995, **117**, 10645.
- 19 β -Fragmentation and 1,2-H shift rate constants of alkoxy radicals are expected to increase in the same direction by methyl substitution, for example **16** vs. **5**.
- 20 T. Toraya, S. Honda and K. Mori, *Biochemistry*, 2010, **49**, 7210.
- 21 J. S. LaKind, E. A. McKenna, R. P. Hubner and R. G. Tardiff, *Crit. Rev. Toxicol.*, 1999, **29**, 331.
- 22 P. Navasumrit, T. H. Ward, N. J. F. Dodd and P. J. O'Connor, *Carcinogenesis*, 2000, **21**, 93.
- 23 C. Moncada, V. Torres, G. Varghese, E. Albano and Y. Israel, *Mol. Pharmacol.*, 1994, **46**, 786.
- 24 J. W. T. Spink and R. J. Woods, in: *An Introduction to Radiation Chemistry*, Wiley, New York, 3rd edn, 1990, p.100.