


Organic & Biomolecular Chemistry

www.rsc.org/obc

Volume 11 | Number 2 | 14 January 2013 | Pages 181–384



ISSN 1477-0520

RSC Publishing

COMMUNICATION

Tushar Kanti Chakraborty *et al.*

Total synthesis of (29S,37S)-isomer of malevamide E, a potent ion-channel inhibitor

Total synthesis of (29S,37S)-isomer of malevamide E, a potent ion-channel inhibitor†‡

Cite this: *Org. Biomol. Chem.*, 2013, **11**, 257

Received 3rd August 2012,
Accepted 11th September 2012

DOI: 10.1039/c2ob26533h

www.rsc.org/obc

The first total synthesis of (29S,37S)-malevamide E (**1**), a potent ion channel inhibitor, has been achieved in a convergent fashion involving Julia–Kocienski olefination, Urpi acetal aldol and Shiina macrolactonization reactions as the key steps. The strategy developed herein is amenable for the synthesis of the other possible isomers in search for the correct stereoisomer of the naturally occurring molecule.

Introduction


Extracts of cyanobacteria belonging to the genus *Symploca* (Oscillatoriaceae) have given rise to scores of biologically active natural products with interesting structures.¹ One such example is a new depsipeptide, malevamide E, an *N*-methylated analogue of dolastatin 14, isolated from field-collected colonies of the filamentous cyanobacterium *Symploca laete-viridis*.² Dolastatin 14 was previously reported by Pettit and co-workers³ in exceptionally low yield from *D.auricularia*. Malevamide E demonstrated a dose-dependent (2–45 μ M) inhibition of store-operated Ca^{2+} entry in thapsigargin-treated human embryonic kidney (HEK) cells indicating an inhibitory effect on Ca^{2+} release-activated Ca^{2+} (CRAC) channels.

Although, there has been considerable interest in the total synthesis of dolastatin 14, to date, neither the total synthesis nor the stereochemical configuration of dolastatin 14 or malevamide E has been reported. All four diastereomers of its polyketide-derived subunit, dolatrienoic acid were earlier synthesized as part of the effort toward the full synthesis of dolastatin 14 by two independent groups.⁴ Encouraged by the properties and coupled with our interest in total synthesis of marine natural products,⁵ we embarked on the challenge of

Praveen Kumar Gajula,^a Shrikant Sharma,^b Ravi Sankar Ampapathi^b and Tushar Kanti Chakraborty^{*a}

taking up a simple approach to make all possible diastereomers of the polyketide subunit. Herein, we report the first total synthesis of (29S,37S)-isomer of malevamide E (**1**).

Retrosynthetically (Scheme 1), (29S,37S)-malevamide E (**1**) can be obtained by macrolactonization of acyclic hydroxy acid **3**, which, in turn, could be prepared by coupling the two key intermediates **4** and **5**, the former with the long chain fatty



Scheme 1 Retrosynthetic analysis.

^aCSIR-Central Drug Research Institute, Lucknow-226 001, India.

E-mail: chakraborty@cdri.res.in

^bNMR Centre, SAI, CSIR-Central Drug Research Institute, Lucknow-226 001, India

†CDRI communication no. 8314.


‡Electronic supplementary information (ESI) available: Experimental procedure and characterisation data for all the new compounds. See DOI: 10.1039/c2ob26533h

acid moiety and the latter possessing the heptapeptide component. The salient feature of our synthesis is the stereoselective construction of the aliphatic fragment involving two key steps: (i) enantioselective addition of a titanium enolate derived from a chiral thiazolidinethione which was expected to fix the C29 stereocenter; and (ii) construction of C32–C33 bond using Julia–Kocienski olefination of the subunits 6 and 7.


Results and discussion

Our synthetic endeavour began with the synthesis of aldehyde 7 (Scheme 2) which started with the preparation of 4-(*tert*-butyldiphenylsiloxy)butanal dimethyl acetal 8 from butanediol according to the reported procedure.⁶ An acetal aldol reaction between the resultant dimethyl acetal 8 and (*S*)-*N*-acetyl-4-isopropyl-1,3-thiazolidine-2-thione (9), according to the conditions described by Urpi,⁷ provided the aldol product 10 in 72% yield in 3 : 1 dr in favour of the required isomer. The diastereomerically pure products could be easily separated using simple column chromatography.⁸ Reductive cleavage of the chiral auxiliary of the major isomer (the minor isomer can be used for the synthesis of another diastereomer of the molecule) afforded aldehyde, which was subjected to a Wittig–Horner reaction with *N*-methyldiethylphosphoacetamide (11) at $-40\text{ }^{\circ}\text{C}$ in presence of NaH to obtain the α,β -unsaturated amide 12 in 78% yield exclusively as an *E*-isomer.⁹ The amide 12 was reduced with DIBAL-H to the corresponding aldehyde followed by Wittig homologation with $\text{Ph}_3\text{P}=\text{CH}(\text{CH}_3)\text{CO}_2\text{Et}$ in benzene under reflux conditions to provide the α,β -unsaturated ester 13 in 80% yield. Desilylation of 13 with TBAF in THF gave a primary alcohol 14 in 75% yield, which on oxidation under Parikh–Doering conditions gave the aldehyde 7.

On the other hand, sulfone 6 was synthesized (Scheme 3) starting from a known compound 15,⁴ which was prepared by silyl protection of the corresponding alcohol. Hydroboration and oxidation of the terminal olefin moiety of 15 afforded primary alcohol 16 in 80% yield, which was further converted into the desired sulfone 6 in 85% yield *via* Mitsunobu reaction



Scheme 3 Synthesis of sulfone 6. (i) (a) BH_3SMe_2 , THF, $0\text{ }^{\circ}\text{C}$, rt, 2 h, (b) H_2O_2 , NaOH, 80%; (ii) (a) PTSH, DIAD, PPh_3 , THF, rt, 30 min, (b) $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot\text{H}_2\text{O}$, H_2O_2 , EtOH, rt, 15 min, 85%.




Scheme 4 Synthesis of compound 4. (i) LiHMDS, THF:HMPA, $-78\text{ }^{\circ}\text{C}$, aldehyde 7, $-78\text{ }^{\circ}\text{C}$, 3 h, 72%; (ii) TBAF:AcOH (1 : 1), H_2O , DMF, $50\text{ }^{\circ}\text{C}$, 12 h, 85%; (iii) LiOH, THF:MeOH: H_2O (3 : 1 : 1), $0\text{ }^{\circ}\text{C}$, rt, 1 h, 74%.


using 1-phenyl-1*H*-tetrazol-5-thiol (PT-SH) and diisopropyl azodicarboxylate (DIAD) and oxidation in one pot using catalytic amounts of ammonium molybdate and H_2O_2 .

Synthesis of compound 17, as depicted in Scheme 4, was performed through crucial Julia–Kocienski olefination.¹⁰ Accordingly, sulfone 6 was treated with LiHMDS in dry THF:HMPA (4 : 1) at $-78\text{ }^{\circ}\text{C}$ followed by slow addition of aldehyde 7, over a period of 10 min and the same temperature was maintained for 3 h to furnish the desired *E*-olefinic compound 17 in 72% yield along with 7% of the *Z*-isomer. An attempt to accomplish the deprotection of silyl group with TBAF in THF was ineffective and no desired product was obtained even at higher temperatures. Alternatively, desilylation of 17 using TBAF and AcOH (1 : 1) in presence of H_2O in DMF for 12 h gave deprotected 18 in 85% yield.¹¹ Saponification of 18 with LiOH in THF:MeOH: H_2O (3 : 1 : 1) at $0\text{ }^{\circ}\text{C}$ afforded corresponding hydroxy acid 4, which was used directly in the next step without further characterization.

The synthesis of the peptide segment, depicted in Scheme 5, started from the tetrapeptide 19, which was synthesized from commercially available protected amino acids by standard solution phase peptide synthesis conditions using 1-ethyl-3-(3-(dimethylamino)propyl)carbodiimide hydrochloride (EDCI) and 1-hydroxybenzotriazole hydrate (HOBT) as coupling agents and dry CH_2Cl_2 as solvent. For synthesis of 24, *N*-methylation of dipeptide 22 and tetrapeptide 19 was carried out using silver oxide and MeI in DMF to provide the corresponding *N*-methylated dipeptide 23 and tetrapeptide 20, respectively. From the 2D-TOCSY and ROESY spectrum of 20, at least 4 conformations were present in the NMR time scale out of which conformations 1A, 1B and 2A, 2B showed exchange peaks with each other (see ESI, Fig. 1†). For 1A,1B strong nOes between $^4\text{Pro H}_\delta \rightarrow ^3\text{Val H}_\alpha$ confirmed *trans* configuration and absence of such nOes for 2A, 2B suggests *cis*



Scheme 2 Synthesis of compound 7. (i) 9, TiCl_4 , DIPEA, SnCl_4 , CH_2Cl_2 , $-78\text{ }^{\circ}\text{C}$, 1 h, 72%; (ii) (a) DIBAL, CH_2Cl_2 , $-78\text{ }^{\circ}\text{C}$, 15 min, 96%; (b) 11, NaH, THF, $-40\text{ }^{\circ}\text{C}$, 1 h, 78%; (iii) (a) DIBAL, CH_2Cl_2 , $-78\text{ }^{\circ}\text{C}$, 15 min, 97%; (b) $\text{Ph}_3\text{P}=\text{C}(\text{CH}_3)\text{CO}_2\text{Et}$, benzene, reflux, 3 h, 80%; (iv) TBAF, THF, $0\text{ }^{\circ}\text{C}$, rt, 1 h, 85%; (v) $\text{SO}_3\cdot\text{Py}$, Et_3N , DCM: DMSO (2 : 1), $0\text{ }^{\circ}\text{C}$, rt, 30 min, 96%.



Scheme 5 Synthesis of peptide fragment and completion of the total synthesis. (i) Ag_2O , MeI, DMF, 0 °C, rt, 12 h, 72%; (ii) (a) LiOH, THF : MeOH : H_2O (3 : 1 : 1) 0 °C, rt, 1 h; (b) TFA salt of Boc-deprotected **23**, BOP-Cl, DIPEA, CH_2Cl_2 , 0 °C, rt, 12 h, 76%; (iii) Ag_2O , MeI, DMF, 0 °C, rt, 12 h, 68%; (iv) (a) TFA, CH_2Cl_2 , 0 °C, rt, 1 h; (b) Boc-D-Val-OH, BOP-Cl, DIPEA, CH_2Cl_2 , 0 °C, rt, 12 h, 79%; (v) (a) TFA, CH_2Cl_2 , 0 °C, rt, 1 h; (b) **4**, EDCI, HOBT, CH_2Cl_2 , 0 °C, rt, 15 min, then added amine from step (a), DIPEA, CH_2Cl_2 , rt, 12 h, 75%; (vi) LiOH, THF : MeOH : H_2O (3 : 1 : 1), 0 °C, rt, 1 h, 96%; (vii) MNBA, DMAP, CH_2Cl_2 , rt, 12 h, 62%.

configuration around proline peptide bond. Other two conformations may be attributed to the *cis/trans* isomerism of the one of the tertiary amides of **20** as these classes of peptides are prone to exhibit *cis/trans* conformations.¹² Coupling of the *N*-methylated tetrapeptide **20** with *N*-methylated dipeptide **23** using BOP-Cl gave the hexapeptide **21** in 76% yield.

Finally, Boc deprotection of **21** using TFA in CH_2Cl_2 followed by coupling with Boc-D-Val-OH using BOP-Cl as a coupling reagent gave the heptapeptide component **24**. NMR studies on hexapeptide **21** and heptapeptide **24** suggested presence of at least 8 conformations and may be attributed again to the *cis/trans* isomerisation of the tertiary amides (see ESI, Fig. 2 and 3†).

With both the halves of the molecule now ready to be coupled, the heptapeptide **24** was treated with 30% TFA in CH_2Cl_2 at 0 °C to effect Boc deprotection and coupled with the hydroxy acid **4** following the EDCI/HOBT protocol to obtain the coupled product **25** in 75% yield. The hydroxy ester **25** on saponification with LiOH in THF : MeOH : H_2O (3 : 1 : 1) at 0 °C afforded corresponding hydroxy acid **3**, which was used directly in the next step without further characterization. The stage was now set to carry out the crucial macrolactonization reaction. Lactonization was successfully carried out using Shiina anhydride protocol¹³ using 2-methyl-6-nitrobenzoic anhydride (MNBA) in presence of DMAP in CH_2Cl_2 at room temperature to afford the target molecule **1** in 62% yield.

NMR spectral data, both ^1H and ^{13}C , and specific rotation of the synthetic **1** were entirely not in agreement with the data reported for the isolated natural compounds. However, the detailed 2D-NMR analysis of **1** showed similar chemical shifts for the aliphatic region and deviated for certain residues of the peptoid portion.¹⁴ From the assigned major conformation, it is worth noting that chemical shifts of the Ala, Asn and Phe match well with the natural product.

In conclusion, we have developed a method for the total syntheses of (2*S*,3*S*)-isomer of malevamide **E** (**1**) in a convergent fashion involving Julia-Kocienski olefination, Urpi acetal aldol and Shiina macrolactonization as the key steps. Our strategy is amenable for the synthesis of other possible diastereomers whose synthesis is under progress and will be reported in due course.

Experimental section

Compound 25

To a solution of **18** (50 mg, 0.15 mmol) in THF : MeOH : H_2O (3 : 1 : 1, 3 mL) at 0 °C, LiOH· H_2O (18.5 mg, 0.44 mmol) was added and stirred from 0 °C to room temperature for 1 h. The reaction mixture was then acidified to pH ~ 2 with 1 N HCl. It was diluted with EtOAc, washed with brine, dried (Na_2SO_4), filtered and concentrated *in vacuo* to obtain the crude acid **4**, which was used directly in the next step without further characterization.

In round bottom flask, a solution of **24** (99.8 mg, 0.12 mmol) in CH_2Cl_2 (2 mL) was taken. To this solution, trifluoroacetic acid (1 mL) was added at 0 °C and slowly warmed to room temperature and stirred for 1 h. The reaction mixture was then concentrated *in vacuo* and azeotroped with CH_2Cl_2 3 times (3 × 10 mL) to give the Boc-deprotected compound **5**.

The above prepared acid **4** was dissolved in CH_2Cl_2 (3 mL) and cooled to 0 °C. It was then sequentially treated with HOBT (29.7 mg, 0.22 mmol) and EDCI (42.0 mg, 0.22 mmol). After 10 min, compound **5**, prepared above and dissolved in CH_2Cl_2 (2 mL), was added to the reaction mixture followed by the addition of DIPEA (0.08 mL, 0.44 mmol). After stirring for 12 h at room temperature, the reaction mixture was diluted with CHCl_3 , washed with saturated NH_4Cl solution, 1 N HCl, saturated NaHCO_3 solution, water, brine, dried (Na_2SO_4), filtered and concentrated *in vacuo*. Purification by column chromatography (silica gel, 2% to 3% MeOH in CHCl_3 as eluent) afforded **25** (100 mg, 75%) as a white solid. R_f = 0.3 (SiO_2 , 10% MeOH in CHCl_3); specific rotation $[\alpha]_D^{25} = -66.6$ (c 0.15, CHCl_3); IR (neat): ν_{max} 3404, 2361, 2111, 1637, 1445, 1218, 1020, 769, 669 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ see Table 6;† ^{13}C NMR (400 MHz, CDCl_3): δ 173.09, 171.82, 170.66, 170.21, 170.04, 169.55, 169.14, 168.32, 137.62, 137.16, 133.89, 130.68, 130.24, 130.14, 129.98, 129.54, 129.04, 128.59, 127.89, 79.97, 68.08, 60.04, 58.73, 56.92, 56.73, 55.47, 54.97, 53.72, 52.44, 52.27, 49.85, 47.36, 39.04, 38.89, 37.32, 35.73, 33.73, 32.60, 32.08, 31.88, 30.97, 30.06, 29.81, 29.46, 28.60, 28.30, 27.06, 26.47, 25.94, 25.07, 23.63, 22.79, 19.95, 19.81, 19.52, 19.44, 19.23, 18.09, 16.98, 14.26, 12.98; MS (ESIMS): m/z (%): 1157 (100) $[\text{M} + \text{Na}]^+$; HRMS (ESIMS): calcd for $\text{C}_{61}\text{H}_{99}\text{N}_8\text{O}_{12}$ $[\text{M} + \text{H}]^+$: 1135.7282, found: 1135.7255.

Compound 1

To a solution of **25** (55 mg, 0.05 mmol) in THF : MeOH : H_2O (3 : 1 : 1, 3 mL) at 0 °C, LiOH· H_2O (6.11 mg, 0.14 mmol) was added and stirred from 0 °C to room temperature for 1 h. The reaction mixture was then acidified to pH ~ 2 with 1 N HCl. It

was diluted with EtOAc, washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The hydroxy acid **3**, thus obtained, after flash chromatography (52 mg, 96%), was directly used in the next step without further characterization.

To a stirred solution of 2-methyl-6-nitrobenzoic anhydride (MNBA, 18.98 mg, 0.06 mmol) and DMAP (13.48 mg, 0.11 mmol) in CH₂Cl₂ (16 mL) was added a solution of hydroxy acid **3** (52 mg, 0.05) in CH₂Cl₂ (10 mL) over a period of 12 h at room temperature using a syringe pump. Thereafter, saturated NaHCO₃ solution (20 mL) was added and the aqueous layer was extracted with CHCl₃ (3 × 50 mL). The combined organic layers were dried with Na₂SO₄, filtered and concentrated *in vacuo*. Purification by column chromatography (silica gel, 2% to 3% MeOH in CHCl₃ as eluent) afforded **1** (31 mg, 62%) as a colorless oil. *R*_f = 0.3 (SiO₂, 10% MeOH in CHCl₃); specific rotation [α]_D²⁵ = -22.6 (*c* 0.09, CH₂Cl₂); IR (neat): ν_{\max} 3703, 3572, 3332, 2926, 2567, 2361, 1635, 1445, 1219, 1096, 770 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ see Table 8 in ESI; ¹³C NMR (150 MHz, CDCl₃): δ see Table 8 in the ESI; ⁺ MS (ESIMS): *m/z* (%): 1103 (100) [M + H]⁺; HRMS (ESIMS): calcd for C₆₀H₉₅N₈O₁₁ [M + H]⁺: 1103.7120, found: 1103.7085.

Acknowledgements

The authors wish to thank the UGC (P.K.G.) and CSIR, New Delhi (S.S.) for research fellowships. P.K.G thanks Dr Dipankar Koley for his support. The authors are thankful to SAIF, CSIR-CDRI for providing the spectroscopic and analytical data and also to Prof Wesley Yoshida, University of Hawaii, Manoa for providing the proton spectrum of malevamide E.

Notes and references

- (a) G. G. Harrigan, H. Luesch, W. Y. Yoshida, R. E. Moore, D. G. Nagle, V. J. Paul, S. L. Mooberry, T. H. Corbett and F. A. Valeriote, *J. Nat. Prod.*, 1998, **61**, 1075–1077; (b) G. G. Harrigan, H. Luesch, W. Y. Yoshida, R. E. Moore, D. G. Nagle and V. J. Paul, *J. Nat. Prod.*, 1999, **62**, 655–658; (c) H. Luesch, W. Y. Yoshida, R. E. Moore, V. J. Paul, S. L. Mooberry and T. Corbett, *J. Nat. Prod.*, 2002, **65**, 16–20; (d) F. D. Horgen, W. Y. Yoshida and P. J. Scheuer, *J. Nat. Prod.*, 2000, **63**, 461–467; (e) F. D. Horgen, E. B. Kazmierski, H. E. Westenburg, W. Y. Yoshida and P. J. Scheuer, *J. Nat. Prod.*, 2002, **65**, 487–491 The malevamide is named after the Hawaiian word “malewa” (ma lé va), which means “to sway”. M. K. Pukui and S. H. Elbert, *Hawaiian Dictionary, Revised and Enlarged Edition*, University of Hawaii Press, Honolulu, 1986, p. 232, referring to the graceful movement of the filamentous *Symploca* colonies caused by the ever-constant ocean swells; (f) P. G. Williams, W. Y. Yoshida, R. E. Moore and V. J. Paul, *J. Nat. Prod.*, 2002, **65**, 1336–1339; (g) P. G. Williams, W. Y. Yoshida, R. E. Moore and V. J. Paul, *J. Nat. Prod.*, 2003, **66**, 1006–1009; (h) P. G. Williams, W. Y. Yoshida, R. E. Moore and V. J. Paul, *J. Nat. Prod.*, 2003, **66**, 620–624; (i) P. G. Williams, W. Y. Yoshida, R. E. Moore and V. J. Paul, *J. Nat. Prod.*, 2004, **67**, 49–53; (j) T. L. Simmons, K. L. McPhail, E. Ortega-Barrá, S. L. Mooberry and W. H. Gerwick, *Tetrahedron Lett.*, 2006, **47**, 3387–3390.
- B. Adams, P. Pörzgen, E. Pittman, W. Y. Yoshida, H. E. Westenburg and F. D. Horgen, *J. Nat. Prod.*, 2008, **71**, 750–754.
- G. R. Pettit, Y. Kamano, C. L. Herald, C. Dufresne, R. B. Bates, J. M. Schmidt, R. L. Cerny and H. Kizu, *J. Org. Chem.*, 1990, **55**, 2989–2990.
- (a) S. Mouné, G. Niel, M. Busquet, I. Eggleston and P. Jouin, *J. Org. Chem.*, 1997, **62**, 3332–3339; (b) J. J. Duffield and G. R. Pettit, *J. Nat. Prod.*, 2001, **64**, 472–479.
- (a) T. K. Chakraborty, V. R. Reddy and P. K. Gajula, *Tetrahedron*, 2008, **64**, 5162–5167; (b) T. K. Chakraborty, V. R. Reddy and A. K. Chattopadhyay, *Tetrahedron Lett.*, 2006, **47**, 7435–7438; (c) T. K. Chakraborty and V. R. Reddy, *Tetrahedron Lett.*, 2006, **47**, 2099–2102; (d) T. K. Chakraborty and S. Das, *Tetrahedron Lett.*, 2001, **42**, 3387–3390; (e) T. K. Chakraborty and S. Das, *Chem. Lett.*, 2000, 80–81; (f) T. K. Chakraborty, S. Ghosh and S. Dutta, *Tetrahedron Lett.*, 2001, **42**, 5085–5088; (g) T. K. Chakraborty, S. Ghosh, P. Laxman, S. Dutta and R. Samanta, *Tetrahedron Lett.*, 2005, **46**, 5447–5550; (h) T. K. Chakraborty and V. R. Suresh, *Tetrahedron Lett.*, 1998, **39**, 9109–9112; (i) T. K. Chakraborty and V. R. Suresh, *Tetrahedron Lett.*, 1998, **39**, 7775–7778; (j) T. K. Chakraborty and D. Thippeswamy, *Synlett*, 1999, 150–152; (k) T. K. Chakraborty, D. Thippe swamy, V. R. Suresh and S. Jayaprakash, *Chem. Lett.*, 1997, 563.
- H. Okada, T. Mori, Y. Saikawa and M. Nakata, *Tetrahedron Lett.*, 2009, **50**, 1276–1278.
- (a) A. Cosp, E. Llacer, P. Romea and F. Urpí, *Tetrahedron Lett.*, 2006, **47**, 5819–5823; (b) A. Cosp, P. Romea, F. Urpí and J. Vilarrasa, *Tetrahedron Lett.*, 2001, **42**, 4629–4631.
- To prove the stereochemistry of adduct, compound **10** was converted to a known compound (+)-(3*S*)-1,6-Dihydroxy-3-methoxy-hexane, the spectral data of which was matched with the reported data. For reference see: P. Ackermann, H. Tobler and C. Ganter, *Helv. Chim. Acta*, 1972, **55**, 2731–2743.
- T. J. Heckrodt and J. Mulzer, *J. Am. Chem. Soc.*, 2003, **125**, 4680–4681.
- J. Liu, K. Xu, J. He, L. Zhang, X. Pan and X. She, *J. Org. Chem.*, 2009, **74**, 5063–5066.
- H. Higashibayashi, K. Shinko, T. Ishizu, K. Hashimoto, H. Shirahama and M. Nakata, *Synlett.*, 2000, 1306–1308.
- Q. Sui, D. Borchardt and D. L. Rabenstein, *J. Am. Chem. Soc.*, 2007, **129**, 12042–12048.
- (a) I. Shiina, M. Kubota, H. Oshiumi and M. Hashizume, *J. Org. Chem.*, 2004, **69**, 1822–1830; (b) I. Shiina, M. Hashizume, Y. S. Yamai, H. Oshiumi, T. Shimazaki, Y. J. Takasuna and R. Ibuka, *Chem.–Eur. J.*, 2005, **11**, 6601–6608; (c) For a review, see: .I. Shiina, *Chem. Rev.*, 2007, **107**, 239–273.
- For δ values see ESI Table 8.†