

Asymmetric Synthesis of the Balanol Heterocycle via a Palladium-Mediated Epimerization and Olefin Metathesis


Gregory R. Cook,* P. Sathya Shanker, and Scott L. Peterson

Department of Chemistry, North Dakota State University,
Fargo, North Dakota 58105-5516

grcook@plains.nodak.edu

Received June 2, 1999

ABSTRACT



The enantioselective formal synthesis of balanol, a potent protein kinase C inhibitor, was accomplished from D-serine utilizing a Pd-catalyzed equilibration of diastereomeric 5-vinyloxazolines to set the stereochemistry of the vicinal amino and hydroxyl groups. A ruthenium-catalyzed ring-closing metathesis was employed to form the seven-membered nitrogen heterocycle.

Balanol (**1**, Figure 1) was first isolated in 1993 from the fungus *Verticillium balanoides*¹ and again in 1994 from a species of *Fusarium*.² It is a potent inhibitor of human protein kinase C displaying IC₅₀ values of 4–9 nM.³ Due to its high inhibitory activity, balanol has been the target of a number of synthetic efforts. Most recently, a samarium-mediated radical cyclization of an oxime ether afforded the nitrogen heterocycle in racemic form as a 6.6:1 mixture of diastereomers, and the major isomer was subsequently resolved at a later stage.⁴ The Nicolaou group has reported an asymmetric synthesis of balanol beginning with D-serine. Key to the synthesis was a chiral allylation of a protected serinal with

(Ipc)₂B-allyl to afford a 12:1 mixture of diastereomers.⁵ The Tanner group utilized regio- and stereoselective opening of chiral epoxides and aziridines to control the vicinal amino alcohol stereochemistry.⁶ Preparation of (2*S*,3*R*)-3-hydroxylysine in enantiomerically pure form⁷ was key to the total synthesis of balanol reported by Lampe and Hughes.⁸ Other total syntheses relying on chiral resolution of racemates have





Figure 1.

- (1) Kulanthaivel, P.; Hallock, Y. F.; Boros, C.; Hamilton, S. M.; Janzen, W. P.; Ballas, L. M.; Loomis, C. R.; Jiang, J. B. *J. Am. Chem. Soc.* **1993**, *115*, 6452.
- (2) Ohshima, S.; Yanagisawa, M.; Katoh, A.; Fujii, T.; Sano, T.; Matsukuma, S.; Furumai, T.; Fujii, M.; Watanabe, K.; Yokose, K.; Arisawa, M.; Okuda, T. *J. Antibiot.* **1994**, *47*, 639.
- (3) Boros, C.; Hamilton, S. M.; Katz, B.; Kulanthaivel, P. *J. Antibiot.* **1994**, *47*, 1010.
- (4) (a) Miyabe, H.; Torieda, M.; Inoue, K.; Tajiri, K.; Kiguchi, T.; Naito, T. *J. Org. Chem.* **1998**, *63*, 4397. (b) Miyabe, H.; Torieda, M.; Kiguchi, T.; Naito, T. *Synlett* **1997**, 580. (c) Naito, T.; Torieda, M.; Tajiri, K.; Ninomiya, I.; Kiguchi, T. *Chem. Pharm. Bull.* **1996**, *44*, 624.
- (5) Nicolaou, K. C.; Bunnage, M. E.; Koide, K. *J. Am. Chem. Soc.* **1994**, *116*, 8402.
- (6) (a) Tanner, D.; Almario, A.; Högberg, T. *Tetrahedron* **1995**, *51*, 6061. (b) Tanner, D.; Tedenborg, L.; Almario, A.; Pettersson, I.; Csöreg, I.; Kelly, N. M.; Andersson, P. G.; Högberg, T. *Tetrahedron* **1997**, *53*, 4857.
- (7) (a) Hughes, P. F.; Smith, S. H.; Olson, J. T. *J. Org. Chem.* **1994**, *59*, 5799. (b) Coulon, E.; Caño de Andrade, M. C.; Ratovelomanana-Vidal, V.; Genêt, J.-P. *Tetrahedron Lett.* **1998**, *39*, 6467.
- (8) (a) Lampe, J. W.; Hughes, P. F.; Biggers, C. K.; Smith, S. H.; Hu, H. *J. Org. Chem.* **1994**, *59*, 5147. (b) Lampe, J. W.; Hughes, P. F.; Biggers, C. K.; Smith, S. H.; Hu, H. *J. Org. Chem.* **1996**, *61*, 4572.


appeared.^{8a,9} The hexahydroazepine **2** (Figure 1) has been targeted in several formal syntheses of balanol.¹⁰

We have recently shown¹¹ that chiral 5-vinylloxazolidinones (**3**) derived from α -amino acids react with palladium(0) catalysts to afford 5-vinylloxazolines (**6**) via oxidative insertion, loss of CO₂, and subsequent cyclization of the amide oxygen (Scheme 1). The oxazoline products were obtained



with enhanced diastereomeric ratios, which suggested that the intermediate π -allyl palladium complexes **4** and **5** were undergoing equilibration. Oxazolines **6** were also ionized by the palladium catalyst and were in equilibrium with **4** and **5**, thus giving rise to thermodynamic product ratios.¹² We envisioned that this equilibration could be utilized to set the vicinal amino alcohol stereochemistry of the balanol heterocycle. The pendant vinyl group could be employed in a transition metal catalyzed ring-closing metathesis to generate the hexahydroazepine ring. This approach to the synthesis of **2** is presented in this Letter.


The synthesis of **2** began with the protected D-serine derivative **7** as shown in Scheme 2. Diisobutylaluminum hydride reduction of the ester and addition of vinylmagnesium bromide to the crude aldehyde afforded the allylic



^a (a) DIBAL-H, Tol, -78 °C; (b) vinylmagnesium bromide, THF, 58% for two steps; (c) NaH, THF, 91%; (d) 4-BnO-C₆H₄COCl, NaH, 90%; (e) Pd₂(dba)₃CHCl₃ (1 mol %), dppp (4 mol %), THF, 35 °C, 79% isolated pure *trans* diastereomer.

alcohol **8**^{13,14} in 58% yield (two steps) as a 67:33 mixture of *syn* and *anti* diastereomers, respectively.^{15,16} Oxazolidinone formation was achieved by treatment with sodium hydride in THF, and acylation gave the required 5-vinylloxazolidinone **9** in high yield. The palladium-catalyzed oxazoline formation/epimerization provided **10** as a 94:6 (*trans*:*cis*) mixture of diastereomers.¹⁵ The isolated yield of the purified major diastereomer was 79% (*cis* isomer, 3%). The optimal epimerization conditions were 35 °C in THF. When toluene was employed as the solvent, a slightly lower selectivity was obtained. Higher temperatures resulted in the formation of trace amounts of elimination products with no improvement in the diastereomer ratio.

Removal of the *tert*-butyldimethylsilyl protecting group from **10** with tetrabutylammonium fluoride afforded the free alcohol in 83% yield (Scheme 3). Mitsunobu reaction with



^a (a) TBAF, THF, 83%; (b) DPPA, Ph₃P, DEAD, 88%; (c) Ph₃P, THF-H₂O; then BnOCCl, Et₃N, 70%; (d) NaH, allyl bromide, 87% (11% recovered starting material); (e) Cl₂(PCy₃)₂Ru=CHPh (2 × 5 mol %), 45 °C, CH₂Cl₂, 77%; (f) KO₂CNNCO₂K, AcOH, 50%; (g) 1 N HCl, EtOH, 60 °C; then excess Et₃N, MeOH, rt, 63%; (h) 2 N HCl, THF, rt; then excess Et₃N, MeOH, rt, 72%; (i) (Ph₃P)₃RhCl (5 mol %) H₂ (25 psi), benzene, 94%.

diphenylphosphoryl azide¹⁷ was employed to introduce the nitrogen in 88% yield (**11**). Reduction with triphenylphos-

(9) Adams, C. P.; Fairway, S. M.; Hardy, C. J.; Hibbs, D. E.; Hursthouse, M. B.; Morley, A. D.; Sharp, B. W.; Vicker, N.; Warner, I. *J. Chem. Soc. Perkin Trans. 1* **1995**, 2355.

(10) For asymmetric syntheses of protected (3*R*,4*R*)-3-amino-4-hydroxyhexahydroazepines, see: (a) Müller, A.; Takyar, D. K.; Witt, S.; König, W. A. *Liebigs Ann. Chem.* **1993**, 651. (b) Tuch, A.; Saniere, M.; Le Merrer, Y.; Depezay, J.-C. *Tetrahedron: Asymmetry* **1996**, 7, 2901. (c) Wu, M. H.; Jacobsen, E. N. *Tetrahedron Lett.* **1997**, 38, 1693. (d) Albertini, E.; Barco, A.; Benetti, S.; De Risi, C.; Pollini, G. P.; Zanirato, V. *Tetrahedron* **1997**, 53, 17177. (e) Morie, T.; Kato, S. *Heterocycles* **1998**, 48, 427.

(11) Cook, G. R.; Shanker, P. S. *Tetrahedron Lett.* **1998**, 39, 3405.

(12) A similar Pd-mediated thermodynamic equilibration of vinyl aziridines has been reported: (a) Ibuka, T.; Mimura, N.; Aoyama, H.; Akaji, M.; Ohno, H.; Miwa, Y.; Taga, T.; Nakai, K.; Tamamura, H.; Fujii, N.; Yamamoto, Y. *J. Org. Chem.* **1997**, 62, 999. (b) Ibuka, T.; Mimura, N.; Aoyama, H.; Ohno, H.; Nakai, K.; Akaji, M.; Habashita, H.; Tamamura, H.; Miwa, Y.; Taga, T.; Fujii, N.; Yamamoto, Y. *J. Org. Chem.* **1997**, 62, 2982.

phine and water, protection with benzyl chloroformate, and allylation with allyl bromide gave the diene **12**. The olefin metathesis reaction with 10 mol % of the Grubbs ruthenium alkylidene catalyst¹⁸ provided the desired seven-membered nitrogen heterocycle **13** in a respectable yield (77%). The experimental conditions for the olefin metathesis reaction are worthy of comment. Optimal results were obtained when 5 mol % of the catalyst was used followed by the addition of another 5 mol % after 4 h. The reaction would not proceed to completion with a single loading of the catalyst, even up to 20 mol %. Presumably, the catalyst decomposed and an additional loading was required to obtain complete conversion.

Diimide reduction of the olefin was accomplished with potassium azodicarboxylate in acetic acid to afford the saturated nitrogen heterocycle **14** in a moderate yield. Hydrolysis of the oxazoline was carried out with 1 N HCl in ethanol at 60 °C. The formation of **2** required subsequent treatment with base. Under simple aqueous workup conditions, the benzoate derivative was isolated as the amine hydrochloride salt. Stirring with triethylamine in methanol for 36 h was necessary to effect the benzoyl transfer to the amine. A higher overall yield from **13** to **2** was realized by reversing the hydrolysis and reduction steps. Acidic hydroly-

sis and base treatment afforded **15** in 72% yield, and hydrogenation with Wilkinson's catalyst gave **2** with no observed cleavage of the protecting groups.

We have described an asymmetric synthesis of the balanol heterocycle from D-serine which incorporated a palladium-mediated equilibration to set the relative stereochemistry. Seven-membered ring formation was accomplished with a ruthenium-catalyzed olefin metathesis reaction in high yield. The utility of intermediate **13** for the synthesis of balanol analogues¹⁹ and novel azasugar derivatives is currently under investigation and will be reported in due course.

Acknowledgment. We thank NSF (OSR-9452892, CHE-9875013), NIH (GM58470-01), and North Dakota State University for support of our programs. We are grateful to Boulder Scientific, Inc. for a generous gift of Grubbs' catalyst. We thank Dr. Peter Wuts for helpful discussions.

Supporting Information Available: Experimental procedures and characterization data for compounds **2** and **9–15** and unnumbered intermediates isolated. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL990705+

(13) The alcohol **8** is a known compound. See ref 12a and references therein.

(14) All compounds displayed satisfactory spectroscopic (NMR, IR) data consistent with their structures.

(15) Diastereomer ratios were determined by ¹H NMR (400 MHz).

(16) Careful chelation controlled addition of vinylolithium reagents to aminoaldehydes generated in situ from DIBAL/TRIBAL in non-ether solvents have been reported to proceed with high (>20:1) selectivity. (a) Sames, D.; Polt, R. *J. Org. Chem.* **1994**, *59*, 4596. See also: (b) Angle, S. R.; Henry, R. H. *J. Org. Chem.* **1997**, *62*, 8549. (c) Ibuka, T.; Habashita, H.; Otaka, A.; Fujii, N.; Oguchi, Y.; Ueyehara, T.; Yamamoto, Y. *J. Org. Chem.* **1991**, *56*, 4370.

(17) Lal, B.; Pramanik, B. N.; Manhas, M. S.; Bose, A. K. *Tetrahedron Lett.* **1977**, 1977.

(18) For recent reviews on ring-closing metathesis, see: (a) Schuster, M.; Blechert, S. *Angew. Chem.* **1997**, *109*, 2144; *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2036. (b) Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413. (c) Armstrong, S. K. *J. Chem. Soc., Perkin Trans. 1* **1998**, 371.

(19) For studies on analogues of balanol, see: (a) Heerding, J. M.; Lampe, J. W.; Darges, J. W.; Stamper, M. L. *Biorg. Med. Chem. Lett.* **1995**, *5*, 1839. (b) Jagdmann, G. E., Jr.; Defauw, J. M.; Lai, Y.-S.; Crane, H. M.; Hall, S. E.; Buben, J. A.; Hu, H.; Gosnell, P. A. *Biorg. Med. Chem. Lett.* **1995**, *5*, 2015. (c) Lai, Y.-S.; Menaldino, D. S.; Nichols, J. B.; Jagdmann, G. E., Jr.; Mylott, F.; Gillespie, J.; Hall, S. E. *Biorg. Med. Chem. Lett.* **1995**, *5*, 2151. (d) Lai, Y.-S.; Mendoza, J. S.; Hubbard, F.; Kalter, K. *Biorg. Med. Chem. Lett.* **1995**, *5*, 2155. (e) Mendoza, J. S.; Jagdmann, G. E., Jr.; Gosnell, P. A. *Biorg. Med. Chem. Lett.* **1995**, *5*, 2211. (f) Hu, H.; Hollinshead, S. P.; Hall, S. E.; Kalter, K.; Ballas, L. M. *Biorg. Med. Chem. Lett.* **1996**, *6*, 973. (g) Jagdmann, G. E., Jr.; Defauw, J. M.; Lampe, J. W. *Biorg. Med. Chem. Lett.* **1996**, *6*, 1759. (h) Defauw, J. M.; Murphy, M. M.; Jagdmann, G. E., Jr.; Hu, H.; Lampe, J. W.; Hollinshead, S. P.; Mitchell, T. J.; Crane, H. M.; Heerding, J. M.; Mendoza, J. S.; Davis, J. E.; Dargess, J. W.; Hubbard, F. R.; Hall, S. E. *J. Med. Chem.* **1996**, *39*, 5215. (i) Nielsen, J.; Lyngsø, L. O. *Tetrahedron Lett.* **1996**, *37*, 8439. (j) Lai, Y.-S.; Mendoza, J. S.; Jagdmann, G. E., Jr.; Menaldino, D. S.; Biggers, C. K.; Heerding, J. M.; Wilson, J. W.; Hall, S. E.; Jiang, J. B.; Janzen, W. P.; Ballas, L. W. *J. Med. Chem.* **1997**, *40*, 226.

