

A Self-Assembled Porphyrin-Based Dimeric Capsule Constructed by a Pd(II)–Pyridine Interaction Which Shows Efficient Guest Inclusion


Atsushi Ikeda,^{*,†} Masatsugu Ayabe,[†] Seiji Shinkai,^{*,†} Shigeru Sakamoto,[‡] and Kentaro Yamaguchi[‡]

Department of Chemistry & Biochemistry, Graduate School of Engineering, Kyushu University, Fukuoka 812-8581, Japan, and Chemical Analysis Center, Chiba University, Chiba 263-8522, Japan

seijitcm@mbox.nc.kyushu-u.ac.jp

Received September 19, 2000

ABSTRACT



Novel self-assembled molecular capsules were constructed from two moles of pyridine-containing porphyrin derivatives and four moles of Pd(II) complexes utilizing a pyridine–Pd(II) interaction. The ¹H NMR spectral studies established that these self-assembled molecular capsules 5 and 6 have a highly symmetrical *D*_{4h} structure as well as a large inside cavity. It was shown that molecular capsule 6 can include a large bipyridine guest by a two-point simultaneous pyridine–Zn(II) interaction.

A great deal of effort has been devoted toward multiporphyrin arrays such as molecular wires,¹ molecular switches,² photosynthetic systems,³ photosensitizers for DNA cleavage,⁴ and photocurrent generation.⁵ A combination of these excellent devices with host–guest chemistry has a large future

potential but is much less developed so far. A few cyclic host compounds composed of covalently linked multiporphyrins have been synthesized, and some of them can include guest molecules such as pyridine derivatives^{6,7} and fullerenes.⁸ The findings suggest that molecular capsules with a three-dimensional cavity would have larger association constants and kinetically slower exchange rates for specific guest molecules than cyclic compounds with a two-dimensional cavity, but the syntheses of such molecular capsules are

[†] Kyushu University.

[‡] Chiba University.

(1) (a) Anderson, H. L. *Chem. Commun.* **1999**, 2323–2330. (b) Wagner, R. W.; Lindsey, J. S. *J. Am. Chem. Soc.* **1994**, *116*, 9759–9760. (c) Aratani, N.; Osuka, A.; Kim, Y. H.; Jeong, D. H.; Kim, D. *Angew. Chem., Int. Ed.* **2000**, *39*, 1458–1462. (d) Tsuda, A.; Furuta, H.; Osuka, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 2549–2552.


(2) Gosztola, D.; Niemczyk, M. P.; Wasielewski, M. R. *J. Am. Chem. Soc.* **1998**, *120*, 5118–5119.

(3) (a) Wasielewski, M. R. *Chem. Rev.* **1992**, *92*, 435–461. (b) Steinbergfrach, G.; Liddell, P. A.; Hung, S. C.; Moore, A. L.; Gust, D.; Moore, T. A. *Nature* **1997**, *385*, 239–241. (c) Martín, N.; Sánchez, L.; Illescas, B.; Pérez, I. *Chem. Rev.* **1998**, *98*, 2527–2547. (d) Van Patten, P. G.; Shreve, A. P.; Lindsey, J. S.; Donohoe, R. J. *J. Phys. Chem. B* **1998**, *102*, 4209–4216.

(4) (a) Sousa, C.; Maziere, C.; Maziere, J. C. *Cancer Lett.* **1998**, *128*, 177–182. (b) deVree, W. J. A.; Essers, M. C.; Sluiter, W. *Cancer Res.* **1997**, *57*, 2555–2558. (c) Jasat, A.; Dolphin, D. *Chem. Rev.* **1997**, *97*, 2267–2340. (d) Suenaga, H.; Nakashima, K.; Mizuno, T.; Takeuchi, M.; Hamachi, I.; Shinkai, S. *J. Chem. Soc., Perkin Trans. 1* **1998**, 1263–1267.

(5) (a) Uosaki, K.; Kondo, T.; Zhang, XQ.; Yanagida, M. *J. Am. Chem. Soc.* **1997**, *119*, 8367–8368. (b) Imahori, H.; Yamada, H.; Ozawa, S.; Ushida, K.; Sakata, Y. *Chem. Commun.* **1999**, 1165–1166. (c) Imahori, H.; Yamada, H.; Nishimura, Y.; Yamazaki, I.; Sakata, Y. *J. Phys. Chem. B* **2000**, *104*, 2099–2108.

frequently very troublesome. Here, it occurred to us that the utilization of coordination bonds has been somewhat neglected, given that Fujita et al.⁹ and Stang et al.¹⁰ have shown a number of attractive examples in which coordination bonds are employed for the construction of self-assembled supramolecular architectures.¹¹ We previously found that two homooxalix[3]arenes dimerize with three *cis*-Pd(II) complexes into a molecular capsule according to a self-assembled manner.^{12,13} The spectroscopic studies have shown that the molecular capsule thus formed can specifically include [60]-fullerene, the selectivity of [60]fullerene vs [70]fullerene being nearly “perfect”.^{12,14} Utilizing this class of concept, several self-assembled multiporphyrin arrays were recently reported in organic solvents.¹⁵ However, most of the preceding examples are two-dimensional macrocycles without a sufficient inclusion cavity, whereas the examples for three-dimensional molecular capsules have been very limited.^{16,17} If such a novel molecular capsule with “porphyrin walls” is successfully constructed, it follows that a guest is shielded inside the cavity while electrons are injected only via these “porphyrin walls”. With this object in mind, we here report novel self-assembled molecular capsules constructed from porphyrin **1** or **2** through the pyridyl–Pd(II) interaction. Very interestingly, we have found that these molecular capsules can include bipyridine derivatives with relatively large association constants.



Compound **1** was obtained¹⁸ in 22% yield by the reaction of 3-methoxy-4-*n*-octyloxy-5-pyridin-4-yl ethynyl benzaldehyde

(6) Sanders et al. have reported that a large number of multiporphyrin hosts can include pyridine guests by a multipoint simultaneous pyridine–Zn(II) interaction: (a) Anderson, S.; Anderson, H. L.; Sanders, J. K. M. *J. Chem. Soc., Perkin Trans. 1* **1995**, 2255–2267. (b) Vidal-Ferran, A.; Clyde-Watson, Z.; Bampos, N.; Sanders, J. K. M. *J. Org. Chem.* **1997**, *62*, 240–241. (c) Nakash, M.; Clyde-Watson, Z.; Feeder, N.; Davies, J. E.; Teat, S. J.; Sanders, J. K. M. *J. Am. Chem. Soc.* **2000**, *122*, 5286–5293 and references therein.

(7) Ambrose, A.; Li, J.; Lianhe Yu, Lindsey, J. S. *Org. Lett.* **2000**, *2*, 2563–2566.

(8) Tashiro, K.; Aida, T.; Zheng, J. Y.; Kinbara, K.; Saigo, K.; Sakamoto, S.; Yamaguchi, K. *J. Am. Chem. Soc.* **1999**, *121*, 5645–5647.

(9) (a) Fujita, M.; Yazaki, J.; Ogura, K. *J. Am. Chem. Soc.* **1990**, *112*, 5645–5647. (b) Fujita, M.; Nagao, S.; Ogura, K. *J. Am. Chem. Soc.* **1995**, *117*, 1649–1650.

(10) (a) Stang, P. J.; Cao, D. H. *J. Am. Chem. Soc.* **1994**, *116*, 4981–4982. (b) Stang, P. J.; Cao, D. H.; Saito, S.; Arif, A. M. *J. Am. Chem. Soc.* **1995**, *117*, 6273–6283.

with pyrrole in propionic acid. Compound **2** was obtained in 99% yield by the reaction of **1** with Zn(OAc)₂. These compounds were identified by IR, ¹H NMR, and MALDI-TOF mass ($[M + H]^+ = 1651.9$ and 1713.8 for **1** and **2**, respectively) spectral evidence and elemental analyses.

As shown in Figure 1b, the simple ¹H NMR splitting pattern was obtained when **1** and **3** were mixed in a 1:2 ratio




Figure 1. Partial ¹H NMR spectra of (a) **1** (2.2 mM) and (b) [**1**]:[**3**] = 1:2 (2.2 mM/4.4 mM): CDCl₃, 27 °C, 600 MHz.

in CDCl₃. When the ratio was higher or lower than this value, the ¹H NMR spectra gave additional peaks and became very complicated. Careful examination of Figure 1b and the ¹H–¹H COSY spectrum reveals that all peaks of **1**·**3** complex can be assigned to one kind of signals, supporting the 2:4 **1**/**3** complex (**5**) with a *D*_{4h} symmetrical structure but not the 1:2 complex with a *C*_{2v} symmetrical structure as inconceivable because the rigid tetraphenylporphyrin skeleton of **1** suppresses the pyridyl groups to get close to each other and thus prevents **1** from the formation of the intramolecular bonds with two *cis*-Pd(II) complexes. Meanwhile, a solution of **2** in CDCl₃ gave a very complicated and very broadened ¹H NMR spectrum, suggesting that the pyridyl groups act as axial ligands to bind Zn(II) intermolecularly.¹⁹ When **3** was added, **2** gave a ¹H NMR spectral splitting pattern very

(11) Fox, O. D.; Drew, M. G. B.; Wilkinson, E. J. S.; Beer, P. D. *Chem. Commun.* **2000**, 391–392.

(12) Ikeda, A.; Yoshimura, M.; Tani, F.; Naruta, Y.; Shinkai, S. *Chem. Lett.* **1998**, 587–588.

(13) Ikeda, A.; Yoshimura, M.; Udzu, H.; Fukuhara, C.; Shinkai, S. *J. Am. Chem. Soc.* **1999**, *121*, 4296–4297.

(14) Ikeda, A.; Udzu, H.; Yoshimura, M.; Shinkai, S. *Tetrahedron* **2000**, *56*, 1825–1832.

(15) (a) Drain, C. M.; Lehn, J. M. *J. Chem. Soc., Chem. Commun.* **1994**, 2313–2314. (b) Stang, P. J.; Fan, J.; Olenyuk, B. *Chem. Commun.* **1997**, 1453–1454. (c) Fan, J.; Whiteford, J. A.; Olenyuk, B.; Levin, M. D.; Stang, P. J.; Fleischer, E. B. *J. Am. Chem. Soc.* **1999**, *121*, 2741–2752.

(16) Slone, R. V.; Hupp, J. T. *Inorg. Chem.* **1997**, *36*, 5422–5423.

(17) Fujita, N.; Fujita, M.; Sakamoto, S.; Yamaguchi, K. *Abstracts, XI International Symposium on Supramolecular Chemistry, Fukuoka, Japan, 2000*; PD-37.

(18) Synthetic details and characterization data relating to **1** and **2** can be found in the Supporting Information.

(19) Supramolecular complexes consisting of tripyridine derivatives and tris{Zn(II) porphyrin} derivatives were reported. However, they cannot include the guest molecules in their cavities utilizing Zn(II)–metal coordination: (a) Felluga, F.; Tecilla, P.; Hillier, L.; Hunter, C. A.; Licini, G.; Scrimin, P. *Chem. Commun.* **2000**, 1087–1088. (b) Ikeda, A.; Sonoda, K.; Shinkai, S. *Chem. Lett.* **2000**, 1220–1221.




Figure 2. Schematic illustration of 1:2 and 2:4 complexes.

similar to that of 2:4 1/3 complex (Figure 3b), which can be also assigned to the molecular capsule **6**.

The formation of molecular capsule **6** was also supported by coldspray ionization mass spectrometry (CIS-MS).²⁰ When a CH₂Cl₂ solution containing **2** and **3** in a 1:2 ratio was subjected to the CIS-MS measurement, strong peaks appeared at 967, 1190, and 1525, which are assignable to [6 - 6CF₃SO₃⁻]⁶⁺, [6 - 5CF₃SO₃⁻]⁵⁺, and [6 - 4CF₃SO₃⁻]⁴⁺, respectively.




Figure 3. Partial ¹H NMR spectra of (a) **2** (1.5 mM) and (b) [**2**]:[**3**] = 1:2 (1.5 mM/3.0 mM): CDCl₃, 27 °C, 600 MHz.

Here, we evaluated whether these novel molecular capsules are capable of including some guest molecules. NMR spectroscopic studies have provided clear evidence that **6** can form complex with 4,4'-trimethylenedipyridine (**4**). The ¹H NMR spectrum of **6** in the presence of **4** is shown in Figure 4. The proton signals for free **6** and **6**·**4** complex




Figure 4. Partial ¹H NMR spectra of (a) **4** (0.8 mM), (b) [**6**]:[**4**] = 2:1 (0.8 mM/0.4 mM), and (c) [**6**]:[**4**] = 1:1 (0.8 mM/0.8 mM): CDCl₃, 27 °C, 600 MHz. Open and filled circles denote the signals for free **4** and **6** and those for **6**·**4** complex, respectively.

appear separately. The peak separation implies that the complexation–decomplexation exchange rate is slower than the ¹H NMR time-scale. The stoichiometry of **6**·**4** complex was estimated to be 1:1 from the peak intensities. These results consistently support the view that **4** resides in the cavity of **6**.²¹ Large upfield shift is observed for the α- and β-protons in the pyridyl groups of **4** (1.91 and 4.69 ppm, respectively). These changes are ascribed to the strong shielding effect of the porphyrin π-systems on the α- and β-protons of the included **4**.

Figure 5 shows the influence of added **4** on the absorption spectral change in **6** (25 °C, CHCl₃). It is seen from Figure




Figure 5. Absorption spectral change in **6** (2.0×10^{-6} M): [**4**] = 0– 5.0×10^{-6} M. Inset: ΔA_{436} vs [**4**] plot in CHCl₃ at 25 °C.

5 that the λ_{\max} for the Soret band (431 nm) shifts to longer wavelength (436 nm) with a tight isosbestic point (433 nm in the Soret band region). The result supports the view that two pyridine units in **4** simultaneously coordinate to Zn(II) in **6**. From a plot of ΔA_{436} vs $[\mathbf{4}]$ (inserted in Figure 5), one can obtain $K = 2.6 \times 10^6 \text{ M}^{-1}$ for the formation of the 1:1 complex from **4** and **6** (the formation of **6·4** complex is 65% at $[\mathbf{4}] = [\mathbf{6}] = 2.0 \times 10^{-6} \text{ M}$). Since the K for the formation of the 1:1 complex from pyridine and ZnTPP (TPP = tetraphenylporphyrin) (estimated under the similar measurement conditions) is $1.1 \times 10^3 \text{ M}^{-1}$, one can regard that the two-point simultaneous binding dramatically enhances the K value. Moreover, the formation of **6·4** complex was also supported by coldspray ionization mass spectrometry. When a CH_2Cl_2 solution containing **4** and **6** in a 1:1 ratio was subjected to the CIS-MS measurement, strong peaks appeared at 1000, 1223, and 1574, which are assignable to $[\mathbf{6}\cdot\mathbf{4} - 6 \text{ CF}_3\text{SO}_3^-]^{6+}$, $[\mathbf{6}\cdot\mathbf{4} - 5 \text{ CF}_3\text{SO}_3^-]^{5+}$, and $[\mathbf{6}\cdot\mathbf{4} - 4 \text{ CF}_3\text{SO}_3^-]^{4+}$, respectively.

(20) Sakamoto, S.; Fujita, M.; Kim, K.; Yamaguchi, K. *Tetrahedron* **2000**, *56*, 955–964.

(21) Although **4** is capable of coordination to Pd(II) to decomposed the capsular structure, it favorably interacts with porphyrin–Zn(II) for the steric suitability.

In conclusion, two porphyrin-based building blocks (**1** or **2**) intermolecularly bind four *cis*-Pd(II) complexes, resulting in a novel molecular capsule according to a self-assembled manner. Owing to rigid acetylene spacers between *meso*-phenyl moieties and pyridyl moieties in **2**, molecular capsule **6** can hold an unusually large cavity enough to bind large bipyridine guest **4**. These results show that porphyrins can act as powerful building blocks for constructing molecular capsules in a self-assembled manner. The further applications of these systems to porphyrin-mediated molecular recognition, redox reactions, photochemical reactions, etc. are currently under investigation in these laboratories.

Acknowledgment. The present work is supported by the Sumitomo Foundation and by a Grant-in-Aid for COE Research “Design and Control of Advanced Molecular Assembly Systems” from the Ministry of Education, Science and Culture, Japan (No. 08CE2005).

Supporting Information Available: Synthetic scheme and assignment of the ^1H NMR spectra of **1**, **2**, **5**, **6**, and **6·4** complex. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL006620M