

Enantiodivergent Synthesis of Either Enantiomer of ABCDE-Ring Analogue of Antitumor Antibiotic Fredericamycin A via Intramolecular [4 + 2] Cycloaddition Approach


Shuji Akai, Toshiaki Tsujino, Nobuhisa Fukuda, Kiyosei Iio, Yoshifumi Takeda, Ken-ichi Kawaguchi, Tadaatsu Naka, Kazuhiro Higuchi, and Yasuyuki Kita*[†]

Graduate School of Pharmaceutical Sciences, Osaka University, 1-6, Yamadaoka, Suita, Osaka 565-0871, Japan

kita@phs.osaka-u.ac.jp

Received September 5, 2001

ABSTRACT



An intramolecular enantiodivergent synthesis of both enantiomers of the ABCDE-ring analogue **22** of fredericamycin A is reported. Key steps involve an intramolecular [4 + 2] cycloaddition of **17** and an aromatic Pummerer-type reaction of **19**. A lipase-catalyzed enantioselective desymmetrization of prochiral diol **2** using 1-ethoxyvinyl 2-furoate **3** led to the pivotal intermediate (*R*)-**4**.

Due to potent antitumor activity against a variety of in vivo tumor models and negative mutagenicity in the Ames test, an antibiotic, fredericamycin A (**1**), has drawn much attention as a leading compound for developing novel types of chemotherapeutic drugs for human cancers.¹ **1** consists of two *peri*-hydroxy tricyclic aromatic moieties, connected through a chiral, spiro, quaternary carbon center, and its chirality arises from the presence of a single methoxy group

at the farthest position of the A-ring. Intensive efforts have been made toward asymmetric total synthesis of **1**, including the total syntheses of racemic **1** by five research groups^{2–6} and the HPLC separation of a racemic intermediate of **1** using a chiral column.⁷ Until recently, no one had succeeded in

[†] Fax: (+81) 6-6879-8229.

(1) Isolation and structure elucidation: (a) Pandey, R. C.; Toussaint, M. W.; Strohane, R. M.; Kalita, C. C.; Aszalos, A. A.; Garretson, A. L.; Wei, T. T.; Byrne, K. M.; Geoghegan, R. F., Jr.; White, R. J. *J. Antibiot.* **1981**, *34*, 1389. (b) Misra, R.; Pandey, R. C.; Silvertown, J. V. *J. Am. Chem. Soc.* **1982**, *104*, 4478. Studies on biological activity: (c) Latham, M. D.; King, C. K.; Gorycki, P.; Macdonald, T. L.; Ross, W. E. *Cancer Chemother. Pharmacol.* **1989**, *24*, 167. (d) Dalal, N. S.; Shi, X. *Biochemistry* **1989**, *28*, 748.

(2) Kelly, T. R.; Bell, S. H.; Ohashi, N.; Armstrong-Chong, R. J. *J. Am. Chem. Soc.* **1988**, *110*, 6471.


(3) Clive, D. L. J.; Tao, Y.; Khodabocus, A.; Wu, Y.-J.; Angoh, A. G.; Bennett, S. M.; Boddy, C. N.; Bordeleau, L.; Kellner, D.; Kleiner, G.; Middleton, D. S.; Nichols, C. J.; Richardson, S. R.; Vernon, P. G. *J. Am. Chem. Soc.* **1994**, *116*, 11275.

(4) Rama Rao, A. V.; Singh, A. K.; Rao, B. V.; Reddy, K. M. *Heterocycles* **1994**, *37*, 1893.


(5) Saint-Jalmes, L.; Lila, C.; Xu, J. Z.; Moreau, L.; Pfeiffer, B.; Eck, G.; Pelsez, L.; Rolando, C.; Julia, M. *Bull. Soc. Chim. Fr.* **1993**, *130*, 447.

(6) Wendt, J. A.; Gauvreau, P. J.; Bach, R. D. *J. Am. Chem. Soc.* **1994**, *116*, 9921.

the asymmetric total synthesis of either **1** or its analogue⁸ because of the lack of an efficient distinction of the enantioface of the highly symmetrical AB-plane.⁹



We have planned two different asymmetric approaches for **1** through an optically pure quaternary carbon intermediate with a definite stereochemistry at an early stage and the completion of the total synthesis while retaining its chiral integrity (Scheme 1). Accordingly, we have quite recently



succeeded in the asymmetric total synthesis of natural **1** by the intermolecular cycloaddition approach, and the absolute stereochemistry of the natural product was determined for the first time.¹⁰ Simultaneously, we have also achieved an intramolecular total synthesis of racemic **1**,^{10c,11} which is believed to be useful for the asymmetric synthesis of optically

(7) Boger, D. L.; Hüter, O.; Mbiya, K.; Zhang, M. *J. Am. Chem. Soc.* **1995**, *117*, 11839.

(8) Studies in this field are well summarized in ref 7. For recent synthetic studies on the racemates, see: Baskaran, S.; Nagy, E.; Braun, M. *Liebigs Ann./Recl.* **1997**, 311. Clive, D. L. J.; Kong, X.; Paul, C. C. *Tetrahedron* **1996**, *52*, 6085. Evans, P. A.; Brandt, T. A. *Tetrahedron Lett.* **1996**, *37*, 1367. Perumal, P. T.; Venugopal, M.; Velusamy, T. P. *Indian J. Chem., Sect. B* **1996**, *35B*, 242.

(9) Quite a few methods applicable to asymmetric synthesis of **1** have been developed, see: (a) Toyota, M.; Terashima, S. *Tetrahedron Lett.* **1989**, *30*, 829. (b) Trypke, W.; Steigel, A.; Braun, M. *Synlett* **1992**, 827.


(10) (a) Kita, Y.; Higuchi, K.; Yoshida, Y.; Iio, K.; Kitagaki, S.; Akai, S.; Fujioka, H. *Angew. Chem., Int. Ed.* **1999**, *38*, 683. (b) Kita, Y.; Higuchi, K.; Yoshida, Y.; Iio, K.; Kitagaki, S.; Ueda, K.; Akai, S.; Fujioka, H. *J. Am. Chem. Soc.* **2001**, *123*, 3214. (c) For a review, see: Kita, Y.; Akai, S.; Fujioka, H. *J. Synth. Org. Chem. Jpn.* **1998**, *56*, 963.

(11) Kita, Y.; Iio, K.; Kawaguchi, K.; Fukuda, N.; Takeda, Y.; Ueno, H.; Okunaka, R.; Higuchi, K.; Tsujino, T.; Fujioka, H.; Akai, S. *Chem. Eur. J.* **2000**, *6*, 3897.


active **1**. In this Letter we report an application of this intramolecular cycloaddition approach involving the intermediates **I–III** to the asymmetric synthesis of the optically active ABCDE-ring analogue **22**. Either enantiomer (*R*)- or (*S*)-**22** was obtained from a single key intermediate, (*R*)-**4**, with perfect retention of its chiral integrity.

Preparation of both enantiomers of the optically active keto aldehyde, (*R*)- and (*S*)-**10**, corresponding to **II**, was the first objective in this study. Creation of the chiral quaternary carbon center was performed by the lipase-catalyzed desymmetrization of prochiral diol **2** using 1-ethoxyvinyl 2-furoate **3**.¹² The desired (*R*)-**4** (21% yield, 96% ee) was obtained along with the diester **5** (78% yield), and **5** was recycled to **2** almost quantitatively. Silylation of (*R*)-**4** followed by saponification afforded the monosilyl ether (*S*)-**6** (95% yield, 96% ee). Stepwise oxidation of (*S*)-**6** by the Dess–Martin periodinane and then by NaClO₂ afforded the carboxylic acid (*R*)-**7** (88% yield), which was treated with BF₃·Et₂O to give (*R*)-**8** (99% yield, 96% ee).¹³ Reaction of (*R*)-**8** with MeLi at 0 °C followed by quenching with aqueous NH₄Cl solution at –78 °C gave the keto-carbinol (*R*)-**9** (80% yield, 96% ee).¹³ The Dess–Martin oxidation of (*R*)-**9** afforded (*R*)-**10** (96% yield). A similar two-step oxidation of (*R*)-**4** gave the carboxylic acid (*S*)-**11** (92% yield), which was then converted to (*S*)-**10** (49% yield, two steps) (Scheme 2).

To prepare the key intermediate (*S*)-**16**, the installation of the acetylene and the A-ring moieties to (*R*)-**10** without loss of its chiral integrity was troublesome. When the reaction




^a (a) **3**, *Candida rugosa* lipase (Meito MY), iPr₂O, 30 °C; (b) K₂CO₃, MeOH; (c) 1. TBSCl, pyridine, DMF; 2. K₂CO₃, MeOH; (d) 1. Dess–Martin periodinane, MeCN, 2. NaClO₂, NaH₂PO₄, 2-methyl-2-butene, *t*-BuOH, H₂O; (e) BF₃·Et₂O, CH₂Cl₂; (f) MeLi, HMPA, THF; (g) Dess–Martin periodinane, MeCN.

Scheme 3^a

^a (a) LiC≡CSPH, HMPA–THF, –78 °C; (b) **13**, DMAP, CH₂Cl₂; (c) LiN(TMS)₂, THF; (d) Moffatt oxidation; (e) 1. Co₂(CO)₈, CH₂Cl₂; 2. BCl₃, CH₂Cl₂; (f) 1. Me₂SiCl₂, Et₃N, chloranil, toluene, 100 °C; 2. (*t*-Bu)₂Si(OTf)₂, Et₃N, DMF; (g) *m*-CPBA, CH₂Cl₂; (h) **20**, catalytic *p*-TsOH, toluene; (i) 1. Bu₄NF, THF–H₂O; 2. BBr₃, CH₂Cl₂; 3. 80% aqueous CF₃CO₂H.

of (*R*)-**10** (96% ee) with lithium phenylthioacetylde was carried out at room temperature, unexpected racemization of the quaternary carbon center at the C-1 position was brought about.¹⁴ This problem was overcome by running the reaction at –78 °C and quenching at the same temperature to give (*1R*)-**12** (53% yield, 96% ee) as a single diastereomer. Acylation of (*1R*)-**12** with **13** gave (*1R*)-**14** (96% yield), which was treated with LiN(TMS)₂ (3 equiv) in THF to induce the migration of the A-ring aryl moiety to the methyl ketone terminus giving (*1R*)-**15** (81% yield). Moffatt oxidation of (*1R*)-**15** afforded the trione (*S*)-**16** (79% yield).

The following manipulation of (*S*)-**16** was performed similarly to the total synthesis of the racemate¹¹ without any problem. Thus, treatment of (*S*)-**16** with Co₂(CO)₈ followed by debenzoylation provided the cobalt complex (*S*)-**17** (50%

Scheme 4^a

^a (a) 1. LiC≡CSPH, HMPA–THF, –78 °C; 2. **13**, DMAP, CH₂Cl₂; 3. LiN(TMS)₂, THF; (b) 1. Moffatt oxidation; 2. Co₂(CO)₈, CH₂Cl₂; 3. BCl₃, CH₂Cl₂; (c) 1. Me₂SiCl₂, Et₃N, chloranil, toluene, 100 °C; 2. (*t*-Bu)₂Si(OTf)₂, Et₃N, DMF; (d) 1. *m*-CPBA, CH₂Cl₂; 2. **20**, catalytic *p*-TsOH, toluene; 3. Bu₄NF, THF–H₂O; 4. BBr₃, CH₂Cl₂; 5. 80% aqueous CF₃CO₂H.

yield), which was subjected to intramolecular [4 + 2] cycloaddition under the standard reaction conditions to afford the pentacyclic product (*S*)-**18** (47% yield). The optical purity of (*S*)-**18** was determined to be 96% ee by chiral HPLC analysis, and the perfect retention of the chiral integrity of (*R*)-**4** was confirmed. Oxidation of (*S*)-**18** to (*S*)-**19** followed by aromatic Pummerer-type reaction using 1-ethoxyvinyl chloroacetate **20** in refluxing toluene¹⁵ introduced a hydroxyl group to the B-ring. The crude product (*R*)-**21** was subjected to stepwise deprotection to afford the optically active ABCDE-ring analogue (*S*)-**22** having the same absolute stereochemistry as natural **1** (Scheme 3).¹⁶

The same transformation was applied to (*S*)-**10** (96% ee) to provide the unnatural enantiomer (*R*)-**22** [96% ee based on the optical purity of (*R*)-**18**] (Scheme 4).¹⁶ A symmetrical pair in the CD spectra of (*S*)- and (*R*)-**22** was obtained

(12) Akai, S.; Naka, T.; Fujita, T.; Takebe Y.; Kita, Y. *Chem. Commun.* **2000**, 1461.

(13) Special care was needed for the desilylation of (*R*)-**7** and the reaction of (*R*)-**8** with MeLi, because a partial racemization of them was often observed under basic conditions. For instance, the reaction of (*R*)-**8** with MeLi at 0 °C followed by quenching at the same temperature gave (*R*)-**9** in ≤92% ee.

(14) The racemization may arise from a retro-aldol-aldol process of the intermediate β-oxidoketone corresponding to **12**. A similar reaction was observed during a study of the asymmetric total synthesis of **1**.¹⁰ A detailed investigation about this racemization is now in progress and will be discussed in the near future.

(15) (a) Kita, Y.; Takeda, Y.; Matsugi, M.; Iio, K.; Gotanda, K.; Murata K.; Akai, S. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1529. See also, (b) Kita, Y.; Takeda, Y.; Iio, K.; Yokogawa, K.; Takahashi, K.; Akai, S. *Tetrahedron Lett.* **1996**, *37*, 7545. (c) Akai, S.; Takeda, Y.; Iio, K.; Takahashi, K.; Fukuda, N.; Kita, Y. *J. Org. Chem.* **1997**, *62*, 5526.

(16) Due to the poor solubility of **22** in various organic solvents and its deep color, determination of its optical purity was difficult. All new compounds were fully characterized by spectroscopic means and combustion analysis.




Figure 1. CD spectra of (*S*)- and (*R*)-**22**.

(Figure 1). The spectroscopic data (IR, ^1H NMR, ^{13}C NMR, HRMS) of (*S*)- and (*R*)-**22** showed good agreement with its structure.¹⁷

Acknowledgment. We are grateful to Akiko Okajima and Seiko Morishita for their technical assistance. This work was financially supported by Grants-in-Aid for Scientific Research [Priority Areas (A) “Exploitation of Multi-Element Cyclic Molecules” and No. 13672214] from the Ministry of Education, Culture, Sports, Science, and Technology, Japan, and the Takeda Science Foundation, Japan. Meito Sangyo Co., Ltd., Japan, is thanked for generous gift of lipase MY.

Supporting Information Available: ^1H and ^{13}C NMR data of (*S*)-**22** and racemic **23**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL016696Y

(17) The similarity of the ^1H and ^{13}C NMR data of (*S*)-**22** with those of racemic **23** [Boger, D. L.; Jacobson, I. C. *J. Org. Chem.* **1991**, *56*, 2115] also supports its structure (In detail, see: Supporting Information).

