

## A Forbidden Rearrangement


Martin Leivers, Iris Tam, Kevin Groves, David Leung, Yuli Xie, and  
Ronald Breslow\*

Department of Chemistry, Columbia University, New York, New York 10027

rb33@columbia.edu


Received June 13, 2003

## ABSTRACT




A barrelene derivative fragments to afford benzene and trappable 1,2,3-tricyanocyclopropene. The barrelene anion fragments more easily to liberate benzene and the 1,2,3-tricyanocyclopropenyl anion, which is not trappable or stable in solution. However, the major thermal product from the barrelene anion is a rearranged allyl anion that is formed by disrotatory cleavage of the cyclopropyl ring, a formally Woodward–Hoffmann-forbidden process. Several proposals are offered to rationalize this forbidden rearrangement.

For quite a while, we have been trying to synthesize the tricyanocyclopropenyl anion **1**, which should be strongly stabilized and a ground-state triplet. Our earliest approaches<sup>1</sup> involved the treatment of 1,2,3-tricyanochlorocyclopropane with base to first eliminate HCl and then deprotonate the resulting cyclopropene **2**, but in all cases we obtained only polymeric materials. Apparently, cyclopropene **2** undergoes anionic polymerization, initiated either by the base used or by the anion **1** if indeed that is formed. An attempt to generate **1** by reduction of 1,2,3-tricyanotrichlorocyclopropane was also unsuccessful.<sup>1</sup> Thus, we conceived a plan in which the double bond of **1** would be introduced *after* the negative charge was in place.



We synthesized compound **3**, a formal Diels–Alder adduct of benzene and tricyanocyclopropene, using the procedure described below. We hoped that its anion **4** would fragment to form benzene and **1** and that the alternative ring opening to form anion **5** would not occur since it is forbidden by the Woodward–Hoffmann general treatment,<sup>2</sup> which requires that the interconversion of a cyclopropyl anion with an allyl

anion involve conrotation in the breaking or making of the cyclopropyl single bond. Such conrotation is not possible with anion **4**, since the product would not be **5** but instead a hopelessly strained stereoisomer. Even so, this forbidden rearrangement of **4** into **5** was the principal process.



Fumaronitrile was converted into **6**<sup>3</sup> by addition of methanethiol. Chlorination with *N*-chlorosuccinimide led to **7**,<sup>3</sup> and with base this afforded **8**.<sup>3,4</sup> Then reaction with (cyanomethyl)dimethylsulfonium triflate and NaH afforded **9**,<sup>3,4</sup> converted into **10**<sup>3,4</sup> with MeI and AgBF<sub>4</sub>. (With base, compound **10** affords cyclopropene **2**, but under all conditions tried this again was converted into polymer, not to a stable solution of anion **1**.) The Fe(CO)<sub>3</sub> adduct **11**<sup>3</sup> of 1,3-cyclohexadiene was converted into cation **12**<sup>3</sup> by hydride abstraction with triphenylmethyl fluoborate, and capture of


(1) Breslow, R.; Cortes, D.; Jaun, B.; Mitchell, R. *Tetrahedron Lett.* **1982**, 23, 795.

(2) Woodward, R. B.; Hoffmann, R. *The Conservation of Orbital Symmetry*; Academic Press: New York, 1970.

(3) Experimental details and characterization are found in the Supporting Information.


(4) A major and minor stereoisomer were formed and characterized.

the cation with  $\text{NH}_2\text{Boc}$  afforded **13**.<sup>3</sup> Ceric ammonium nitrate converted this into diene **14**.<sup>3</sup> Reaction of **10** with **14** in the presence of *N,N*-diisopropylethylamine (generating **2**) afforded adduct **15**.<sup>3,4</sup> After removal of the Boc group, the amino group was dimethylated with formaldehyde and sodium triacetoxyborohydride to form **16**,<sup>3</sup> and quaternization with MeI afforded salt **17**.<sup>3</sup>



Treatment of **17** with KOtBu afforded **3**. However, during this process the allyl anion **5** was also formed by rearrangement of the cyclopropyl anion **4**. Thus, we have examined both the use of **3** to form cyclopropene derivatives by reverse Diels–Alder reactions and also the rearrangement of anion **4** into allyl anion **5**, a process involving the forbidden disrotatory cleavage of the cyclopropyl ring bond.

Compound **3** has the stereochemistry shown (NOE between the cyclopropyl proton and the upfield vinyl hydrogens of **3**) and allyl anion **5** was characterized by NMR (only two proton signals at  $\delta$  5.79 and 2.82, in a 2:1 ratio). Acid quenching of anion **5** afforded triene **18**.<sup>3</sup>




On heating above 170 °C in DMSO-*d*<sub>6</sub> in a sealed tube under argon, compound **3** undergoes a reverse Diels–Alder reaction to form benzene and 1,2,3-tricyanocyclopropene **2**.

The latter can be trapped to furnish the known adduct **20**<sup>1</sup> (in equal amount to the benzene) if diphenylisobenzofuran **19** is present in solution during the thermolysis, but not if it is added later. Thus, even under these thermal nonbasic conditions cyclopropene **2** is not stable in solution, apparently again forming a polymer. When KOtBu is added to a solution of **3** in DMSO-*d*<sub>6</sub> under argon at 165 °C there is the immediate formation of some benzene, suggesting that the anion **3** cracks to liberate the cyclopropenyl anion **1**, but this could not be trapped with **19**, either present in the solution or added later with mild acid. In addition, the rearranged allylic anion **5** is formed in this process, in about a 10:1 ratio to the benzene formed. Thus, the principal process for anion **4** is rearrangement, not fragmentation.

Anion **4** can actually be formed at room temperature by treating **3** with NaH in DMSO-*d*<sub>6</sub>. The cyclopropyl H signal of **3** at  $\delta$  3.24 is lost in the NMR, and the vinyl proton signals of **3** at  $\delta$  6.95 and 6.21 and the bridgehead proton signals at  $\delta$  4.35 move upfield to  $\delta$  6.68, 5.95, and 3.74, respectively, in anion **4**. The anion is moderately stable and can be reprotonated to form **3** (but not its epimer) or alkylated with methyl iodide to form the two isomers of **21**.<sup>3,4</sup> (one with NOE of the methyl to vinyl hydrogens). However, on gentle warming or even on standing for 3–4 h at room temperature, anion **4** undergoes rearrangement into allylic anion **5** (15% also fragments to form benzene).

This rearrangement is a formally “forbidden” reaction: the disrotatory cleavage of the cyclopropane sigma bond is forbidden for the anion,<sup>5</sup> allowed for the cation. Since it occurs fairly readily, an explanation is needed. It should be mentioned that a related reaction has been seen previously (such violations of the Woodward–Hoffmann rules are apparently not widely known), the rearrangement of anion **22** into allylic anion **23**.<sup>5</sup> Thus the extra  $\pi$  electrons in our anion **4** are not part of the explanation.




We propose that several factors may be overcoming the electronic problem in the disrotatory ring opening of anion **4**. One is that the unshared carbon electrons in cyanocyclopropyl anions are in hybrid orbitals, not in p orbitals. This is revealed in stereochemical studies of such anions previously, in which deuterium exchanges involving such anions are faster than are inversion processes.<sup>6,7</sup> As a result, the disrotatory opening of the cyclopropane initially pushes the  $\sigma$ -bond electrons away from the carbanion electrons (Figure 1), and indeed the  $\sigma$ -bond electrons begin to overlap with the empty hybrid orbital of the carbanion, not the filled orbital. That is, the system begins to resemble an allylic cation, not just an anion, although later rehybridization of the three carbons to  $\text{sp}^2$  indeed leads to the product allylic anion. In

(5) Boche, G. *Top. Curr. Chem.* **1988**, *146*, 1–156, see p 22.

(6) Walborsky, H. M.; Youssef, A. A.; Motes, J. M. *J. Am. Chem. Soc.* **1962**, *84*, 2465.

(7) Breslow, R.; Douek, M. *J. Am. Chem. Soc.* **1968**, *90*, 2698.



**Figure 1.** As the cyclopropyl single bond breaks by a disrotatory motion, the electrons are pushed away from the carbanion electrons and overlap with the empty hybrid orbital pointing down, not the filled orbital pointing up.

a sense, the factors involved are a little like those that lead to backside displacement reactions in which electron pairs interact with  $\sigma^*$ , avoiding the  $\sigma$  electrons of the leaving group.

A second factor might be special for our system, with its three cyano groups. They are expected to strongly stabilize the antiaromatic cyclopropenyl anion system by withdrawing  $\pi$  electrons, and the antiaromaticity of the forbidden transition state in disrotatory rearrangement of a cyclopropyl anion into an allylic anion might also be diminished by such electron

withdrawal. Finally, it is possible that the rearrangements might be in a sense stepwise, with stretching of the single bond preceding the other geometric changes.

Whichever factors are involved, and we favor the first possibility mentioned above, it is clear that a formally forbidden reaction is actually occurring in this case. It competes with the observed fragmentation of **4** to form benzene and the tricyanocyclopropenyl anion **1** in solution, a fragmentation that is much faster with anion **4** than with the neutral **3**. Thus, the fragmentation is indeed being driven by the delocalization of charge in product **1**. However, we have no evidence for the formation of a *stable* solution of **1**, which remains a challenging target.

**Acknowledgment.** This work was partially supported by the NSF.

**Supporting Information Available:** Experimental details and spectra of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL034997X