


Synthesis of Homoallylic Chiral Tertiary Alcohols via Chelation-Controlled Diastereoselective Nucleophilic Addition on α -Alkoxyketones: Application for the Synthesis of the C₁–C₁₁ Subunit of 8-*epi*-Fostriecin

P. Veeraraghavan Ramachandran,* Haipeng Liu, M. Venkat Ram Reddy, and Herbert C. Brown

Herbert C. Brown Center for Borane Research, 560 Oval Drive,
Department of Chemistry, Purdue University, West Lafayette, Indiana 47907-2084
chandran@purdue.edu

Received August 11, 2003

ABSTRACT



Chiral β -*syn*-alkoxyhomoallylic alcohols derived from alkoxyallylboration of aldehydes upon oxidation provided the corresponding chiral ketones. Chelation-controlled nucleophilic addition to these ketones occurred in a highly stereoselective manner to afford *anti*-homoallylic tertiary alcohols. This methodology has been applied for the synthesis of the C₁–C₁₁ subunit of C₈-*epi*-fostriecin.

Homoallylic alcohols are extremely versatile synthetic intermediates that have been extensively utilized for the stereoselective synthesis of complex natural products.¹ Due to their importance, there have been several methods reported in the literature for the stereoselective synthesis of homoallylic alcohols,² and allylboration³ is a widely used procedure. For the past two decades, we have been developing various α -pinene-based reagents for asymmetric “allyl”

boration, and these reagents have found wide acceptance among the synthetic organic community.⁴ One such reagent is *B*- γ -alkoxyallyldiisopinocampheylborane for the stereoselective alkoxyallylboration of aldehydes.⁵ Although this reagent provides very high levels of ee and de for aldehydes, ketones fail to provide similar results, probably due to less favorable stereoelectronic factors.

Stereoselective synthesis of homoallylic tertiary alcohols is very important since they can be used as building blocks for the synthesis of complex natural products. There has been no general procedure for the preparation of homoallylic tertiary alcohols in high de and ee. We envisaged that oxidation of β -alkoxy homoallylic alcohol to the correspond-

(1) (a) Yamamoto, Y. *Acc. Chem. Res.* **1987**, *20*, 243. (b) Roush, W. R. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: Oxford; Vol. 2.

(2) Yamamoto, Y.; Asao, N. *Chem. Rev.* **1993**, *93*, 2207.


(3) (a) Hoffmann, R. W. *Pure Appl. Chem.* **1988**, *60*, 123. (b) Roush, W. R. ACS Symposium Series 386; American Chemical Society: Washington, DC, 1989; p 242. (c) Roush, W. R. In *Methods of Organic Chemistry (Houben-Weyl)*; Georg Thieme Verlag: Stuttgart, 1995; Vol. E 21, p 1410. (d) Brown, H. C.; Ramachandran, P. V. *J. Organomet. Chem.* **1995**, *500*, 1.

(4) Ramachandran, P. V. *Aldrichim. Acta* **2002**, *35*, 23.

(5) Brown, H. C.; Jadhav, P. K.; Bhat, K. S. *J. Am. Chem. Soc.* **1988**, *110*, 1535.

ing ketone, followed by nucleophilic addition, would furnish homoallylic tertiary alcohols. Chelation-controlled diastereoselective nucleophilic additions on α -alkoxyketones⁶ are well-known. On this basis, we undertook a project involving diastereoselective nucleophilic additions on α -alkoxyketones derived from *syn*-alkoxy homoallylic alcohols.

We started with *syn*-monoalkoxy homoallylic alcohols **4a** and **4b** derived from reaction of *B*- γ -methoxyethoxymethoxyallyldiisopinocampheylborane⁷ **3** with acetaldehyde and benzaldehyde, respectively. The alcohols **4a** and **4b** were oxidized to the corresponding ketones **5a** and **5b** using Dess–Martin periodinane (DMP).⁸ We chose various nucleophiles as representative examples for the present study. Reduction of ketones **5a** and **5b** with Zn(BH₄)₂ at -78 °C afforded *anti*-homoallylic 2°-alcohols **6a** and **6e** in 50 and 95% de, respectively. Addition of organometallic reagents took place very smoothly, and the product *anti*-homoallylic tertiary alcohols were obtained in very high de (Scheme 1



^a Conditions: (a) *sec*-BuLi, THF, -78 °C, 0.5 h. (b) (+)-Ipc₂BOMe, -78 °C, 1 h. (c) BF₃·Et₂O, -78 °C, 5 min. (d) RCHO, -78 °C, 10 h. (e) NaOH/H₂O₂, rt, 6 h. (f) DMP, CH₂Cl₂. (g) Nu⁻.


and Table 1). In all of these cases, metal coordination with carbonyl oxygen and alkoxy oxygen atom leads to a five-membered transition state followed by nucleophilic addition from the opposite face providing the *anti* products.

To demonstrate the utility of this procedure, we converted the *anti*-alcohols **6e** and **6f** into *anti*-diols **7e** and **7f**,

Table 1.

compd	R	Nu	yield	de
6a	Me	Zn(BH ₄) ₂	90	50
6b	Me	EtMgBr	85	>95
6c	Me	(CH ₂ =CH)MgBr	78	>95
6d	Me	CH ₃ -C≡CMgBr	75	85
6e	Ph	Zn(BH ₄) ₂	92	>95
6f	Ph	EtMgBr	88	>95
6g	Ph	(CH ₂ =CH)MgBr	81	>95
6h	Ph	CH ₃ -C≡CMgBr	77	>95

respectively, in good yields by cleaving the MEM group under acidic conditions (Scheme 2).



Application of this methodology was further demonstrated by the synthesis of the C₈ epimer of the C₁–C₁₁ subunit of fostriecin (Figure 1). Fostriecin is a natural product isolated




Figure 1.

from *Streptomyces pulveraceus* exhibiting potent anti-cancer properties against a wide range of cell lines.⁹ Accordingly, there have been several reports of the synthesis of this molecule in the recent past.¹⁰ We have employed a strategy similar to our ongoing program on “allyl”boration–ring-closing metathesis for the synthesis of biologically active natural products.¹¹

- (6) (a) Reetz, M. T. *Angew. Chem., Int. Ed.* **1984**, *23*, 556. (b) Still, W. C.; McDonald, J. H., III. *Tetrahedron Lett.* **1980**, *21*, 1031. (c) Nicolaou, K. C.; Claremon, D. A.; Barnette, W. E. *J. Am. Chem. Soc.* **1980**, *102*, 6611. (d) McGarvey, G. J.; Kimura, M. *J. Org. Chem.* **1982**, *47*, 5422. (e) Kobayashi, Y.; Kumar, G. B.; Kurachi, T.; Acharya, H. P.; Yamazaki, T.; Kitazume, T. *J. Org. Chem.* **2001**, *66*, 2011. (f) Carven, A.; Tapolczay, D. J.; Thomas, E. J.; Whiteland, J. W. F. *Chem. Commun.* **1985**, 145.
- (7) (a) Nicolaou, K. C.; Smith, A. L.; Mizuno, Y. *J. Am. Chem. Soc.* **1993**, *115*, 7612. (b) Ramachandran, P. V. Prabhudas, B.; Pratihari, D.; Chandra, J. S.; Reddy, M. V. R. *Tetrahedron Lett.* **2003**, *44*, 3745.
- (8) (a) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155. (b) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277.
- (9) (a) Tunac, J. B.; Graham, B. D.; Dobson, W. E. *J. Antibiot.* **1983**, *36*, 1595. (b) Stampwala, S. S.; Bunge, R. H.; Hurley, T. R.; Willmer, N. E.; Brankiewicz, A. J.; Steinman, C. E.; Smittka, T. A.; French, J. C. *J. Antibiot.* **1983**, *36*, 1601. (c) Leopold, W. R.; Shillis, J. L.; Nelson, J. M.; Roberts, B. J. *Cancer Res.* **1984**, *44*, 1928.
- (10) (a) Boger, D. L.; Hikota, M.; Lewis, B. M. *J. Org. Chem.* **1997**, *62*, 1748. (b) Boger, D. L.; Ichikawa, S.; Zhong, W. *J. Am. Chem. Soc.* **2001**, *123*, 4161. (c) Chavez, D. E.; Jacobsen, E. N. *Angew. Chem. Int. Ed.* **2001**, *40*, 3667. (d) Reddy, Y. K.; Falck, J. R. *Org. Lett.* **2002**, *4*, 969. (e) Miyashita, K.; Ikejiri, M.; Kawasaki, H.; Maemura, S.; Imanishi, T. *Chem. Commun.* **2002**, 742. (f) Wang, Y.-G.; Kobayashi, Y. *Org. Lett.* **2002**, *4*, 4615. (g) Esumi, T.; Okamoto, N.; Hatakayama, S. *Chem. Commun.* **2002**, 3042. (h) Fujii, K.; Maki, K.; Kanai, M.; Shibasaki, M. *Org. Lett.* **2003**, *5*, 733. (i) Miyashita, K.; Ikejiri, M.; Kawasaki, H.; Maemura, S.; Imanishi, T. *J. Am. Chem. Soc.* **2003**, *125*, 8238.

Monoprotection of *cis*-2-butene-1,4-diol **8**, with TBSCl, followed by PCC oxidation, provided the *trans*-aldehyde **9**.¹² This, upon alkoxyallylboration with (–)-*B*- γ -methoxyethoxymethoxyallyldiisopinocampheylborane **3**, furnished the homoallylic alcohol **10** in >98% de and 94% ee. DMP oxidation provided the α -alkoxyketone, and addition of methylmagnesium bromide took place stereoselectively to provide the anti tertiary alcohol **11** in >90% de. Protection of this alcohol as its TBS ether and subsequent deprotection of the primary TBS group yielded the alcohol **12**. DMP oxidation, followed by allylboration of the resulting aldehyde with *B*-allyldiisopinocampheylborane,¹³ gave the homoallylic alcohol **13** in 87% de. Treatment with acryloyl chloride and ring-closing metathesis of the resulting acrylate with Grubbs' first-generation ruthenium catalyst¹⁴ provided the C₈-epimer of the C₁–C₁₁ subunit of fostriecin **14** (Scheme 3).


In conclusion, we have developed an efficient procedure for the highly stereoselective synthesis of *anti*- β -alkoxy and β -hydroxy homoallylic 3°-alcohols. We have also applied this procedure for the synthesis of the C₈-epimer of the C₁–C₁₁ subunit of fostriecin. Further exploration of this methodology is in progress, and we believe that this protocol for preparing chiral homoallylic 3°-alcohols and diols will find major applications in organic synthesis.

(11) (a) Ramachandran, P. V.; Chandra, J. S.; Reddy, M. V. R. *J. Org. Chem.* **2002**, *67*, 7547. (b) Reddy, M. V. R.; Yucel, A. J.; Ramachandran, P. V. *J. Org. Chem.* **2001**, *66*, 2512. (c) Reddy, M. V. R.; Rearick, J. P.; Hoch, N.; Ramachandran, P. V. *Org. Lett.* **2001**, *3*, 19.

(12) Corey, E. J.; Suggs, J. W.; *Tetrahedron Lett.* **1975**, *16*, 2647.

(13) Brown, H. C.; Jadhav, P. K. *J. Am. Chem. Soc.* **1983**, *105*, 2092.

(14) For recent reviews, see: (a) Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413. (b) Furstner, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 3013. (c) Trnka, T.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, *34*, 18.



^a Conditions: (a) TBSCl, imidazole, 72%. (b) PCC, CH₂Cl₂, 81%. (c) CH₂=CHCH₂OMEM, *sec*-BuLi, (–)-Ipc₂BOMe, BF₃·Et₂O, NaOH, H₂O₂, 69%. (d) DMP, 92%. (e) MeMgBr, –78 °C. (f) TBSOTf, 2,6-lutidine, 79% overall. (g) AcOH, THF, H₂O, 72%. (h) DMP, 88%. (i) (+)-Ipc₂BAlI, NaOH, H₂O₂, 65%. (j) Acrylic acid, DCC, DMAP, 84%. (k) Grubbs first-generation generation catalyst, CH₂Cl₂, 78%.

Acknowledgment. Financial assistance from the Herbert C. Brown Center for Borane Research¹⁵ and the Aldrich Chemical Co. is gratefully acknowledged.

Supporting Information Available: Experimental and spectral data along with ¹H and ¹³C NMR spectra for various compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL035516C

(15) Contribution #29 from Herbert C. Brown Center for Borane Research.