

Lewis Base-Catalyzed Additions of  
Alkynes Using Trialkoxysilylalkynes

Robert B. Lettan II and Karl A. Scheidt\*

Department of Chemistry, Northwestern University, Evanston, Illinois 60208

scheidt@northwestern.edu

Received May 4, 2005


## ABSTRACT



The Lewis base-catalyzed additions of alkynyl nucleophiles to aldehydes, ketones, and imines is described. Mechanistic studies strongly indicate that the use of new triethoxysilylalkynes facilitates access of a reactive hypervalent silicate intermediate. This activated carbon nucleophile subsequently undergoes rapid addition to carbonyl compounds and imines, thus affording the secondary and tertiary propargyl systems in moderate to high yield.

Chemical reactions catalyzed by nucleophilic species (Lewis bases) possess significant potential for new bond-forming strategies. Over the past decade, many advances in Lewis base-catalyzed processes have been realized, including asymmetric acylations,<sup>1</sup> aldol reactions,<sup>2</sup> and the enantioselective synthesis of  $\beta$ -lactams and  $\beta$ -lactones.<sup>3</sup> However, Lewis base-catalyzed reactions have not been explored to the same extent as Lewis acid or transition-metal-promoted processes, and consequently, these organocatalytic nucleophilic manifolds possess considerable promise. Many Lewis base-catalyzed reactions involve the judicious incorporation of a silyl group that can facilitate access to a hypervalent silicate species which ultimately undergoes delivery of a nucleophile.<sup>4</sup> Based on this principle, we reasoned that the mild and general addition of an alkynyl unit to various electrophiles might be accomplished employing Lewis base

catalysts. In this paper, we report that trialkoxysilylalkynes (**1**) are novel alkynyl nucleophiles in the presence of catalytic Lewis bases and undergo mild and efficient addition to aldehydes and ketones (**2**) to cleanly afford propargyl alcohols (**3**, eq 1).



Many highly stereoselective metal-promoted additions of alkynes to aldehydes have been reported, most notably by Carreira and co-workers.<sup>5</sup> However, there are fewer general strategies for the addition of alkynes to both aldehydes and ketones under mild catalytic conditions.<sup>6</sup> Inspired by the silicon-based allyl additions,<sup>7</sup> carbonyl reductions,<sup>8</sup> that are catalyzed by Lewis bases, we envisioned that appropriately

(1) (a) Fu, G. C. *Acc. Chem. Res.* **2000**, *33*, 412–420. (b) Vedejs, E.; Daugulis, O.; MacKay, J. A.; Rozners, E. *Synlett* **2001**, 1499–1505.

(2) (a) Denmark, S. E.; Stavenger, R. A.; Su, X. P.; Wong, K. T.; Nishigaichi, Y. *Pure Appl. Chem.* **1998**, *70*, 1469–1476. (b) Denmark, S. E.; Stavenger, R. A. *Acc. Chem. Res.* **2000**, *33*, 432–440.

(3) (a) France, S.; Weatherwax, A.; Taggi, A. E.; Lectka, T. *Acc. Chem. Res.* **2004**, *37*, 4245–4255. (b) Wynberg, H.; Staring, E. G. J. *J. Am. Chem. Soc.* **1982**, *104*, 166–168 and references therein. (c) Orr, R. K.; Calter, M. A. *Tetrahedron* **2003**, *59*, 3545–3565.

(4) (a) Chuit, C.; Corriu, R. J. P.; Reye, C.; Young, J. C. *Chem. Rev.* **1993**, *93*, 1371–1448. (b) Holmes, R. R. *Chem. Rev.* **1996**, *96*, 927–950. (c) Brook, M. A. *Silicon in Organic, Organometallic, and Polymer Chemistry*; Wiley-Interscience: New York, 1999.


(5) (a) Frantz, D. E.; Fassler, R.; Carreira, E. M. *J. Am. Chem. Soc.* **2000**, *122*, 1806–1807. (b) Anand, N. K.; Carreira, E. M. *J. Am. Chem. Soc.* **2001**, *123*, 9687–9688. (c) Pu, L. *Tetrahedron* **2003**, *59*, 9873–9886 and references therein. (d) Li, X. S.; Lu, G.; Kwok, W. H.; Chan, A. S. C. *J. Am. Chem. Soc.* **2002**, *124*, 12636–12637.

(6) (a) Cozzi, P. G. *Angew. Chem., Int. Ed.* **2003**, *42*, 2895–2898. For specific asymmetric alkynyl additions to acetophenone derivatives, see: (b) Lu, G.; Li, X. S.; Jia, X.; Chan, W. L.; Chan, A. S. C. *Angew. Chem., Int. Ed.* **2003**, *42*, 5057–5058. (c) Saito, B.; Katsuki, T. *Synlett* **2004**, 1557–1560.

functionalized alkynes might productively interact with an electron-rich catalyst to generate a reactive nucleophilic species.<sup>9,10</sup>

To test this hypothesis, electron-deficient trialkoxysilyl-alkynes (such as **5a**) were easily accessed from the corresponding terminal alkynes<sup>11</sup> and subjected to various nucleophilic species in the presence of an aldehyde (**4**, Table 1, eq 2). The alkoxy substituents were placed on the silicon

**Table 1.** Lewis Base-Catalyzed Additions of Alkynes<sup>a</sup>



entry	Lewis base	equiv	time (h) ((T (°C)) <sup>b</sup>	yield <sup>c</sup> (%)
1	<i>n</i> -Bu <sub>4</sub> N·F <sub>2</sub> SiPh <sub>3</sub>	1.0	16 (23)	50
2	LiOMe	1.0	24 (23)	NR <sup>d</sup>
3	NaOMe	1.0	19 (23)	50
4	KOEt	1.0	1 (23)	57
5	KOEt	0.1	2 (0)	84
6	KO- <i>t</i> -Bu	0.2	2 (0)	75
7	(±)-KOCH(CH <sub>3</sub> )Ph	0.2	2 (0)	75

<sup>a</sup> All reactions were performed under inert atmosphere at 0.2 M.  
<sup>b</sup> Reactions halted after 100% conversion. <sup>c</sup> Isolated yields after purification.  
<sup>d</sup> No reaction.

to induce significant hypervalency in the presence of a Lewis base catalyst while maintaining ease of synthetic accessibility.<sup>12</sup> Fluoride sources were initially surveyed as potential catalysts, but stoichiometric quantities of various salts resulted in limited success (entry 1). Prompted by the strong Si–O bond strength, we examined simple alkoxides with various counterions. Interestingly, while LiOMe afforded no product (entry 2), the use of alkoxides with more electro-positive counterions (Na and K) cleanly provided the desired propargyl alcohol **6**, with KOEt affording a significant rate enhancement over NaOMe (entries 3 and 4). During these early investigations, we noticed product decomposition as a

(7) (a) Kobayashi, S.; Nishio, K. *Tetrahedron Lett.* **1993**, *34*, 3453–3456. (b) Nakajima, M.; Saito, M.; Shiro, M.; Hashimoto, S. *J. Am. Chem. Soc.* **1998**, *120*, 6419–6420. (c) Iseki, K.; Mizuno, S.; Kuroki, Y.; Kobayashi, Y. *Tetrahedron* **1999**, *55*, 977–988. (d) Denmark, S. E.; Fu, J. P. *J. Am. Chem. Soc.* **2001**, *123*, 9488–9489 and references therein.

(8) (a) Kobayashi, S.; Yasuda, M.; Hachiya, I. *Chem. Lett.* **1996**, 407–408. (b) Schiffrers, R.; Kagan, H. B. *Synlett* **1997**, 1175–1178. (c) LaRonde, F. J.; Brook, M. A. *Tetrahedron Lett.* **1999**, *40*, 3507–3510. (d) Iwasaki, F.; Onomura, O.; Mishima, K.; Maki, T.; Matsumura, Y. *Tetrahedron Lett.* **1999**, *40*, 7507–7511. (e) Nishikori, H.; Yoshihara, R.; Hosomi, A. *Synlett* **2003**, 561–563.

(9) For early contributions in this area, see: (a) Nakamura, E.; Kuwajima, I. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 498–499. (b) Kuwajima, I.; Nakamura, E.; Hasimoto, K. *Tetrahedron* **1983**, *39*, 975–982.

(10) With trialkylsilylalkynes, see: (a) Busch-Petersen, J.; Bo, Y.; Corey, E. J. *Tetrahedron Lett.* **1999**, *40*, 2065–2068. (b) Baldwin, J. E.; Pritchard, G. J.; Rathmell, R. E. *J. Chem. Soc., Perkin Trans. 1* **2001**, 2906–2908. (c) Kraus, G. A.; Bae, J. *Tetrahedron Lett.* **2003**, *44*, 5505–5506.

(11) See the Supporting Information for details.

(12) The use of trialkylsilylalkynes with KOEt afforded minimal amounts of alkynyl addition products with aldehydes and was not pursued.

function of increasing reaction time and Lewis base concentration. Gratifyingly, catalytic reaction conditions with only 10 mol % of KOEt at 0 °C generated alcohol **6** in good yield (entry 5). In addition, sterically hindered tertiary and secondary alkoxides can also be employed as catalysts (entries 6 and 7).<sup>13</sup>

With catalytic conditions identified for this process, the structure of the alkyne nucleophile was varied (Table 2, eq

**Table 2.** Scope of Alkyne Addition

entry	alkyne	R	product	yield (%)
1	<b>5a</b>	<i>n</i> -butyl	<b>6</b>	84
2	<b>5b</b>	<i>i</i> -Pr	<b>7</b>	75
3	<b>5c</b>	<i>t</i> -Bu	<b>8</b>	73
4	<b>5d</b>	Ph	<b>9</b>	77
5	<b>5e</b>	CH <sub>2</sub> OBn	<b>10</b>	76
6	<b>5f</b>	CH <sub>2</sub> OTIPS	<b>11</b>	76


3). The reaction is facile at 0 °C, and the alkyne can accommodate linear or branched alkyl groups (entries 1–3) as well as aryl substitution (entry 4). Propargyl systems employing benzyl and triisopropylsilyl protecting groups smoothly afford the desired carbinols in good yields.

Under Lewis base-catalyzed conditions (10 mol % of KOEt), triethoxysilylalkyne **5a** undergoes facile addition to various aldehydes in good yields (eq 4, Table 3). The reaction is high yielding with numerous electron-rich and electron-deficient aromatic aldehydes, is mild enough to accommodate enolizable aldehydes (entries 9 and 10), and notably affords selective additions in the presence of esters (entry 11).<sup>14</sup>

A distinctive and important attribute of this process is the capability of this new alkynyl nucleophilic reagent to undergo

(13) The control experiment with 1-hexyne, KOEt, and aldehyde does not produce desired propargyl alcohol under the reaction conditions. The addition of Si(OEt)<sub>4</sub> in this system yields no product either. See the Supporting Information for details.

(14) In contrast, the lithium alkyne addition to methyl 4-formylbenzoate affords a complex mixture from the addition to both carbonyl carbons.


**Table 3.** Triethoxysilylhexyne (**5a**) Additions to Aldehydes

entry	RCHO	product	yield (%) <sup>a</sup>
1			<b>12</b> 74
2			<b>6</b> 84
3			<b>13</b> 82
4			<b>14</b> 76
5			<b>15</b> 76
6			<b>16</b> 80
7			<b>17</b> 85
8 <sup>b</sup>			<b>18</b> 59
9			<b>19</b> 56
10			<b>20</b> 58
11			<b>21</b> 78
12			<b>22</b> 93

<sup>a</sup> Isolated yields after chromatography. <sup>b</sup> >95% (*E*)-cinnamaldehyde.

addition to ketones (Table 4, eq 5). The conditions that afford the best yields employ a higher catalyst loading (20 mol %) and a crown ether (18-crown-6).<sup>15</sup> By utilizing these Lewis basic conditions, undesired aldol products are not observed with enolizable ketones as substrates, indicating that the

(15) The addition of 18-crown-6 allows for catalytic KOEt to be used and presumably promotes turnover of the nucleophilic alkoxide product. See the Supporting Information for details.


**Table 4.** Triethoxysilylhexyne (**5a**) Additions to Ketones

entry	ketone	product	yield (%) <sup>b</sup>
1			<b>23</b> 73
2 <sup>c</sup>			<b>24</b> 54 <sup>d</sup>
3			<b>25</b> 54
4			<b>26</b> 71
5			<b>27</b> 85
6			<b>28</b> 64
7			<b>29</b> 73

<sup>a</sup> 20 mol % of 18-crown-6. <sup>b</sup> Isolated yields after chromatography. <sup>c</sup> >95% (*E*)-chalcone. <sup>d</sup> Only 1,2-addition product observed.

basicity of these new reagents is attenuated relative to standard alkynyl organometallic reagents.

Our preliminary investigations of this process have provided important clues about the operative reaction mechanism. We propose that the active alkynyl nucleophile is a hypervalent silicate intermediate resulting from reversible addition of alkoxide. Evidence for this species was obtained by the low-temperature <sup>29</sup>Si NMR experiment of combining trialkoxysilylalkyne **5a**, 1.0 equiv of KOEt, and 1.0 equiv of 18-crown-6 (Figure 1). A new and distinct signal at −126

**Figure 1.** <sup>29</sup>Si NMR (−60 °C) of **5a**, KOEt, and 18-crown-6 (1:1:1).


ppm is similar to previously studied pentavalent silicate intermediates such as  $(\text{EtO})_4\text{SiPh}\cdot\text{K}$  (−117 ppm).<sup>16</sup>

An intriguing aspect of the reaction is that the propargyl product is initially an alkoxide and may well activate the

(16) Swamy, K. C. K.; Chandrasekhar, V.; Harland, J. J.; Holmes, J. M.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 2341–2348.

trialkoxysilylalkyne in a manner similar to KOEt. To probe this product-activation possibility, the potassium salt of propargyl alcohol **6** was added (10 mol %) to silylalkyne **5a** and benzaldehyde (Scheme 1). Interestingly, propargyl


Scheme 1



alcohol **14** is isolated in good yield after aqueous workup (74%).

Drawing from the data observed in both Figure 1 and Scheme 1, we propose a plausible mechanistic description of this reaction involving Lewis base initiation of an autocatalytic cycle (Scheme 2). The addition of a Lewis base

Scheme 2



to **5** would lead to formation of a pentavalent silicon intermediate **II**, which could mediate the transfer of an alkynyl nucleophile via hexacoordinate organization (**III**).


The resulting silylated alkyne-addition product (**IV**) could dissociate, leading to generation of alkoxide base **I**, thereby perpetuating the reaction. Further evidence for self-promotion was the direct observation of **V** by gas chromatography under catalytic reaction conditions.<sup>17</sup> A mechanism involving self-promotion is also supported by the necessity of crown ether

(17) Generation of **V** is not observed in the presence of stoichiometric amounts of potassium ethoxide.

for catalytic turnover in the addition to ketones, where a more sterically congested tertiary propargyl alkoxide (**I**) is generated.<sup>15</sup>

The unique properties and synthetic potential of these trialkoxysilylalkynes can be observed in the formation of secondary propargylamines<sup>18</sup> (Scheme 3). The combination

Scheme 3



of **5a** and *tert*-butylsulfinyl imine **30**<sup>19</sup> affords alkyne **31** in exceptional yield (95%) with high selectivity (20:1 favoring the (*S<sub>S</sub>*,*R*) diastereomer. Surprisingly, alternative alkynyl organometallic nucleophiles are either less selective (Met = K) or prefer the *opposite* stereoisomer (Met = Li or MgEt).<sup>11</sup>

In conclusion, we have reported an efficient nucleophile-catalyzed addition reaction of alkynes. This new strategy utilizes trialkoxysilylalkynes as stable nucleophile precursors with preliminary mechanistic data implicating a hypervalent organosilane as the active reagent. The new alkynyl species accessed by the addition of a Lewis base possess reactivity to undergo smooth, and in some cases highly selective, additions to aldehydes, ketones, and imines. Mechanistic investigations and applications of this Lewis base-catalyzed strategy based on trialkoxyorganosilane activation are currently underway.

**Acknowledgment.** This work has been partially supported by Northwestern University and the NSF (CAREER Award to K.A.S.). We are grateful to Abbott Laboratories, Amgen, 3M, and Boehringer-Ingelheim for research support and to FMCLithium for providing organolithium reagents used in this research. We thank Mr. David Ballweg for assistance with preparing this manuscript.

**Supporting Information Available:** Experimental procedures and spectral data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL051030F

(18) Asymmetric alkyne additions to imines: (a) Gommermann, N.; Knochel, P. *Chem. Commun.* **2004**, 2324–2325. (b) Fassler, R.; Frantz, D. E.; Oetiker, J.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2002**, *41*, 3054–3056. (c) Traverse, J. F.; Hoveyda, A. H.; Snapper, M. L. *Org. Lett.* **2003**, *5*, 3273–3275.

(19) Ellman, J. A.; Owens, T. D.; Tang, T. P. *Acc. Chem. Res.* **2002**, *35*, 984–995.