

Cycloisomerization of Dienes with Carbophilic Lewis Acids: Mimicking Terpene Biosynthesis with Pt(II) Catalysts

William D. Kerber and Michel R. Gagné*

Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599-3290

mgagne@unc.edu

Received May 31, 2005

ABSTRACT



Dicationic Pt(II) complexes containing triphosphine pincer ligands are excellent catalysts for the cycloisomerization of 1,6- and 1,7-dienes into bicyclopropane carbocycles. In analogy to the biosynthetic route to these monoterpene-like compounds, carbocation intermediates are proposed and supported by trapping experiments. Reactivation of the trapped intermediates indicates that cation generation by C–C bond formation is both rapid and reversible.

The generation of carbocations from unsaturated poly-prenoids, followed by C–C bond/ring-forming cation-olefin reactions, constitutes the key mechanism for the biosynthesis of most, if not all, of the terpene-derived natural products.¹ Under enzyme control, the position of cation generation, the sequence and regioselectivity of the C–C bond-forming steps, and the method of cation quenching all contribute to the skeletal structure and functionality of the cyclization products.

Electrophilic Pt(II) dications supported by a tridentate pincer ligand have recently been shown to catalyze several of the reactions typically associated with terpene biosynthesis (cation generation, cation-olefin, hydride shift, cyclopropanation).² Key to the activity of these catalysts is their highly electrophilic, but also carbophilic, electronic structures, coupled with the all-*cis* pincer ligands, which block β -hydride elimination in any Pt-alkyl intermediates. The result is a buildup of intermediates that follow pathways not otherwise

competitive in catalysts capable of facile β -H elimination. In this communication, we report that a family of 1,5- and 1,6-dienes reacts under Pt-catalysis in a fashion that seemingly parallels the biosynthesis of the [3.1.0] bicyclic monoterpenes. Additionally, we disclose a trapping experiment that supports the intermediacy of organometallic carbocations.


The [3.1.0] bicyclic skeleton is the key structural feature of many monoterpenes (thujane, thujene, sabinene, sabinene hydrate, sabina ketone, etc.), which are found in numerous essential oils and consequently find importance in flavors and fragrances.³ Considerable effort has been devoted to the construction of this ring system, and the available methods include both intra-⁴ and intermolecular⁵ addition of carbonyls to alkenes, intramolecular alkylation of enolates,⁶ a tandem Li-ene cyclization/thiophenoxide elimination,⁷ a Ti-mediated enyne cyclization,⁸ and an unusual rearrangement after reaction of silylketene acetals with bromoform-

(1) In *Biosynthesis of Isoprenoid Compounds*; Porter, J. W., Spurgeon, S. L., Eds.; John Wiley & Sons: New York, 1981; Vol. 1.

(2) (a) Hahn, C.; Cucciolito, M. E.; Vitagliano, A. *J. Am. Chem. Soc.* **2002**, *124*, 9038. (b) Kerber, W. D.; Koh, J. H.; Gagné, M. R. *Org. Lett.* **2004**, *6*, 3013.

(3) (a) Croteau, R. *Chem. Rev.* **1987**, *87*, 929–954. (b) Croteau, R. In *Recent Developments in Flavor and Fragrance Chemistry: Proceedings of the 3rd International Harman & Reimer Symposium*; VCH: Weinheim, 1993; p 263.

diethylzinc.⁹ Each of these approaches requires multiple steps and usually leads to oxygenated monoterpenes that are most amenable to the synthesis of the oxidized congeners.



On the basis of previous cycloisomerization studies with **1**, we postulated that it should also be capable of directly providing the [3.1.0]-bicyclic core¹⁰ from simple 1,6-dienes. Gratifyingly, a variety of dienes with the substitution pattern shown in Table 1 are converted to the saturated bicyclic

Table 1. Cycloisomerization of Dienes with 5 mol % of **1**^a

	diene	t (h)	yield ^b	product
1		25	61	
2		15	65	
3		12	53	
4		36	53	
5		5 ^d	11 ^e	

^a Compound **1** generated in situ from [(PPP)PtMe]BF₄, acetone (1 equiv), and HNTf₂ in MeNO₂, 5% catalyst loading, 40 °C. See Supporting Information. ^b Isolated. ^c By GC. ^d 23 °C. ^e By GC; successive chromatography afforded pure **11** in 5% isolated yield.


products when exposed to 5 mol % of **1**. In the case of the 1,6-diene β -citronellene (**4**), a highly diastereoselective

rearrangement occurs and the natural product *cis*-thujane¹¹ (**5**) is obtained in 47:1 dr.

In keeping with the high carbophilicity of these strong Lewis acids, ester-containing compounds did not poison the catalyst and the expected product was obtained (entry 3). The [4.1.0] bicyclic compound **9** could also be obtained but at a decreased rate (entry 4). A [3.1.0] bicyclic product was also obtained from the cycloisomerization of *O*-Bn linalool **10** (entry 5). In this case, the major product resulted from allylic isomerization to a mixture of *O*-Bn geraniol/nerol; however, a single diastereomer of **11** was also obtained in low yield. A skeletal rearrangement has obviously taken place that will require additional study and optimization. Despite the moderate isolated yields,¹² these one-step reactions represent the most efficient syntheses of the [3.1.0] bicyclic core from simple acyclic starting materials.

The initiating step in the cycloisomerization reactions was presumed to involve electrophilic activation of the terminal alkene and intramolecular addition of the trisubstituted alkene to cyclogenerate intermediate carbocations that were stable to β -H elimination.² These putative Pt(II)-alkyl carbocations had not been observed directly, but we hoped that a trapping nucleophile could intercept them for characterization. In the event, excess benzyl alcohol and Ph₂NMe rapidly converted **12**, the catalyst's resting state,¹³ to a mixture of endo- and exocyclic Pt-alkyls.^{14–16} The composition of this mixture changed with time to ultimately favor the endocyclic product **13a** (see table in Scheme 1).¹⁷ Characterization of the

Scheme 1. Trapping of Carbocationic Intermediates



carbocyclic fragment was achieved by reductive removal with NaBH₄.^{15,18,19}

The data in Scheme 1 indicate not only that C–C bond formation was reversible but that both endo- and exocyclic

(4) (a) Hodgson, D. M.; Chung, Y. K.; Paris, J. M. *J. Am. Chem. Soc.* **2004**, *126*, 8664. (b) Harrak, Y.; Blaszykowski, C.; Bernard, M.; Cariou, K.; Mainetti, E.; Mouries, V.; Dhiman, A.-L.; Fensterbank, L.; Malacria, M. *J. Am. Chem. Soc.* **2004**, *126*, 8656. (c) Barberis, M.; Perez-Prieto, J. *Tetrahedron Lett.* **2003**, *44*, 6683. (d) Mori, K.; Ohki, M.; Matsui, M. *Tetrahedron* **1970**, *26*, 2821. (e) Vig, O. P.; Bhatia, M. S.; Gupta, K. C.; Matta, K. L. *J. Ind. Chem. Soc.* **1969**, *46*, 991.

(5) (a) Galopin, C. C. *Tetrahedron Lett.* **2001**, *42*, 5589. (b) Cheng, D.; Krethadumrongdat, T.; Cohen, T. *Org. Lett.* **2001**, *3*, 2121. (c) Fanta, W. I.; Erman, W. F. *J. Org. Chem.* **1968**, *33*, 1656. (d) Guha, P. C.; Krishnamurthy, S. *Chem. Ber.* **1937**, *70*, 0B, 2112.

(6) (a) Hamon, D. P. G.; Shirley, N. J. *Aust. J. Chem.* **1987**, *40*, 1321. (b) Gaoni, Y. *Tetrahedron* **1972**, *28*, 5525. (c) Alexandre, C.; Rouessac, F. *Bull. Chem. Soc. Jpn.* **1972**, *45*, 2241. (d) Nelson, N. A.; Mortimer, G. A. *J. Org. Chem.* **1957**, *22*, 1146.

(7) Cheng, D.; Knox, K. R.; Cohen, T. *J. Am. Chem. Soc.* **2000**, *122*, 412.

(8) Urabe, H.; Suzuki, K.; Sato, F. *J. Am. Chem. Soc.* **1997**, *119*, 10014.

(9) Rousseau, G.; Slougui, N. *J. Am. Chem. Soc.* **1984**, *106*, 7283.


(10) Ene-yne cycloisomerization with PtCl₂ and Au(III) catalysts can also generate (through different mechanisms) related bicyclopropane structures. See, for example: (a) Harrak, Y.; Blaszykowski, C.; Bernard, M.; Cariou, K.; Mainetti, E.; Mouries, V.; Dhiman, A.-L.; Fensterbank, L.; Malacria, M. *J. Am. Chem. Soc.* **2004**, *126*, 8656. (b) Mamane, V.; Gress, T.; Krause, H.; Fürstner, A. *J. Am. Chem. Soc.* **2004**, *126*, 8654. (c) Fürstner, A.; Hennen, P. *Chem. Commun.* **2004**, 2546. (d) Nieto-Oberhuber, C.; Munoz, M. P.; Bunuel, E.; Nevado, C.; Cardenas, D. J.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2004**, *43*, 2402. (e) Luzung, M. R.; Markham, J. P.; Toste, F. D. *J. Am. Chem. Soc.* **2004**, *126*, 10858.

(11) Rees, J. C.; Whittaker, D. *Org. Magn. Reson.* **1981**, *15*, 363.

(12) The balance of material in these reactions is usually alkene-containing isomers that appear to originate from the consumption of cyclopropane.


(13) The trapping reactions are complete within minutes as observed by ³¹P NMR.

regiochemistries were kinetically viable. When **13a** was treated with 1 equiv of the strong Lewis acid $B(C_6F_5)_3$, benzyloxide abstraction occurred concomitant with retro-cyclization to **12** (eq 1) (**13a**:**12** ~3:1 at early times).²⁰ This



mixture is eventually driven to **3** and $(\text{PPP})\text{Pt}(\text{OBn})^+$.²¹ The observation that **13a** returns to the catalyst resting state (**12**) upon OBn abstraction and the time-dependent ratio of **13a**/**13b** together suggest that cyclization is rapid and reversible and that a postcyclization step (H^- shift or cyclopropanation) is turnover-limiting.

Since both endo- and exocyclization geometries are demonstrably viable, two plausible mechanisms for the cycloisomerization of **2** to **3** must be considered (Scheme 2). The two scenarios differ in the timing of ring construction,



with the 6-endo pathway (**b**) most closely following the sequence of thujane biosynthesis.³ Distinguishing the two mechanisms may require computational help, but in both cases the cyclopropanation event is proposed to proceed via

(14) This nomenclature refers to the position of the electrophilic alkene in the C—C bond-forming transition state.

(15) Both 6-exo- and 6-endo-cyclization modes were observed in stoichiometric cascade cyclization of polyenes. See: Koh, J. H.; Gagné, M. R. *Angew. Chem., Int. Ed.* **2004**, 43, 3459–3461.

a key carbocation that is γ to the metal (from a 1,2-hydride shift of the kinetic cation). Related, stereospecific, double inversion processes are known for Sn-,²² Fe-,²³ and Ti-alkyls²⁴ with γ -carbocations.²⁵

We have described the cycloisomerization of dienes to [3.1.0] and [4.1.0] bicyclic compounds, including the one-step synthesis of the natural product *cis*-thujane. The catalyst for these reactions, a Lewis acidic Pt(II) pincer complex, activates alkenes for cation-olefin cyclizations, as evidenced by the trapping of intermediate organo-platinum carbocations.

Acknowledgment. We thank Mr. Au Jin Kim for synthesizing **6** and Dr. Gary Glish for performing CI MS and gratefully acknowledge the NIGMS (GM60578) for funding. W.D.K thanks Glaxo-Smith-Kline for a graduate fellowship. M.R.G. is a Camille-Dreyfus Teacher-Scholar.

Supporting Information Available: Full experimental procedures, including catalytic methods and characterization of products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL051277C

(16) For catalytic asymmetric alkoxymercuration of enynes with Pt(II) and Au(I), see: (a) Charruault, L.; Véronique, M.; Taras, R.; Gladiali, S.; Genêt, J.-P. *Chem. Commun.* **2004**, 851. (b) Muñoz, M. P.; Adrio, J.; Carretero, J. C.; Echavarren, A. M. *Organometallics* **2005**, 24, 1293.

(17) The balance of the material was two Pt-alkyls that remain uncharacterized.

(18) Hahn, C.; Morvillo, P.; Vitagliano, A. *Eur. J. Inorg. Chem.* **2001**, 419.

(19) A single diastereomer was obtained from reductive cleavage of **13b**. Although *trans* relative stereochemistry seems most likely, this has not been conclusively determined.

(20) A similar OBn abstraction likely mediates the isomerization of **10** to *O*-Bn geraniol/nerol (entry 5, Table 1).

(21) As **2** is consumed, **1** abstracts alkoxide from boron to generate $[(\text{PPP})\text{Pt}(\text{OBn})]^+$.

(22) (a) Davis, D. D.; Johnson, H. T. *J. Am. Chem. Soc.* **1974**, 96, 7576. (b) Fleming, I.; Urch, C. J. *Tetrahedron Lett.* **1983**, 24, 4591. (c) McWilliam, D. C.; Balasubramanian, T. R.; Kuivila, H. G. *J. Am. Chem. Soc.* **1978**, 100, 6407. (d) Lambert, J. B.; Salvador, L. A.; So, J. H. *Organometallics* **1993**, 12, 697.

(23) (a) Casey, C. P.; Smith, L. J. *Organometallics* **1989**, 8, 2288. (b) Casey, C. P.; Smith Vosejka, L. J. *Organometallics* **1992**, 11, 738. (c) Brookhart, M.; Liu, Y.; Goldman, E. W.; Timmers, D. A.; Williams, G. D. *J. Am. Chem. Soc.* **1991**, 113, 7. (d) Brookhart, M.; Liu, Y. *J. Am. Chem. Soc.* **1991**, 939.

(24) Casey, C. P.; Strotman, N. A. *J. Am. Chem. Soc.* **2004**, 126, 1699.

(25) An alternative possibility is direct metalation to form a Pt(IV) metallacyclobutane followed by reductive elimination. Evidence disfavoring this mechanism includes the observation of cyclopropane products with related Pd(II) catalysts, which would implicate the involvement of a Pd(IV) intermediate; see ref 2b. However, this possibility has not been rigorously disproven.