

Photodimerizable Ditopic Ligand


Damien Jouvenot, Edith C. Glazer, and Yitzhak Tor*

*Department of Chemistry and Biochemistry, University of California, San Diego,
La Jolla, California 92093*

ytor@ucsd.edu

Received January 29, 2006

ABSTRACT



The synthesis, photophysical properties, and structural characterization of a photodimerizable ditopic ligand are described. Upon irradiation at 366 nm, ligand **1** dimerizes to the head-to-tail tetra-bpy ligand **2**. This thermally stable photodimer can be dissociated back to **1** using higher energy irradiation (254 nm).

Bridging ligands, capable of coordinating multiple metal ions, have been attracting considerable interest because of their potential use as building blocks for the fabrication of functional multimetallic assemblies.¹ Of particular interest are systems that can respond to external stimuli, such as photons,^{2,3} protons,^{4,5} electrons,⁶ or other reactive species. Such responsive ligands and their complexes can, in principle, be employed for the on/off switching or fine tuning of the interactions between the coordinating centers. In contrast to the abundance of unreactive polypyridine multitopic ligands, the repertoire of responsive ditopic bridging ligands that can be modulated by external stimuli is limited.

(1) (a) Ruben, M.; Rojo, J.; Romero-Salguero, F. J.; Uppadine, L. H.; Lehn, J.-M. *Angew. Chem., Int. Ed.* **2004**, *43*, 3644–3662. (b) Sauvage, J.-P.; Collin, J.-P.; Chambron, J.-C.; Guillerez, S.; Coudret, C.; Balzani, V.; Barigelletti, F.; De Cola, L.; Flamigni, L. *Chem. Rev.* **1994**, *94*, 993–1019.

(2) Collin, J.-P.; Jouvenot, D.; Koizumi, M.; Sauvage, J.-P. *Eur. J. Inorg. Chem.* **2005**, 1850–1855.

(3) Brouwer, A.; Frochot, C.; Gatti, F. G.; Leigh, D. A.; Mottier, L.; Paolucci, F.; Roffia, S.; Wurpel, G. W. H. *Science* **2001**, *291*, 2124–2128.

(4) Sakamoto, A.; Hirooka, A.; Namiki, K.; Kurihara, M.; Murata, M.; Sugimoto, M.; Nishihara, H. *Inorg. Chem.* **2005**, *44*, 7547–7558.

(5) Haga, M.-A.; Takasugi, T.; Tomie, A.; Ishizuya, M.; Yamada, T.; Hossain, M. D.; Inoue, M. *J. Chem. Soc., Dalton Trans.* **2003**, *10*, 2069–2079.

(6) Zelikovich, L.; Libman, J.; Shanzer, A. *Nature* **1995**, *374*, 790–792.

Here, we report the synthesis and reactivity of a new photodimerizable ligand **1**, where a diimine core is incorporated into an anthracene unit. Near-ultraviolet irradiation ($\lambda \sim 360$ nm) results in exclusive formation of a head-to-tail tetra-bpy ligand **2**.

Anthracene photodimerization is one of the oldest photochemical reactions known.⁷ Since it was first observed in 1867, it has been employed in numerous responsive systems, including metal coordinating ones.⁸ Most anthracene-containing ligands, however, consist of isolated coordinating sites that are covalently linked to an anthracene unit, where the latter serves as the dimerizing module only. Examples include anthracene-containing crown ethers or cryptands,^{9,10} as well as systems with pendant polypyridyl ligands.^{11,12} In contrast to the majority of these systems, ligand **1** encompasses a conjugating diazaanthracene unit that serves as the common core of the ditopic bis-bpy ligand. As such, the electronic

(7) Becker, H. D. *Chem. Rev.* **1993**, *93*, 145–172.


(8) Bouas-Laurent, H.; Castellan, A.; Desvergne, J. P.; Lapouyade, R. *Chem. Soc. Rev.* **2000**, *29*, 43–55.

(9) (a) Hiraga, H.; Morozumi, T.; Nakamura, H. *Eur. J. Org. Chem.* **2004**, 4680–4687. (b) Desvergne, J. P.; Lahrahar, N.; Gotta, M.; Zimmermann, Y.; Bouas-Laurent, H. *J. Mater. Chem.* **2005**, *15*, 2873–2880.

(10) Another system based on a [2.2](9,10)anthracenophane is used to coordinate [Ru(Cp)]⁺ in an η^6 fashion. See: Glatzhofer, D. T.; Liang, Y.; Khan, M. A. *J. Chem. Soc., Chem. Commun.* **1993**, 742–743.

properties of the ligand and its metal complexes are expected to change upon photodimerization.

To facilitate the synthesis of the bridging ligand **1**, a concise synthesis of the hitherto unknown dibromodiazanthracene building block **7** was needed (Scheme 1). It was



prepared from the diester **3** which, in turn, can be made from resorcinol using established procedures.^{13,14} Following activation of the phenolic groups as triflates to give **4**, a Sonogashira cross-coupling reaction with trimethylsilylacetylene yielded **5**. A sealed tube reaction with saturated ethanolic ammonia afforded the bislactam **6**.¹⁵ As **6** is rather insoluble, a simple filtration provided the pure product. Bromination of the lactam using phosphorus oxybromide and potassium carbonate in acetonitrile¹⁶ produced the desired 1,8-dibromo-2,7-diazanthracene **7**. These milder conditions were necessary because refluxing the bislactam in pure phosphorus oxybromide led primarily to the tribrominated derivative. Final assembly of the bridging ligand **1** was achieved by a Negishi cross-coupling reaction¹⁷ between 2-bromopyridine and the dibromo derivative **7** to give the desired product in 88% yield (Scheme 1).^{14,18,19}

Single crystals of **1**, suitable for X-ray crystallography, were obtained by slowly diffusing diethyl ether into a




Figure 1. Crystal structure of **1**.

solution of the ligand in chloroform in the dark. As seen in the solved structure (Figure 1), the bipyridine units adopt the expected trans geometry, thus minimizing the global dipole moment of the molecule and the electrostatic interaction between the nitrogen atoms' lone pairs. In addition, the diazanthracene core and the individual pyridine rings are also out of plane (torsion angles: 134.1° and 151.5°). Inspection of the crystal lattice reveals intermolecular contacts between the pyridinic moieties of the anthracene unit. The average distance between planes is 3.44 \AA , allowing for an efficient π - π stacking along the *b* axis. Note, however, the absence of stacked diazanthracene units, an arrangement that is likely to be necessary for photodimerization in the solid state.²⁰

The absorption spectrum of ligand **1** is shown in Figure 2a. It is characterized by an intense absorption at 250 nm with a shoulder centered around 290 nm . There is also a broad, featureless band tailing into the visible region with a maximum at 392 nm . The molecule's emission spectrum (Figure 2a, inset) lacks the fine features associated with typical anthracene derivatives, exhibiting a broad band with a λ_{max} at 470 nm .²¹ The excitation spectrum agrees well with the absorption spectrum, with a peak maximum at 397 nm (Figure 2a, inset).

When crystallization attempts were carried out in the presence of light, the photodimer **2**, less soluble than **1**, crystallized out exclusively. The head-to-tail dimeric structure of the photoproduct could be unambiguously determined by X-ray crystallography (Figure 3). As with the parent ligand, the bipyridine units adopt an expected trans geometry with significant deviation from coplanarity (torsion angles: 143.0° and 165.4°). In the crystal structure, the dimer displays an inversion center located in the center of the molecule. The crystal packing is ensured mainly by two π - π interactions (Figure 4). The principal π - π stacking is observed between

(11) Beyerler, A.; Belser, P.; De Cola, L. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2779–2781.

(12) Trouts, T. D.; Tyson, D. S.; Pohl, R.; Kozlov, D. V.; Waldron, A. G.; Castellano, F. N. *Adv. Funct. Mater.* **2003**, *13*, 398–402.

(13) Zeng, H.; Miller, R. S.; Flowers, R. A.; Gong, B. *J. Am. Chem. Soc.* **2000**, *122*, 2635–2644.

(14) See Supporting Information for additional experimental details and data.

(15) Sakamoto, T.; An-naka, M.; Kondo, Y.; Yamanaka, H. *Chem. Pharm. Bull.* **1986**, *34*, 2754–2759.

(16) Janin, Y. L.; Roulland, E.; Beurdeley-Thomas, A.; Decaudin, D.; Monneret, C.; Poupon, M.-F. *J. Chem. Soc., Perkin Trans. 1* **2002**, 529–532.

(17) Loren, J. C.; Siegel, J. S. *Angew. Chem., Int. Ed.* **2001**, *40*, 754–757.

(18) A successful double cross-coupling reaction requires thoroughly dried zinc chloride (24 h at 55°C under reduced pressure).

(19) Interestingly, the usual stain employed to visualize bipyridines ($\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2$ in water) does not yield the typical red color but rather a bright green color. See Supporting Information Figure S.15.

(20) Single-crystal–single-crystal reactions are very rare for anthracene derivatives; however, dimerization can be performed in the crystalline state followed by recrystallization. See: Turowska-Tyrk, I.; Grzesniak, K. *Acta Crystallogr.* **2004**, *C60*, o146–o148.

(21) The emission quantum yield was determined to be 0.02.




Figure 2. (a) Absorption spectrum of ligand **1** in CH_3CN . Inset: emission (blue line) (excitation at 395 nm) and excitation (green line) (monitoring at 470 nm) spectra of **1** in CH_3CN . (b) Fluorescence intensity continuously decreases upon irradiation at 366 nm: $t = 0$ (red line); $t = 45$ min (blue line); $t = 90$ min (green line); $t = 180$ min (black line); $t = 360$ min (yellow line). Sample temperature is initially stabilized at 25°C . A photostationary state can be estimated around 50%. Irradiation was performed in degassed acetonitrile solution in a quartz cell with a portable UV lamp of 4 W at 366 nm.

two adjacent bipyridine fragments with a distance of 3.53 \AA between the planes. An additional π - π interaction involves the other pyridyl substituent stacking with its analogue of the neighboring molecule. This effect is weaker as the interplanar separation is 3.62 \AA .




Figure 3. Crystal structure of ligand **2**.




Figure 4. Packing of ligand **2** in the crystal.

Upon irradiation at 366 nm, ligand **1** undergoes dimerization at the 9 and 10 carbons. This photochemical transformation can be easily monitored by fluorescence spectroscopy as the anthracene core responsible for the molecule's emission at 470 nm is disrupted upon dimerization, thereby providing a probe for following the progress of the reaction (Figure 2b).²² A considerable amount of dimer **2** was prepared by irradiation (366 nm) of ligand **1** in a degassed acetonitrile solution followed by purification using preparative TLC.¹⁴ Most signals in the ^1H NMR spectrum of the dimer **2** feature an upfield shift when compared to the corresponding protons in ligand **1** (Figure 5). As expected, protons




Figure 5. ^1H NMR spectra in CD_2Cl_2 and assignment of ligand **1** (top) and the photodimer **2** (bottom).

9 and 10 display the most dramatic shift as their hybridization changes from sp^2 to sp^3 (Figure 5, blue line). Their multiplicity also changes from singlets to doublets, as expected for a head-to-tail dimer. A group of three protons (4', 5', and 6') are only slightly shifted (red line), consistent with a minimal change in their environment upon dimerization. Protons 3 and 4, along with proton 3', experience a significant shift (0.4–0.6 ppm) (green line). For protons 3 and 4, the significant shielding can be attributed to ring current effects of the aromatic rings located directly below. On the other hand, the shift of proton 3' could be attributed to the absence of the deshielding cone produced by the central ring as it loses aromaticity upon dimerization.

The photodimer **2** was found to be thermally stable, with no indication for dissociation to monomer **1** over 12 h in solution at room temperature. Preliminary results suggest, however, that the dimerization may be reversed by irradiation at 254 nm (Scheme 2 and Supporting Information Figure

S.12).¹⁴ This finding is in agreement with previous literature reports of photoreversible dimerization.^{11,23} Note, however, that further optimization is required because under the current unoptimized conditions the reaction is only partially reversible and repetitive cycling appears to result in sample degradation.¹⁴


To the best of our knowledge, ligand **1** is the first example of a photodimerizable ligand, in which the diimine core is embedded within the bridging anthracene unit. Akin to such systems that undergo photoinduced structural modifications (e.g., *cis*–*trans* isomerization,²⁴ photocyclization²⁵), this paradigm system could be considered as a photoswitch to which added properties may eventually be incorporated upon binding to metal ions.

Acknowledgment. The authors thank the National Institutes of Health (GM 069773) and NSF (CHE 0213323) for support.

Supporting Information Available: Full synthetic details and analytical data for all new compounds; photoinduced dimerization and dissociation; crystal data and refinement information for **1** and **2**; spectra of mono- and dinuclear Ru^{2+} complexes of **1**; and reactions of **1** and **2** with Fe^{2+} . This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL060253I

Scheme 2. Reversible Photoinduced Dimerization and Dissociation Reactions



(22) Note the photodimer **2** is nonemissive under these conditions.

(23) European Patent Application 0 394 846 A2.

(24) Shinkai, S.; Nakaji, T.; Nishida, Y.; Ogawa, T.; Manabe, O. *J. Am. Chem. Soc.* **1980**, *102*, 5860–5865.

(25) (a) Irie, M.; Fukaminato, T.; Sasaki, T.; Tamai, N.; Kawai, T. *Nature* **2002**, *420*, 759–760. (b) Fraysse, S.; Coudret, C.; Launay, J.-P. *Eur. J. Inorg. Chem.* **2000**, 1581–1590.