

Stereoselective Synthesis of Pyridinones: Application to the Synthesis of (–)-Barrenazines


Thilo Focken and André B. Charette*

Department of Chemistry, Université de Montréal, P.O. Box 6128, Station Downtown, Montréal, Quebec, H3C 3J7, Canada

andre.charette@umontreal.ca

Received April 13, 2006

ABSTRACT



The stereoselective synthesis of pyridinones was accomplished by the nucleophilic addition of Grignard reagents to a chiral pyridinium salt derived from 4-methoxyphenylpyridine. This methodology was applied to an expedient synthesis of (–)-barrenazine A and B. After N-functionalization and 1,4-reduction of the pyridinone system, the corresponding α -amino piperidinones readily undergo dimerization to give the hexahydrodipyrindopyrazine skeleton of the barrenazine alkaloids.

In 2003, Kashman and co-workers isolated two novel cytotoxic alkaloids, barrenazine A (**1**) and B (**2**), from an unidentified tunicate collected at Barren Islands in Madagascar.¹ These two compounds have a unique 1,3,4,6,8,9-hexahydrodipyrindino[3,4-*b*:3',4'-*e*]pyrazine skeleton. Related naturally occurring structures are rare, and among those are the hexahydrodipyrano-pyrazines palythazin and isopalythazine^{2c} and the trihydropyrano-pyrazine clavulazine.^{2a} A synthesis of palythazine in six steps was reported starting from tetra-O-benzoylated D-glucose.^{2b} Herein, we report an expedient, enantioselective synthesis of both barrenazines from a common intermediate.

Our retrosynthetic analysis of barrenazines A and B is depicted in Figure 1. It was envisioned that the pyrazine core could be assembled through the dimerization of amino pyridinone **3**. Fragment **3** would be elaborated by the stereoselective functionalization of 4-methoxyphenylpyridine (**4**).

A strategy that has been frequently used to prepare substituted piperidine and piperidinone alkaloids³ consists

of the addition of a nucleophile to a chiral or achiral *N*-alkyl- or *N*-acylpyridinium salt.⁴ As part of our ongoing research program directed toward the expedient synthesis of enantiopure, polysubstituted piperidines from readily available starting materials, we developed a new methodology for the generation of pyridinium salts from pyridine, secondary amide **5**, and triflic anhydride.⁵ These salts undergo highly diastereo- and regioselective addition of carbon nucleophiles to produce dihydropyridines.





Figure 1. Retrosynthetic analysis of barrenazine A and B.

(1) Chill, L.; Akin, M.; Kashman, Y. *Org. Lett.* **2003**, *5*, 2433–2435.
(2) (a) Watanabe, K.; Iguchi, K.; Fujimori, K. *Heterocycles* **1998**, *49*, 269–274. (b) Jarglis, P.; Lichtenthaler, F. W. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 141–142. (c) Uemura, D.; Toya, Y.; Watanabe, I.; Hirata, Y. *Chem. Lett.* **1979**, *12*, 1481–1482.

The addition of 4-methoxypyridine (**4**, 2.3 equiv) to a mixture of amide **5** (1.0 equiv) and triflic anhydride (1.1 equiv) at $-78\text{ }^{\circ}\text{C}$ led to the formation of the desired pyridinium salt (Table 1).

Table 1. Nucleophilic Addition to Pyridinium Salts Derived from 4-Methoxypyridine (**4**) and Amide **5**



entry	RMgBr	product	yield (%) ^a	dr ^b
1	Me	6a	85	95:5
2	Et	6b	76	93:7
3	<i>n</i> -Bu	6c	71	>95:5
4	<i>t</i> -Bu	6d	61	93:7
5	(CH ₂) ₆ OTBS	6e	70	>95:5
6	(CH ₂) ₅ CH=CH ₂	6f	84 ^c	93:7
7	CH=CH ₂	6g	64	92:8
8	C≡CCH ₃	6h	52 + 8	83:17
9	C≡CPh	6i	65	89:11
10	Ph	6j	66	91:9
11	furanyl	6k	65	95:5

^a Products isolated as a single diastereomer. ^b Determined by ¹H NMR of the crude product. ^c Mixture of diastereomers (dr = 13:1).

NMR monitoring of the pyridinium salt formation⁶ indicated that the activation process was complete within 5 h at room temperature. The pyridinium salt solution was then cooled to $-20\text{ }^{\circ}\text{C}$, and an ether solution of the Grignard reagent was added. After acidic hydrolysis of the intermediate methyl enol ethers, pyridinones **6a–k** were obtained in excellent diastereoselectivities and good yields.⁷ We found that using THF as a cosolvent in the reaction led to a decrease in diastereoselectivity.⁸ The addition proceeded well not only

(3) For recent reviews on the stereoselective synthesis of piperidines, see: (a) Buffat, M. G. P. *Tetrahedron* **2004**, *60*, 1701–1729. (b) Felpin, F.-X.; Lebreton, J. *Eur. J. Org. Chem.* **2003**, 3693–3712. (c) Weintraub, P. M.; Sabol, J. S.; Kane, J. M.; Borcherding, D. R. *Tetrahedron* **2003**, *59*, 2953–2989.

(4) For selected, recent references on the syntheses of pyridinones by addition of nucleophiles to pyridinium salts and the use of pyridinones in synthesis, see: (a) Comins, D. L.; Sahn, J. *J. Org. Lett.* **2005**, *7*, 5227–5228. (b) Comins, D. L.; King, L. S.; Smith, E. D.; Février, F. C. *Org. Lett.* **2005**, *7*, 5059–5062. (c) Knapp, S.; Yang, C.; Pabbaraja, S.; Rempel, B.; Reid, S.; Withers, S. G. *J. Org. Chem.* **2005**, *70*, 7715–7720. (d) Kuethe, J. T.; Comins, D. L. *J. Org. Chem.* **2004**, *69*, 5219–5231. (e) Ege, M.; Wanner, K. T. *Org. Lett.* **2004**, *6*, 3553–3556. (f) Kuethe, J. T.; Comins, D. L. *J. Org. Chem.* **2004**, *69*, 2863–2866.

(5) (a) Lemire, A.; Charette, A. B. *Org. Lett.* **2005**, *7*, 2747–2750. (b) Lemire, A.; Beaudoin, D.; Grenon, M.; Charette, A. B. *J. Org. Chem.* **2005**, *70*, 2368–2371. (c) Lemire, A.; Grenon, M.; Pourashraf, M.; Charette, A. B. *Org. Lett.* **2004**, *6*, 3517–3520. (d) Charette, A. B.; Grenon, M.; Lemire, A.; Pourashraf, M.; Martel, J. *J. Am. Chem. Soc.* **2001**, *123*, 11829–11830.


(6) Charette, A. B.; Grenon, M. *Can. J. Chem.* **2001**, *79*, 1694–1703.

(7) See Supporting Information for details.

with simple alkylmagnesium reagents such as methylmagnesium bromide (entry 1) but also with bulkier nucleophiles such as *tert*-butylmagnesium bromide (entry 4). Furthermore, the addition of vinyl-, aryl-, and furylmagnesium bromides led to adducts **6g**, **6j**, and **6k**, respectively, with excellent stereocontrol (entries 7, 10, and 11). Only the addition of alkynylmagnesium reagents resulted in a lower selectivity (entries 8 and 9).⁹

The synthesis of (–)-barrenazines A and B started with the introduction of the side chain using an appropriate organomagnesium reagent (Scheme 1). Two approaches were

Scheme 1. Synthesis of α -Iodo Pyridinone **8** and Crystal Structure of **7a**



pursued in parallel starting from **6e** (Table 1, entry 5) and from **6f** (Table 1, entry 6).

The α -iodination of pyridinone **6e** as well as that of ethyl derivative **6b** were efficiently performed using ICl in the presence of Cs₂CO₃ to give **7a** and **7b**, respectively.¹⁰ The generation of the unsaturated side chain of (–)-barrenazine B (**2**) was completed by desilylation using TBAF, Swern oxidation, and Wittig olefination to give **8**.

Alternatively, the terminal olefin moiety of (–)-barrenazine B (**2**) could be directly introduced when using the organomagnesium reagent derived from 7-iodohept-1-ene. Thus, the corresponding pyridinone **6f** was obtained in high diastereoselectivity (Table 1, entry 6). The chemoselective

(8) The organomagnesium reagents were either purchased as solutions in Et₂O or freshly prepared prior to use from the corresponding alkyl iodides, vinyl, or phenyl bromides using *t*-BuLi or by deprotonation using *n*-BuLi, followed by transmetalation to MgBr₂·OEt₂. See Supporting Information for details.


(9) THF had to be added in these reactions to solubilize the organomagnesium reagents.

(10) ICl for iodination of pyridinones: (a) Comins, D. L.; Hiebel, A.-C.; Huang, S. *Org. Lett.* **2001**, *3*, 769–771. (b) See also: Comins, D. L.; Joseph, S. P.; Chen, X. *Tetrahedron Lett.* **1995**, *36*, 9141–9144.

conversion of **6f** to **8** was then accomplished using I₂ in the presence of pyridine.¹¹

α -Iodo, α,β -unsaturated carbonyl compounds are valuable intermediates in organic synthesis, especially in transition-metal-mediated reactions.¹² For the purpose of the synthesis of **1** and **2**, α -iodo **7a** and **8** were transformed into the Boc-amino enones **9a,b** via a Buchwald's Cu-catalyzed C–N cross-coupling using *tert*-butyl carbamate¹³ (Scheme 2).

Scheme 2. Synthesis of Protected Barrenazine A and B



Originally, we intended to use **9** directly in a reductive cyclization to construct the pyrazine core of barrenazine. However, the α -amino enone obtained from **9a** upon TFA deprotection did not undergo the desired reaction using Pd/C and hydrogen gas.¹⁴ Presumably, the enone system is too deactivated because of conjugation of the free amino group with the amidine moiety. It was anticipated that reduction

(11) I₂/pyridine reagent: (a) Krafft, M. E.; Cran, J. W. *Synlett* **2005**, 1263–1266. (b) Souza, F. E. S.; Sutherland, H. S.; Carlini, R.; Rodrigo, R. *J. Org. Chem.* **2002**, *67*, 6568–6570. (c) Johnson, C. R.; Adam, J. P.; Braun, M. P.; Senanayake, C. B. W.; Wovkulich, P. M.; Uskokovic, M. R. *Tetrahedron Lett.* **1992**, *33*, 917–918.

(12) For review of β -iodo carbonyl compounds in Pd-catalyzed cross-coupling reactions, see: Negishi, E. *J. Organomet. Chem.* **1999**, *576*, 179–194.

(13) Jiang, L.; Job, G. E.; Klapars, A.; Buchwald, S. L. *Org. Lett.* **2003**, *5*, 3667–3669.

(14) For cyclizations of β -amino cyclohexenones, see: (a) Drögemüller, M.; Flessner, T.; Jautelat, R.; Scholz, U.; Winterfeldt, E. *Eur. J. Org. Chem.* **1998**, *2811*, 1–2831. (b) Kramer, A.; Ullmann, U.; Winterfeldt, E. *J. Chem. Soc., Perkin Trans. 1* **1993**, 2865–2867.

of the Boc derivatives **9** to the saturated piperidinone **10** would circumvent this problem. The construction of symmetrical pyrazines from α -amino aldehydes and ketones is well-known, and condensation usually proceeds fairly easily.¹⁵ The 1,4-reduction of **9** using known procedures such as L-selectride¹⁶ or Stryker's reagent proved to be difficult.¹⁷ Poor selectivity of 1,4- vs 1,2-reduction and/or low yields of the corresponding piperidinones were observed. The only reagent that gave complete 1,4-selectivity and full conversion was a combination of CuBr and LiAl(O^{*t*}Bu)₃H.¹⁸ Thus, we could obtain compounds **10a** and **10b** in excellent yields and good diastereoselectivities. Although having diastereomerically enriched **10** is meaningless for the remainder of the synthesis, as the stereogenic center is lost in the subsequent step, the Boc-amino ketones **10** are interesting chiral building blocks.

The configuration of piperidinone **10a** was determined by analyzing the coupling constants and by NOESY experiments. Nitrogen heterocycles protected with acyl, carbamate, or imidate groups are known to place substituents at the 2-position axially, to minimize A^{1,3} strain (Figure 2).¹⁹ NMR




Figure 2. Configuration of *N*-Boc-amino ketone **10**.

analysis indicated that diastereomer **10** was the major product arising from the reduction (Figure 2).

Having Boc-amino ketones **10** in hand, the next steps consisted of the amine deprotection with TFA followed by the oxidative cyclization under basic conditions in air to afford **11**.

The amidine auxiliary was then removed via an intramolecular, alkoxy-assisted strategy using BBr₃ (Scheme 3).

(15) Selected examples: (a) Tonsiengsom, F.; Miyake, F. Y.; Yakushijin, K.; Horne, D. A. *Synthesis* **2005**, in print. (b) Gosh, U.; Ganessunker, D.; Sattigeri, V. J.; Carlson, K. E.; Mortensen, D. J.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. *Bioorg. Med. Chem.* **2003**, *11*, 629–657. (c) Chiba, T.; Sakagami, H.; Murata, M.; Okimoto, M. *J. Org. Chem.* **1995**, *60*, 6764–6770.


(16) Selected examples: (a) Alcaide, B.; Almendros, P.; Alonso, J. M.; Aly, M. F. *Chem.–Eur. J.* **2003**, *9*, 3415–3426. (b) Tasber, E. S.; Garbaccio, R. M. *Tetrahedron Lett.* **2003**, *44*, 9185–9188. (c) Lim, S. H.; Curtis, M. D.; Beak, P. *Org. Lett.* **2001**, *3*, 711–714. (d) Comins, D. L.; Libby, A. H.; Al-awar, R. S.; Foti, C. J. *J. Org. Chem.* **1999**, *64*, 2184–2185.

(17) Selected examples: (a) Lipshutz, B. H.; Chrisman, W.; Noson, K.; Papa, P.; Sclafani, J. P.; Vivian, R. W.; Keith, J. M. *Tetrahedron* **2000**, *56*, 2779–2788. (b) Lipshutz, B. H.; Keith, J.; Papa, P.; Vivian, R. *Tetrahedron Lett.* **1998**, *39*, 4627–4630. (c) Mahoney, W. S.; Brestensky, D. M.; Stryker, J. M. *J. Am. Chem. Soc.* **1988**, *110*, 291–293.

(18) 1,4-Reduction of enones with CuX/LiAlH: (a) Comins, D. L.; LaMunyon, D. H. *Tetrahedron Lett.* **1989**, *30*, 5053–5056. (b) Comins, D. L.; Abdullah, A. H. *J. Org. Chem.* **1984**, *49*, 3392–3394. (c) Semmelhack, M. F.; Stauffer, R. D.; Yamashita, A. *J. Org. Chem.* **1977**, *42*, 3180–3188.

(19) (a) Watson, P. S.; Jiang, B.; Scott, B. *Org. Lett.* **2000**, *2*, 3679–3681. (b) Hoffmann, H. R. *Chem. Rev.* **1989**, *89*, 1841–1860. (c) Johnson, F. *Chem. Rev.* **1968**, *68*, 375–413. See also the X-ray crystal structures in the Supporting Information.

Scheme 3. Amidine Group Cleavage and Isolation of Barrenazine A (**15a**) and B (**15b**) Bis(trifluoroacetate)^a



^a TFA salts were obtained after HPLC purification; see Supporting Information for details.


After cleavage of the methoxy groups on the amidine moiety of **11a**, **12** spontaneously fragmented to afford oxazoline **14** and free hexahydrodipyrindopyrazine, which was isolated as its Boc derivative **13**. The same procedure applied to pyrazine **11b**, obtained after hydrogenation of the terminal alkenes, produced barrenzine A bis(trifluoroacetate) (**15a**) in 55% yield.

The amidine cleavage with the unsaturated side chain still in place proved to be more challenging. A mixture with varying amounts of brominated side chains was observed if the same reaction conditions as before were used. Fortunately, a combination of BBr₃ and 2,6-lutidine turned out to cleave the methoxy groups efficiently while leaving the terminal olefin untouched. Thus, barrenzine B (**15b**) was isolated as its bis(trifluoroacetate) salt in 51% yield.²⁰ The spectroscopic data for both **15a** and **15b** were identical to those reported by Kashman.¹ Neutralization upon treatment with saturated aqueous NaHCO₃ led to barrenzine A and B.

(20) The ratio of barrenzine/TFA was determined by NMR using α,α,α -trifluorotoluene as the internal standard.

Pyrazine **11b** is an interesting compound to apply ring-closing metathesis conditions.²¹ Applying Grubbs' second generation catalyst, followed by hydrogenation of the double bond and removal of the chiral auxiliary, allowed the isolation of the cyclophane-type macrocycle **16** in 51% yield (Scheme 4).²²

Scheme 4. RCM of **11b** to Give Cyclophane **16**^a



^a TFA salt was obtained after HPLC purification; see Supporting Information for details.

In summary, an expedient synthesis of (–)-barrenzine A and B was accomplished in 30% (eight steps) and 28% (seven steps) overall yield, respectively, from 4-methoxypyridine. The synthetic route allows for the preparation of various analogues as exemplified by the synthesis of the ethyl side chain analogue.

Acknowledgment. This work was supported by the National Science and Engineering Research of Canada (NSERC), Merck Frosst Canada & Co., Boehringer Ingelheim (Canada), Ltd., and the Université de Montréal. We are also grateful to Professor Yoel Kashman from Tel Aviv University for providing the spectra of the isolated natural products. We also thank Dr. M. Sales for helpful discussions.

Supporting Information Available: Experimental procedures, characterization data, and NMR spectra for all new compounds and crystal structure data for **6i** and **7a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0609006

(21) For reviews on the olefin metathesis reaction, see: (a) Deiters, A.; Martin, S. F. *Chem. Rev.* **2004**, *104*, 2199–2238. (b) Grubbs, R. H. *Tetrahedron* **2004**, *60*, 7117–7140. (c) Connor, S. J.; Blechert, S. *Angew. Chem., Int. Ed.* **2003**, *42*, 1900–1923. (d) Trnka, T. M.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, *34*, 18–29. (e) Fürstner, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 3012–3043.

(22) A mixture of *E/Z* double-bond isomers in a ratio of 8:1 is isolated, if the hydrogenation step is avoided.