

A New Entry to Polycyclic Indole Skeletons via Palladium-Catalyzed Intramolecular Heteroannulation


Jianxiu Liu, Ming Shen, Yongda Zhang, Guisheng Li, Ahmad Khodabocus, Sonia Rodriguez, Bo Qu, Vittorio Farina,[†] Chris H. Senanayake,[†] and Bruce Z. Lu^{*}

Department of Chemical Development, Boehringer-Ingelheim Pharmaceuticals, Inc.,
800 East Leigh Street, Suite 205, Richmond, Virginia 23219

zlu@rdg.boehringer-ingelheim.com

Received June 12, 2006

ABSTRACT




A one-step method was developed for elaboration of a variety of polycyclic indole skeletons via a novel palladium-catalyzed intramolecular indolization of 2-chloroanilines bearing tethered acetylenes. This novel intramolecular indolization method unveils an unusual *syn* amidopalladation pathway of a tethered alkyne.

Alkaloids containing the indole structure motif are found in numerous natural products and synthetic compounds of vital medicinal value.¹ Construction of polycyclic indoles usually requires multistep approaches.² The exploration of new methodologies that allow rapid establishment of these indole skeletons in a single operation remains an important challenge facing organic chemists.³

Recently we have developed a novel Pd-catalyzed regioselective intermolecular indolization of 2-bromo- or chloroanilines with internal alkynes,⁴ based on Larock's protocol.⁵ The feasibility of using chloroaniline derivatives as the

starting material has significant practical and economical value from a cost and throughput standpoint. We were curious whether the same Pd-catalyzed alkyne insertion protocol could be applied for the preparation of polycyclic indoles via an intramolecular indolization process (Scheme 1). Herein we report our first results on this program.

Scheme 1. Synthesis of Polycyclic Indoles via Pd-Catalyzed Intramolecular Heteroannulation



Mechanistically, to achieve the successful execution of this catalytic process, one prerequisite is to form the desired

(5) (a) Larock, R. C.; Yum, E. K. *J. Am. Chem. Soc.* **1991**, *113*, 6689. (b) Larock, R. C.; Yum, E. K.; Refvik, M. D. *J. Org. Chem.* **1998**, *63*, 7652.

[†] Current address: Boehringer-Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, Ridgefield, CT 06877.

(1) For recent reviews of indole-containing natural products, see: (a) Lounasmaa, M.; Tolvanen, A. *Nat. Prod. Rep.* **2000**, *17*, 175. (b) Faulkner, D. J. *Nat. Prod. Rep.* **1999**, *16*, 155. (c) Michael, J. P. *Nat. Prod. Rep.* **1998**, *15*, 571.


(2) (a) Vice, S. F.; Copeland, C. R.; Forsey, S. P.; Dmitrienko, G. I. *Tetrahedron Lett.* **1985**, *26*, 165. (b) Vice, S. F.; Friesen, R. W.; Dmitrienko, G. I. *Tetrahedron Lett.* **1985**, *26*, 5253. (c) Artis, D. R.; Cho, I.-S.; Jaime-Figueroa, S.; Muchowski, J. M. *J. Org. Chem.* **1994**, *59*, 2456.

(3) Hiroi, K.; Hiratsuka, Y.; Watanabe, K.; Abe, I.; Kato, F.; Hiroi, M. *Synlett* **2001**, 263.

(4) Shen, M.; Li, G.; Lu, B. Z.; Hossain, A.; Roschangar, F.; Farina, F.; Senanayake, C. H. *Org. Lett.* **2004**, *6*, 4129.

amidopalladacycle **5** via the “*endo-dig*” carbopalladation, which would afford the indoles by subsequent C–N reductive elimination (path C).⁵ Given the strained geometry of amidopalladacycle **5**, as well as the potential competing “*exo-dig*”-carbopalladation of **3** (path B),⁶ we expected that the reaction may work only for large rings (i.e., $n =$ a rather large number). Surprisingly, we have observed that the desired product is formed even when n (Schemes 1 and 2)

Scheme 2. Mechanism Insight into Pd-Catalyzed Intramolecular Heteroannulation



is very small ($n = 1$). This observation points, therefore, to a new and unanticipated mechanism being responsible for this novel transformation. In this preliminary Letter, we disclose a novel domino process that involves, most likely, the following steps: (a) formation of the Pd-containing zwitterion **6** (Scheme 2, path D) by standard oxidative addition of Pd(0) to **1**, probably through depot form **2**;⁷ (b) formation of the bicyclic palladacycle **7** by *syn* amidopalladation of the acetylene into **6**;^{8–10} and (c) reductive elimination of the palladacycle **7** to afford the indole ring.

(6) (a) Zhang, Y.; Negishi, E.-I. *J. Am. Chem. Soc.* **1989**, *111*, 3454. (b) Brown, D.; Grigg, R.; Sridharan, V.; Tambyrajah, V.; Thornton-Pett, M. *Tetrahedron* **1998**, *54*, 2595. (c) Wang, R.-T.; Chou, F.-L.; Luo, F.-T. *J. Org. Chem.* **1990**, *55*, 4846 and references therein. (d) Luo, F.-T.; Wang, R.-T. *Heterocycles* **1991**, *32*, 2365. (e) Hayashi, M.; Sai, H.; Horikawa, H. *Heterocycles* **1998**, *48*, 1331. (f) Zhang, H.; Larock, R. C. *Org. Lett.* **2002**, *4*, 3035. (g) Brase, S.; Meijere, A. *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; John Wiley & Sons: New York, 2002; Vol 1, pp 1369 and 1405. (h) Bolton, G. L.; Hodges, J. C., *J. Comb. Chem.* **1999**, *1*, 130.

(7) Palladation of anilides yields six-membered Pd cycles such as **2**, see: Horino, H.; Inoue, N. *J. Org. Chem.* **1981**, *46*, 4416.


(8) (a) Ney, J. E.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 8644. (b) Hay, M. B.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 16468. (c) Nakhla, J. S.; Kampf, J. W.; Wolfe, J. P. *J. Am. Chem. Soc.* **2006**, *128*, 2893. (d) Brice, J. L.; Harang, J. E.; Timokhin, V. I.; Anastasi, N. R.; Stahl, S. S. *J. Am. Chem. Soc.* **2005**, *127*, 2868.

(9) For *anti* addition in aminopalladation, see: Cacchi, S.; Marinelli, F. *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; John Wiley & Sons: New York, 2002; Vol. 2, pp 2227–2244.

To the best of our knowledge, intramolecular *syn* amidopalladation of this type has never been reported in the literature.

We started the investigation with the 2-chloroaniline derivative **9** by evaluating the influence of several important reaction parameters. On the basis of our previous experience on *intermolecular* indolization, DtBPF or Cy₃P is superior to the other conventional electron-rich ligands for the alkyne insertion to the arylpalladium chloride intermediates. However, we envisioned that the choice of the optimum base would be very crucial in order to overcome the “*exo-dig*” carbopalladation pathway (to yield **4**, path B, Scheme 2). Thus, our initial efforts were mainly focused on the investigation of base effects. These results are summarized in Table 1.

Table 1. Conditions Screened for the Intramolecular Heteroannulation of **9**^a

















entry	ligand	base	solvent	temp/time (°C/ min)	solution assay (%)
1	DtBPF	DBU	NMP	130/120	trace
2	DtBPF	TMG	NMP	130/120	trace
3	DtBPF	KHCO ₃	NMP	130/90	25
4	DtBPF	K ₂ CO ₃	NMP	130/40	70
5	DtBPF	Na ₂ CO ₃	NMP	130/90	11
6	DtBPF	K ₃ PO ₄	NMP	130/60	50
7 ^b	DtBPF	K ₂ CO ₃	NMP	130/150	49
8	DtBPF	K ₂ CO ₃	NMP	110/240	41
9	DtBPF	CS ₂ CO ₃	NMP	130/90	26
10	DtBPF	NaH	NMP	130/20	trace
11	CTC-Q-Phos	K ₂ CO ₃	NMP	130 /180	trace
12	<i>t</i> Bu ₃ P	K ₂ CO ₃	NMP	130/40	37
13	DiPPF	K ₂ CO ₃	NMP	130/40	35

^a All reactions were carried out with anilide **9** (1.0 mmol), Pd(OAc)₂ (5 mol %), DtBPF (10 mol %), and K₂CO₃ (5 equiv) in NMP (10 mL).
^b Pd₂(dba)₃ (2.5 mol %) was used.

As anticipated, the optimal results were obtained by using DtBPF as ligand. K₂CO₃ was the optimal base and NMP proved to be the superior solvent. The reaction of **9** was complete within 1 h at 130 °C under these conditions, which is much faster than the intermolecular reaction under the same conditions,⁴ providing the desired product **10** in 70% isolated yield (entry 4, Table 1).

(10) For related chemistry that may have similar mechanism, see: (a) Maassarani, F.; Pfeffer, M.; Spencer, J.; Wehman, E. *J. Organomet. Chem.* **1994**, *466*, 265. (b) Korivi, R. P.; Cheng, C. H. *Org. Lett.* **2005**, *7*, 5179.

Table 2. Synthesis of Polycyclic Indoles via Pd-Catalyzed Intramolecular Heteroannulation of Tethered Alkynes with 2-Chloroanilides^a

entry	substrate	product	yield(%) ^b
1			56
2			70
3			61
4			59
5			78
6			50
7			75

^a All reactions were carried out with 2-chloroanilides (1.0 mmol), Pd(OAc)₂ (5 mol %), D'BPF (10 mol %), and K₂CO₃ (2.5 equiv) in NMP (10 mL). ^b Isolated yield of major product.

The dependence on the base used is rather remarkable, as organic bases are totally ineffective (entries 1 and 2) and other inorganic bases are also quite inefficient. To illustrate

the scope of this methodology, a variety of polycyclic indoles were synthesized with our standard conditions, even though better yield may be obtainable through re-optimization of each individual entry. These results are summarized in Table 2. (2-Chloroaryl)carbamate **11** and ureas **17**, **19**, and **21** are readily prepared from (2-chlorobenzene)isocyanate with alkynyl alcohols or amines. Reaction of the carbamate **11** is very fast. However, reduced yields were obtained due to the decomposition of the starting material under the reaction conditions, presumably through the corresponding isocyanate (Table 2, entry 1). The influence of substituents on both ends of the acetylene and the ring size are illustrated by the amides **9**, **13**, and **15** (Table 2, entries 2–4). Five-, six-, and seven-membered rings were formed uneventfully. The major side product frequently observed is the dechlorinated starting material, the result of a reductive process. Work toward identifying the source of reducing agent is underway as are further studies on the optimization of these processes.

At this moment, we have no mechanistic understanding of this process, but we believe that the rationale presented in Scheme 2 adequately explains our observations and is sufficiently preceded in the *aminopalladation of alkenes*.

In summary, we have developed a one-step method for elaboration of a variety of polycyclic indole skeletons via a novel palladium-catalyzed intramolecular indolization of 2-chloroanilines bearing tethered acetylenes. This novel intramolecular indolization method unveils an unusual *syn* amidopalladation pathway of a tethered alkyne. Further studies on this protocol and its scope are in progress.

Acknowledgment. We acknowledge Dr. Heewon Lee of Boehringer-Ingelheim Pharmaceuticals for analytical support.

Supporting Information Available: Reaction procedures and characterization of products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL061440J