

# A New Approach to the Neoglycopeptides: Synthesis of Urea- and Carbamate-Tethered *N*-Acetyl-*D*-glucosamine Amino Acid Conjugates


Yoshiyasu Ichikawa,\* Fumiyo Ohara, Hiyoshizo Kotsuki, and Keiji Nakano

Faculty of Science, Kochi University, Akebono-cho, Kochi 780-8520, Japan

ichikawa@kochi-u.ac.jp

Received July 8, 2006

## ABSTRACT



A novel approach to the synthesis of Fmoc-protected neoglycopeptide building blocks is described. Oxidation of *N*-acetyl-*D*-glucosamine isonitrile afforded the corresponding highly reactive glycopyranosyl isocyanate, which reacted with amino acid derivatives to furnish the corresponding urea- and carbamate-tethered Fmoc-protected *N*-acetyl-*D*-glucosamine amino acid conjugates in good yields.

Carbohydrates linked to the peptide backbone of proteins have become the focus of bioorganic and/or medicinal research work due to their involvement in diverse biochemical processes such as cellular recognition and adhesion.<sup>1</sup> Carbohydrates in natural glycoproteins are attached to the peptide backbones through the oxygen in the side chain of serine or threonine in *O*-linked glycoproteins or through the carboxamide nitrogen of asparagine in *N*-linked glycoproteins.<sup>2</sup> Although extensive research effort has been devoted to exploring the synthesis of accurately sequenced glycopeptides for biological and structural studies, total synthesis of the native glycopeptides still remains a challenging and time-consuming endeavor due to the difficulty of glycosylation reactions. In parallel, development of glycopeptide mimetics continues to attract much attention in order to supply homogeneous, stable, and readily accessible glycopeptide analogues for biological studies and therapeutic applications.<sup>3</sup> Accordingly, the covalent attachment of car-

bohydrates to peptides replaced with non-native linkages is of particular interest, with special attention having been paid to mimetics where the *O*-glycosidic linkage is replaced by carbon–carbon,<sup>4</sup> carbon–sulfur,<sup>5</sup> and carbon–aminoxy units.<sup>6</sup> For glycopeptide mimetics of natural glycopeptides with *N*-glycosidic linkages, the amide group in *N*-glycosides has been replaced by a retroamide subunit<sup>7</sup> and a urea glycosyl bond.<sup>8</sup>

During the course of developing a synthesis of various glycopeptide mimetics, we planned to pursue the solid-phase

(3) (a) Taylor, C. M. *Tetrahedron* **1998**, *54*, 11317. (b) Davis, B. G. *Chem. Rev.* **2002**, *102*, 579.

(4) (a) Burkhardt, F.; Hoffmann, H.; Kessler, H. *Angew. Chem., Int. Ed.* **1997**, *36*, 1191. (b) Dondoni, A.; Mara, A. *Chem. Rev.* **2000**, *100*, 4395. (c) McGravey, G. J.; Benedum, T. E.; Schmidmann, F. W. *Org. Lett.* **2002**, *4*, 3591.

(5) (a) Davis, N. J.; Flitsch, S. L. *Tetrahedron Lett.* **1991**, *32*, 6793. (b) Jobron, L.; Hummel, G. *Org. Lett.* **2000**, *2*, 2265. (c) Knapp, S.; Myers, D. S. *J. Org. Chem.* **2001**, *66*, 3636. (d) Cohen, S. B.; Halcomb, R. L. *Org. Lett.* **2001**, *3*, 405. (e) Zhu, X.; Pachamuthu, K.; Schmidt, R. R. *J. Org. Chem.* **2003**, *68*, 5641 and references therein.

(6) Rodriguez, E. C.; Marcaurelle, L. A.; Bertozzi, C. R. *J. Org. Chem.* **1998**, *63*, 7134.


(7) Hoffmann, M.; Burkhardt, F.; Hessler, G.; Kessler, H. *Helv. Chim. Acta* **1996**, *79*, 1519.

(1) (a) Dwek, R. A. *Chem. Rev.* **1996**, *96*, 683. (b) Varki, A. *Glycobiology* **1993**, *3*, 97.

(2) (a) Marcaurelle, L. A.; Bertozzi, C. R. *Chem.—Eur. J.* **1999**, *5*, 1384. (b) H. Kunz. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 294.

synthesis of urea-tethered neoglycopeptides, the strategy for which is outlined in Scheme 1. Starting with urea-linked


**Scheme 1.** Synthetic Strategy of Neoglycopeptides



*N*-acetyl-D-glucosamine amino acid conjugate **A**, solid-phase synthesis would give the *N*-acetyl-D-glucosaminyl peptide **B**. Transglycosylation of **B** using endo- $\beta$ -GlcNAc-ase would provide the urea-tethered neoglycopeptide **C**.<sup>9</sup> In this paper, we report on the synthesis of the urea-tethered glycosyl amino acid (**A**, P<sup>1</sup> = Fmoc) and its use as a building block for solid-phase synthesis based on the Fmoc-strategy.


Our retrosynthetic analysis of **A** is shown in Scheme 2. A key feature of our plan was to construct the urea-glycosyl bond through a coupling reaction between isocyanate **1** and various  $\alpha,\beta$ -diamino acid derivatives. Since such a highly functionalized isocyanate **1** was only postulated to be a transient intermediate by Pinter,<sup>10</sup> the crucial step in our approach was the synthesis of the reactive isocyanate **1**. To solve this problem, we planned to employ the oxidation of isonitrile **2** for the generation of **1** under mild reaction conditions.<sup>11</sup>

**Scheme 2.** Retrosynthetic Analysis of **A**



Starting with commercially available *N*-acetyl-D-glucosamine **3** (Scheme 3), Horton's protocol afforded  $\alpha$ -chloro-*N*-acetyl-D-glucosamine acetate **4**,<sup>12</sup> which was further transformed into glycosyl azide **5** by the displacement reaction with sodium azide under phase-transfer conditions.<sup>13</sup> Catalytic hydrogenation of azide **5** followed by treatment of the resulting glycopyranosylamine with acetic formic anhydride furnished the formamide **6** in 73% yield over two steps. Dehydration of **6** with triphosgene/triethylamine gave the *N*-acetyl-D-glucosamine isonitrile **2** in good yield (81%).

**Scheme 3.** Preparation of *N*-Acetyl-D-glucosamine Isonitrile **2**




With the synthesis of isonitrile **2** established, we initially examined the synthesis of urea glycosides by the reaction of **1** with six different amines as summarized in Table 1. In a typical case (entry A), oxidation of **2** was carried out with pyridine *N*-oxide (3 equiv) and a catalytic amount of iodine (7 mol %) in acetonitrile in the presence of water scavenger (MS 3 Å).

The resulting solution, containing highly reactive isocyanate **1**, was immediately treated with phenethylamine (2.0 equiv).<sup>14</sup> To our delight, the urea glycoside **7a** was isolated in 92% yield. Primary amines having alkyl branches at the  $\alpha$ -carbon (entry B and C) as well as secondary amine (entry D) smoothly reacted with **1** to afford the corresponding urea glycosides **7b-d** in good yields (>90%). Even in the case of a sterically hindered secondary amine, such as diisopropylamine (entry E), the corresponding urea glycoside **7e** was obtained in 91% yield. An application of **1** for the synthesis of urea-tethered disaccharide is represented in entry F, where aminosugar reacted with **1** to yield the urea-tethered pseudo-disaccharide **7f** in 92% yield.

(8) Ichikawa, Y.; Nishiyama, T.; Isobe, M. *Synlett* **2000**, 1253.

(9) (a) Seitz, O.; Wong, C.-H. *J. Am. Chem. Soc.* **1997**, *119*, 8766. (b) Wang, L.-X.; Tang, M.; Suzuki, T.; Kitajima, K.; Inoue, Y.; Inoue, S.; Fan, J.-Q.; Lee, Y. *J. Am. Chem. Soc.* **1997**, *119*, 11137. (c) Mizuno, M.; Haneda, K.; Iguchi, R.; Muramoto, I.; Kawakami, T.; Aimoto, S.; Yamamoto, K.; Inazu, T. *J. Am. Chem. Soc.* **1999**, *121*, 284.

(10) Pinter reported reaction of azide **5** with triphenylphosphine and CO<sub>2</sub> and isolated the symmetrical carbodiimide **ii**. Although the intermediacy of isocyanate **1**, formed by reaction of iminophosphorane **i** with CO<sub>2</sub>, was proposed, isolation or even detection of **1** has never been successful due to the rapid reaction of **i** with reactive isocyanate **1**. See: Kovacs, J.; Pinter, I.; Messmer, A. *Carbohydr. Res.* **1987**, *166*, 101.



(11) Ichikawa, Y.; Nishiyama, T.; Isobe, M. *J. Org. Chem.* **2001**, *66*, 4200.

(12) Horton, D.; Johnson, A. L.; McKusick, B. C. *Organic Syntheses*; Wiley: New York, 1973; Collect. Vol. V, p 1.


(13) Tropper, F. D.; Anderson, F. O.; Braun, S.; Roy, R. *Synthesis* **1992**, 619.

(14) Although silica gel TLC analysis of the reaction mixture showed the consumption of **2**, isocyanate **1** could not be observed by TLC. Accordingly, we employed 2 equiv of amines to optimize the yields.

**Table 1.** Synthesis of Urea Glycosides from Isonitrile **2**

entry	product (R=)	yield (%)
A		92
B		92
C		93
D		91
E		91
F		92


Encouraged by the ready formation of urea glycosides by the reaction of **1** with amines, we next explored the synthesis of an Fmoc-protected *N*-acetyl-D-glucosamine amino acid conjugate **13** (Scheme 4). Boc-protection of  $\alpha,\beta$ -diamino acid

**Scheme 4.** Synthesis of Urea-Linked *N*-Acetyl-D-glucosamine Amino Acid Conjugate **13**

derivative **8**<sup>15</sup> followed by a deprotection–protection sequence gave the Fmoc-protected amino acid **10**.


Esterification of **10** furnished the fully protected  $\alpha,\beta$ -diamino acid **11**. Removal of the *N*-Boc protecting group in **11** was carried out by treatment with TFA in  $\text{CH}_2\text{Cl}_2$ , and the resultant TFA salt **12** was treated with isocyanate **1** (2.0 equiv) in the presence of diisopropylethylamine at room temperature for 60 min. Fmoc-protected *N*-acetyl-D-glucosamine amino acid conjugate **13** was isolated in 85% yield after chromatographic purification.


During the course of developing our urea-tethered glycosyl amino acid conjugate, it was envisioned that the highly reactive isocyanate **1** would react with serine derivatives to afford a new type of carbamate-tethered *N*-acetyl-D-glucosamine amino acid conjugates. Accordingly, the reaction of **1** with benzyl alcohol was quickly investigated to check the reactivity of **1** toward alcohols (Scheme 5). To our

**Scheme 5.** Reaction of Isocyanate **1** with Benzyl Alcohol

delight, treatment of **1** with benzyl alcohol (1.0 equiv) at room temperature for 5 h gave rise to the carbamate **14**, albeit in low yield (20%). However, use of excess benzyl alcohol (3.0 equiv) and heating the reaction mixture at 70 °C improved the yields up to 82%.

Our attention then turned toward the synthesis of carbamate-tethered *N*-acetyl-D-glucosamine amino acid conjugate **16** (Table 2). Treatment of isocyanate **1** with serine derivative **15** using reaction conditions similar to those employed in Scheme 5 (entries A–C) furnished the coupling product **16** in moderate yields (36–48%). Although a large excess of **15** (9 molar excess with respect to **1**) improved the yield up to 80% and excess **15** was recovered by chromatography (entry D), we attempted to improve the reaction by activating the isocyanate group in **1**. Although activation of **1** with 4-(*N,N*-dimethylamino)pyridine (DMAP) gave the dimeric product predominantly,<sup>16</sup> use of 20 mol % dibutyltin dilaurate successfully improved the coupling process in good yield (81%, entry F).<sup>17</sup> It should be noted that the reaction conditions in entry F are mild and neutral and that  $\beta$ -elimination and deprotection of acetyl/Fmoc groups was not observed.

(15) Waki, M.; Kitajima, Y.; Izumiya, N. *Synthesis* **1981**, 266.(16) Initial attempts to activate isocyanate **1** by DMAP resulted in the intramolecular attack of nitrogen in the *N*-acetyl group to the neighboring isocyanate (**1** → **i**) followed by dimerization to afford **ii**.(17) (a) Abbate, F. W.; Ulrich, H. *J. Appl. Polym. Sci.* **1969**, *13*, 1929. (b) Greenwald, R. B.; Pendri, A.; Bolikal, D. *J. Org. Chem.* **1995**, *60*, 331.

**Table 2.** Optimization for the Synthesis of **16**

entry	isonitrile <b>2</b> (equiv)	serine <b>15</b> (equiv)	<i>T</i> (°C)	time (h)	catalyst (mol %)	yield (%)
A	1.0	3.0	rt	5	none	40
B	1.0	3.0	70	5	none	48
C	3.0	1.0	rt	5	none	36
D	1.0	10.0	70	15	none	80
E	1.0	3.0	70	12	20	64
F	3.0	1.0	rt	7	20	81

We have demonstrated that *N*-acetyl-D-glucopyranosyl isocyanate **1** is a valuable synthon for the preparation of urea- and carbamate-tethered amino acid conjugates. The highly reactive nature of isocyanate **1**, as demonstrated in the

(18) For the reactivity of glycosyl isothiocyanates, see: Witczak, Z. J. *Adv. Carbohydr. Chem. Biochem.* **1986**, *44*, 91. For recent representative examples of thiourea conjugation strategy, see: (a) Lindhorst, T. K.; Kieburg, C. *Angew. Chem., Int. Ed.* **1996**, *35*, 1953. (b) Benito, J. M.; Gomez-Garcia, M.; Ortiz Mellet, C.; Baussanne, I.; Defaye, J.; Garcia Fernandez, J. M. *J. Am. Chem. Soc.* **2004**, *126*, 10355.

coupling reaction with the hydroxy group of the serine derivative **15** under mild conditions, is noteworthy when compared with the less reactive glycosyl isothiocyanate counterpart.<sup>18</sup> Further studies to the synthesis of neoglycopeptides using **13** and **16** is now in progress.<sup>19</sup>

**Acknowledgment.** We are grateful for financial support from a Grant-in-Aid for Scientific Research on Priority Areas (18032055 and 18037053) from MEXT. This work was supported in part by a Special Research Grant for Green Science from Kochi University.

**Supporting Information Available:** Full experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0616788

(19) In the synthesis of *O*-linked type glycopeptides, removal of the Fmoc group became a serious problem due to the base-catalyzed  $\beta$ -elimination (Kunz, H. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 294). To check this point, Fmoc-deprotection of **13** and **16** were carried out by treatment with 20% piperidine in DMF at room temperature for 10 min. In these reactions,  $\beta$ -elimination has never been observed, and the products and yields were confirmed after acetylation, which shows the compatibility of both **13** and **16** with standard solid-phase peptide synthesis.

