

Design and Synthesis of Novel C_2 -Symmetric Chiral Piperazines and an Application to Asymmetric Acylation of σ -Symmetric 1,2-Diols


Daisuke Nakamura,[‡] Kiyomi Kakiuchi,[‡] Kenji Koga,[‡] and Ryuichi Shirai^{*,‡,†}

Research and Education Center for Materials Science and Graduate School of Materials Science, Nara Institute of Science and Technology, Takayama, Ikoma-shi, Nara 630-0192, Japan, and Faculty of Pharmaceutical Sciences, Doshisha Women's College of Liberal Arts, Kodo, Kyotanabe-shi, Kyoto 610-0395, Japan

rshirai@dwc.doshisha.ac.jp

Received October 27, 2006

ABSTRACT



A novel alicyclic chiral C_2 -symmetric piperazine, (*S,S*)-**7**, is designed and synthesized from L-proline. Benzoylation of a series of cyclic and acyclic *meso*-1,2-diols with a catalytic amount of (*S,S*)-**7** and CuCl_2 provided optically active monobenzoates with high enantioselectivity.

Designing of new chiral ligands to improve enantioselectivity of organic reactions has become an important issue. Chiral tertiarily diamines (**1**–**5**) have been used as efficient chiral inductors, particularly (–)-sparteine (**1**) and bisoxazoline (**3**) have been found to be effective in many asymmetric transformations.¹ Our project has focused on the development of a new class of C_2 -symmetric alicyclic chiral diamines. In this article, we describe the design and the synthesis of a novel C_2 -symmetric chiral alicyclic tertiary diamine, (5*aS*,10*aS*)-5*a*,10*a*-dimethyldecahydrodipyrrolo[1,2-*a*:1',2'-*d*]pyrazine ((*S,S*)-**7**, DMPP), and its application to asymmetric desymmetrization of *meso*-1,2-diols by enantioselective monoacylation.

Prior to the synthesis of (*S,S*)-**7**, we reinvestigated the synthesis of (5*aS*,10*aS*)-decahydrodipyrrolo[1,2-*a*:1',2'-*d*]pyrazine ((*S,S*)-**6**, DHPP), which was reported by Breitmeier and co-worker,² by intramolecular aminolysis of L-proline

methyl ester at high temperature followed by LiAlH_4 reduction. Unfortunately, their experimental procedure was not reproducible and the product was obtained in low yield and contained inseparable unidentified impurity. In addition, their reported ^{13}C NMR spectra indicates ten ^{13}C -signals for the structure of (*S,S*)-**6** even though it has C_2 -symmetry,

- (1) (a) Nozaki, H.; Aratani, T.; Toraya, T. *Tetrahedron Lett.* **1968**, 4097. (b) Mukaiyama, T.; Asami, S. *Top. Curr. Chem.* **1985**, 127, 133. (c) Tomioka, K.; Nakajima, M.; Koga, K. *J. Am. Chem. Soc.* **1987**, 109, 6213. (d) Kerrick, S. T.; Beak, P. *J. Am. Chem. Soc.* **1991**, 113, 9708. (e) Park, Y. S.; Boys, M.; Beak, P. *J. Am. Chem. Soc.* **1996**, 118, 3757. (f) Hodgson, D. M.; Lee, G. P. *Chem. Commun.* **1996**, 1015. (g) Bannan, Y. L.; Hanessian, S. *Chem. Rev.* **1997**, 97, 3161. (h) Park, Y. S.; Beak, P. *J. Org. Chem.* **1997**, 62, 1574. (i) Hoppe, D. M.; Hense, T. *Angew. Chem., Int. Ed.* **1997**, 36, 2282 and references cited therein. (j) Lucet, D.; LeGall, T.; Mioskowski, C. *Angew. Chem., Int. Ed.* **1998**, 37, 2580. (k) Li, X.; Schenkel, L. B.; Kozlowski, M. C. *Org. Lett.* **2000**, 2, 875. (l) Ganguly, B.; Freed, D. A.; Kozlowski, M. C. *J. Org. Chem.* **2001**, 66, 1103. (m) Xu, Z.; Kozlowski, M. C. *J. Org. Chem.* **2002**, 67, 3072. (n) Dearden, M. J.; Firkin, C. R.; Hermet, J.-P. R.; O'Brien, P. *J. Am. Chem. Soc.* **2002**, 124, 1870. (o) Kozlowski, M. C.; Li, X.; Carroll, P. J.; Xu, Z. *Organometallics* **2002**, 21, 4513. (p) Phuan, P.-W.; Ianni, J. C.; Kozlowski, M. C. *J. Am. Chem. Soc.* **2004**, 126, 15473.

- (2) Zadel, G.; Breitmaier, E. *Chem. Ber.* **1994**, 127, 1323.

[‡] Nara Institute of Science and Technology.

[†] Present address: Doshisha Women's College of Liberal Arts.





Figure 1. Chiral tertiary diamines.

which should give five ^{13}C -signals. It is implied that the product obtained by Breitmeier's procedure may not be chemically and/or optically pure. Therefore, we modified the procedure of the multigram scale preparation of (*S,S*)-**6** as follows (Scheme 1). L-Proline methyl ester hydrochloride,



obtained by thionyl chloride in methanol, was neutralized with sodium bicarbonate (powder) in dichloromethane and the resulting precipitate was filtered and the filtrate was concentrated in vacuo. Then, the neat residue was stirred at


room temperature for several days. The resulting diketopiperazine **8a**³ was purified by silica gel column chromatography and recrystallization. Reduction of **8a** with LiAlH_4 and bulb-to-bulb distillation gave pure (*S,S*)-**6**, with only five ^{13}C NMR signals as expected, in high yield.

With the chiral diamine (*S,S*)-**6** in hand, synthesis of a variety of substituted chiral diamines, which may have enhanced potential as the chiral ligand for asymmetric catalyst, was investigated here. Diastereoselective alkylation of **8a** via potassium enolate with Romo's method⁴ gave the monoalkylated products **8b–d** in good yield. Then, an introduction of the same alkyl group on another bridgehead carbon center was attempted to obtain C_2 -symmetric *cis*-dialkylated products. However, obtaining C -methylation of monomethyl diketopiperazine **8b** by conventional methods with LDA or KHMDS as the hindered base and iodomethane in THF was unsuccessful. It was assumed that the methyl group on the bridgehead interfered with the approach of the amide base, therefore we employed lithium diethylamide (LDEA), a more accessible base than the hindered LDA, to give the desired chiral *cis*-dimethyl isomer **9b** in good yield as sole diastereomer (Table 1). Unfortunately, synthesis of *cis*-diallyl (**9c**) and *cis*-dibenzyl (**9d**) analogues was unsuccessful to give undesired achiral *trans*-isomers predominantly, presumably due to severe steric hindrance between the first alkyl group on the bridgehead center and the electrophile.

Finally, reduction of **8b** with LiAlH_4 gave the chiral diamine (*S,S*)-**7**, which was purified as dipicrate and regenerated by neutralization in 67% yield from **8b** (Scheme 2).

Acylation of alcohol is one of the most essential reactions in organic synthesis and many asymmetric acylations have been developed for preparation of small chiral compounds well suited for further manipulation of the optically active molecules. Enzymatic acylation of racemic alcohols by kinetic optical resolution and desymmetrization of prochiral diols have been extensively studied⁵ and almost established as potential methods for the preparation of certain chiral alcohol variants. Recently, catalytic asymmetric acylation of alcohols with achiral acylating agents has emerged in

Table 1. Diastereoselective Alkylation of Diketopiperazine



entry	diketopiperazine	base	RX	yield (%) ^a	cis:trans ^b
1	8b (R = Me)	KHMDS	Mel	N.R.	
2	8b	LDA	Mel	N.R.	
3	8b	LDA + HMPA	Mel	N.R.	
4	8b	LDEA	Mel	75	1:0
5	8c (R = allyl)	LDEA	AllylBr	65	1:8 ^c
6	8d (R = Bn)	LDEA	BnBr	78	0:1

^a Yield of dialkylated product (cis and/or trans) from **8b–d**. ^b Ratio was obtained by ^1H NMR of the crude mixture. ^c Inseparable mixture of epimers.

Table 2. Asymmetric Benzoylation of *meso*-1,2-Diols Catalyzed by CuCl₂-Chiral Diamines


<i>(S,S)</i> -6			<i>(S,S)</i> -7			(-)-sparteine			<i>(R,R)</i> -Ph ₂ box		
major enantiomer	%	% ee ^a	major enantiomer	%	% ee ^a	major enantiomer	%	% ee ^a	major enantiomer	%	% ee ^a
	56	18		73	79		61	66		20	12
	80	91		100	93		80	94		32	56
	82	31		82	70		98	77		98	39
	58	3		77	92		55	94		51	78
	66	70		73	86		98	97		86	59

^a Optical purity was determined by chiral HPLC.

succession.⁶ Matsumura and co-workers reported excellent asymmetric acylation catalyzed by the complex of CuCl₂ and chiral bisoxazoline. Here, we applied chiral piperazine *(S,S)*-7 (DMPP) as the ligand to the above-noted asymmetric acylation.

Benzoylation of *meso*-1,2-diols with 3 mol % of CuCl₂-chiral diamines, such as *(S,S)*-6, *(S,S)*-7, (-)-sparteine, and bisoxazoline, for comparison, provided optically active monobenzoates in moderate to excellent enantioselectivity (Table 2). Desymmetrization of all the *meso*-diols investigated was achieved in high enantioselectivity with CuCl₂-*(S,S)*-7 complex, while use of CuCl₂-bisoxazoline and CuCl₂-*(S,S)*-6 exhibited relatively low optical yield. It was demonstrated that an introduction of two methyl groups on bridgehead centers of *(S,S)*-6 greatly enhanced the enantio-

selectivity. Interestingly, CuCl₂-(-)-sparteine complex showed high enantioselectivity giving the antipodes of mono-

Scheme 2. Reduction of Dimethyldiketopiperazine

- (3) (a) Li, J. P.; Biel, J. H. *J. Heterocycl. Chem.* **1968**, *5*, 703. (b) Vicar, J.; Smolikova, J.; Blaha, K. *Collect. Czech. Chem. Commun.* **1972**, *37*, 4060. (c) Rothe, M.; Mazanek, J. *Angew. Chem., Int. Ed.* **1972**, *11*, 293. (d) Vicar, J.; Budesinsky, M.; Blaha, K. *Collect. Czech. Chem. Commun.* **1973**, *38*, 1940. (e) Ueda, T.; Saito, M.; Kato, T.; Izumiya, N. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 568. (f) Basiuk, V. A.; Gromovoy, T. Yu.; Chunko, A. A.; Soloshonok, V. A.; Kukhar, V. P. *Synthesis* **1992**, 449. (g) Ishihara, K.; Ohara, S.; Yamamoto, H. *J. Org. Chem.* **1996**, *61*, 4196.
- (4) Poullennec, K. G.; Kelly, A. T.; Romo, D. *Org. Lett.* **2002**, *4*, 2645.
- (5) Reviews: (a) Whitesides, G. M.; Wong, C.-H. *Angew. Chem., Int. Ed.* **1985**, *24*, 617. (b) Chen, C.-S.; Sih, C. J. *Angew. Chem., Int. Ed.* **1989**, *28*, 695. (c) Klibanov, A. M. *Acc. Chem. Res.* **1990**, *23*, 114. (d) Drucekhammer, D. G.; Hennen, W. J.; Pederson, R. L.; Barbas, C. F., III; Gautheron, C. M.; Krach, T.; Wong, C.-H. *Synthesis* **1991**, 499. (e) Wong, C.-H.; Whitesides, G. M. *Enzymes in Synthetic Organic Chemistry*; Pergamon: New York, 1994.
- (6) For nonenzymatic asymmetric acylation of *meso*-diols: (a) Mukaiyama, T.; Tomioka, I.; Shimizu, M. *Chem. Lett.* **1984**, 49. (b) Mukaiyama, T.; Yanabe, Y.; Shimizu, M. *Chem. Lett.* **1984**, 404. (c) Ichikawa, J.; Asami, M.; Mukaiyama, T. *Chem. Lett.* **1984**, 949. (d) Ishihara, K.; Kubota, M.; Yamamoto, H. *Synlett* **1994**, 611. (e) Vedejs, E.; Daugulis, O.; Diver, S. T. *J. Org. Chem.* **1996**, *61*, 430. (f) Oriyama, T.; Imai, K.; Hosoya, T.; Sano, T. *Tetrahedron Lett.* **1998**, *39*, 397. (g) Oriyama, T.; Imai, K.; Sano, T.; Hosoya, T. *Tetrahedron Lett.* **1998**, *39*, 3529. (h) Sano, T.; Imai, K.; Ohoishi, K.; Oriyama, T. *Chem. Lett.* **1999**, 265. (i) Traugott, B. M.; Mino, T. *J. Am. Chem. Soc.* **2003**, *125*, 2410. (j) Matsumura, Y.; Maki, T.; Murakami, S.; Onomura, O. *J. Am. Chem. Soc.* **2003**, *125*, 2052. (k) Kawabata, T.; Stragies, R.; Fukaya, T.; Nagaoka, Y.; Schedel, H.; Fujii, K. *Tetrahedron Lett.* **2003**, *44*, 1545. (l) Marzet, C.; Roseblade, S.; Kohler, V.; Pfaltz, A. *Org. Lett.* **2006**, *8*, 1879.

benzoates obtained by use of the CuCl_2 –(*S,S*)-**7** complex. Although the availability of D-proline is severely limited, (–)-sparteine could serve as the pseudoenantiomer of (*S,S*)-**7**.

In conclusion, the synthesis of the novel class of alicyclic chiral piperazine ligand (*S,S*)-**7** has been developed. It was also demonstrated that (*S,S*)-**7** is an effective ligand for copper-catalyzed asymmetric acylation of *meso*-1,2-diols. Mechanistic studies of asymmetric acylation and further application of other asymmetric transformations with (*S,S*)-**7** are currently under investigation.

Acknowledgment. This work was supported by Grant-in-Aid for Scientific Research (C) (No. 18590023) from the Japan Society for the Promotion of Science.

Supporting Information Available: Synthetic procedure, spectral and analytical data for (*S,S*)-**6**, (*S,S*)-**7**, and asymmetric acylation of *meso*-1,2-diols. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0626387