


# Addition of Amines to Nitriles Catalyzed by Ytterbium Amides: An Efficient One-Step Synthesis of Monosubstituted *N*-Arylamidines

Junfeng Wang,<sup>†</sup> Fan Xu,<sup>†</sup> Tao Cai,<sup>†</sup> and Qi Shen<sup>\*†‡</sup>

Key Laboratory of Organic Synthesis of Jiangsu Province, Department of Chemistry and Chemical Engineering, Dushu Lake Campus, Suzhou University, Suzhou 215123, People's Republic of China, and State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China  
qshen@suda.edu.cn

Received November 14, 2007

## ABSTRACT



A one-step synthesis of monosubstituted *N*-arylamidines via addition of amines to nitriles catalyzed by ytterbium amides is reported. The reactions with various substrates give the products in good to excellent yields with 5 mol % ytterbium at 100 °C under solvent-free conditions.

Amidines are of interest as structural units with wide utility in drug design<sup>1</sup> and as synthons for the synthesis of heterocyclic compounds.<sup>2</sup> Several synthetic strategies have been developed<sup>3,4</sup> in which the nucleophilic addition of amine to nitrile is the most convenient and atom-economic method.<sup>3</sup> The one-step synthesis of amidines from nitriles and amines can be realized only if the nitriles are activated by electron-withdrawing groups<sup>3a</sup> or under more forcing conditions in

the presence of Lewis acids<sup>3b</sup> or with aluminum amides<sup>3c</sup> for unactivated nitriles. Trivalent lanthanide triflates have been employed successfully as catalysts in the condensation

(2) (a) Dunn, P. J. In *Comprehensive Organic Functional Group Transformations II*; Katritzky, A. R., Taylor, R. J., Eds.; Elsevier: Oxford, 2005; Vol. 5, pp 655–698. (b) Yokoyama, M.; Menjo, Y.; Wei, H.; Togo, H. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 2735. (c) Lerestif, J. M.; Bazureau, J. P.; Hamelin, J. *Tetrahedron Lett.* **1993**, *34*, 4639. (d) Doise, M.; Blondeau, D.; Sliwa, H. *Synth. Commun.* **1992**, *22*, 2891. (e) Neunhoeffer, H.; Karafiat, U.; Köhler, G.; Sowa, B. *Liebigs Ann. Chem.* **1992**, *115*. (f) Alanine, A. I. D.; Fishwick, C. W. G. *Tetrahedron Lett.* **1989**, *30*, 4443.


(3) (a) Grivas, J. C.; Taurins, A. *Can. J. Chem.* **1961**, *39*, 761. (b) Oxley, P.; Partridge, M. W.; Short, W. F. *J. Chem. Soc.* **1947**, 1110. (c) Garigipati, R. S. *Tetrahedron Lett.* **1990**, *31*, 1969. (d) Fersberg, J. H.; Spaziano, V. T.; Balasubramanian, T. M. *J. Org. Chem.* **1987**, *52*, 1017. (e) Xu, F.; Sun, J.; Shen, Q. *Tetrahedron Lett.* **2002**, *43*, 1867. (f) Rousselet, G.; Capdevielle, P.; Maumy, M. *Tetrahedron Lett.* **1993**, *34*, 6395. (g) Anbazhagan, M.; Boykin, D. W.; Stephens, C. E. *Tetrahedron Lett.* **2001**, *43*, 9089. (h) Anbazhagan, M.; Stephens, C. E.; Boykin, D. W. *Tetrahedron Lett.* **2002**, *43*, 4221. (i) Khanna, I. K.; Yu, Y.; Huff, R. M.; Weier, R. M.; Xu, X.; Koszyk, F. J.; Collins, P. W.; Cogburn, J. N.; Isakson, P. C.; Koboldt, C. M.; Masferrer, J. L.; Perkins, W. E.; Seibert, K.; Veenhuizen, A. W.; Yuan, J.; Yang, D.-C.; Zhang, Y. Y. *J. Med. Chem.* **2000**, *43*, 3168. (j) Zhou, S.; Wang, S.; Yang, G.; Li, Q.; Zhang, L.; Yao, Z.; Zhou, Z.; Song, H. *Organometallics* **2007**, *26*, 3755. (k) Zhang, W.-X.; Nishiura, M.; Hou, Z. *J. Am. Chem. Soc.* **2005**, *127*, 16788.

<sup>†</sup> Suzhou University.

<sup>‡</sup> Shanghai Institute of Organic Chemistry.

(1) (a) Taveras, A. G.; Chao, J.; Biju, P. J.; Yu, Y.; Fine, J. S.; Hipkin, W.; Aki, C. J.; Merritt, J. R.; Li, G.; Baldwin, J. J.; Lai, G.; Wu, M.; Hecker, E. A. WO-2004033440, 2004. (b) Varghese, J.; Maillard, M.; Jagodzinska, B.; Beck, J. P.; Gailunas, A.; Fang, L.; Sealy, J.; Tenbrink, R.; Freskos, J.; Mickelson, J.; Samala, L.; Hom, R. WO-2003040096, 2003. (c) Sugino, M.; Sugita, S. JP-2004175720, 2004. (d) Takamuro, I.; Honma, K.; Ishida, A.; Taniguchi, H.; Onoda, Y. JP-2004115450, 2004. (e) Maguire, M. P.; Dai, M.; Vos, T. J. WO-2002062766, 2002. (f) Sunagawa, M.; Hebeisen, P. WO-2002048149, 2002. (g) Lin, Y. I.; Wang, B. S.; Ruzsala-Mallon, V. M.; Bitha, P.; Fields, T. L. EP-354303, 1990. (h) Moeller, H.; Gloxhuber, C. DE-2364437, 1975. (i) Zhu, B.-Y.; Su, T.; Li, W.; Goldman, E. A.; Zhang, P.; Jia, Z. J.; Scarborough, R. M. WO-2002026734, 2002. (j) Bullock, W. H.; Kluender, H. C. E.; Collibee, W. L.; Dally, R.; Rodriguez, M. E.; Wang, M. WO-2002020526, 2002. (k) Ito, K.; Spears, G. W.; Yamada, A.; Toshima, M.; Kato, M. JP-2001220375, 2001.

Scheme 1. Tested Amides




of nitriles with primary amines and diamines into *N,N'*-disubstituted and cyclic amidines, but monosubstituted amidines cannot be synthesized, and the reaction with secondary amines afforded triazines or pyrimidines via further condensations.<sup>3d</sup> SmI<sub>2</sub> can also catalyze the condensation reaction as a precatalyst, and the same problem as in the case with lanthanide triflates remains to be solved.<sup>3e</sup> Recently, the selective synthesis of monosubstituted alkylamidines from amines and nitriles can be achieved in the presence of CuCl. However, 1.2 equiv of CuCl to amines is needed in this process.<sup>3f</sup> Thus, the development of catalytic addition of amines to nitriles is still a challenge in monosubstituted amidine synthesis.

Prompted by recent successes obtained in C–N bond formation, reactions catalyzed by lanthanide amide complexes, including hydroamination,<sup>5</sup> monocoupling reaction of isocyanides with terminal alkynes,<sup>6</sup> a Cannizzaro-type disproportionation of aromatic aldehydes,<sup>7</sup> and the guanylation of amines<sup>8</sup> and amidination of alkyne,<sup>3j,k</sup> we turned our attention to the possible use of lanthanide amides as catalysts for addition reaction of amines to nitriles.

We have recently synthesized the novel amide complexes supported by a bridged bis(amidinate) **I–III** (Scheme 1) and found that amide complexes **I** and **II** show high reactivity

for controlled polymerization of lactide.<sup>9</sup> Thus, complex **I** was first tested in the condensation reactions of nitriles and amines, and several reaction conditions were examined in the model reaction of **1a** with **2a** (Table 1). The reactions

Table 1. Optimization of the Reaction Conditions<sup>a</sup>



entry	cat. (mol %)	solvent	<b>1a/2a</b>	<i>T</i> (°C)	time (h)	% yield <sup>b</sup>
1	5	toluene	1:1	60	12	35
2	5	toluene	1:1	60	24	45
3	10	THF	1:1	60	12	45
4	10	toluene	1:1	60	12	80
5	5	toluene	1:1	80	24	55
6	5	toluene	1:1	100	24	65
7	5		1:1	100	24	75
8	5		1:2	100	24	88
9	5		1:3	100	24	84
10	5		1:4	100	24	83
11	5		1:10	100	24	65
12	10		1:2	100	12	97

<sup>a</sup> All data were obtained using 1.90 mmol of nitrile. <sup>b</sup> Isolated yields.

(4) Recent examples: (a) Yet, L. *A Survey of Amidine Synthesis*; Technical Report No. 3; Albany Molecular Research, Inc.: New York, 2000; Vol. 4. (b) Tidwell, R. R.; Boykin, D. W. In *Dicationic DNA Minor Groove Binders as Antimicrobial Agents in Small Molecule DNA and RNA Binders: From Synthesis to Nucleic Acid Complexes*; Demeunynck, M., Bailly, C., Wilson, W. D., Eds.; Wiley-VCH: Hoboken, NJ, 2002. (c) Tucker, J. A.; Clayton, T. L.; Chidester, C. G.; Schulz, M. W.; Harrington, L. E.; Conrad, S. J.; Yagi, Y.; Oien, N. L.; Yurek, D.; Kuo, M.-S. *Bioorg. J. Med. Chem.* **2000**, *8*, 601. (d) Stephens, C. E.; Tanious, F.; Kim, S.; Wilson, D. W.; Schell, W. A.; Perfect, J. R.; Franzblau, S. G.; Boykin, D. W. *J. Med. Chem.* **2001**, *44*, 1741. (e) Werbovetz, K.; Brendle, J.; Stephens, C. E.; Boykin, D. W. US Patent 60/246,330, filed 11/7/00. (f) Kusturin, C. L.; Liebeskind, L. S.; Neumann, W. L. *Org. Lett.* **2002**, *4*, 983.

(5) For general reviews on hydroamination reactions, see: (a) Müller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675. (b) Nobis, M.; Driessen-Hölscher, B. *Angew. Chem., Int. Ed.* **2001**, *40*, 3983. (c) Brunet, J. J.; Neibecker, D. In *Catalytic Heterofunctionalization from Hydroamination to Hydrozirconation*; Togni, A., Grützmacher, H., Eds.; Wiley-VCH: Weinheim, Germany, 2001; pp 91–141. (d) Roesky, P. W.; Müller, T. E. *Angew. Chem., Int. Ed.* **2003**, *42*, 2708. (e) Pohlki, F.; Doye, S. *Chem. Soc. Rev.* **2003**, *32*, 104. (f) Beller, M.; Tillack, A.; Seayad, J. In *Transition Metals for Organic Synthesis*, 2nd ed.; Beller, M., Bolm, C., Eds.; Wiley-VCH: Weinheim, Germany, 2004; Vol. 2, pp 403–414.

(6) (a) Komeyama, K.; Sasayama, D.; Kawabata, T.; Takehira, K.; Takali, K. *J. Org. Chem.* **2005**, *70*, 10679. (b) Komeyama, K.; Sasayama, D.; Kawabata, T.; Takehira, K.; Takaki, K. *Chem. Commun.* **2005**, 634.

(7) Zhang, L.; Wang, S.; Zhou, S.; Yng, G.; Sheng, E. *J. Org. Chem.* **2006**, *71*, 3149.

(8) (a) Li, Q.; Wang, S.; Zhou, S.; Yang, G.; Zhu, X.; Liu, Y. *J. Org. Chem.* **2007**, *72*, 6763. (b) Zhang, W.-X.; Nishiura, M.; Hou, Z. *Synlett* **2006**, *8*, 1213. (c) Zhang, W.-X.; Nishiura, M.; Hou, Z. *Chem. Eur. J.* **2007**, *13*, 4037.

took place smoothly under solvent free conditions or in toluene to afford the monosubstituted amidine **3aa** in moderate to excellent yields depending on the conditions used. The reaction in THF provided the lowest yield (Table 1, entry 3). With the increasing temperature, the yields increased (Table 1, entries 2, 5, and 6). The ratio of amine to nitrile showed an effect on the reactivities, and the maximum yields were obtained in the range of 2–4 (Table 1, entries 8–10). The yields increased when the catalyst loading increased and an almost quantitative yield was obtained with 10 mol % of ytterbium (based on **1a**) under solvent-free conditions for 12 h (Table 1, entry 12).


Since **I** performed well in this reaction, we screened other lanthanide amides **II–IV**<sup>10</sup> (Scheme 1) in the reactions with several substrates. All reactions were conducted with 5 mol % of ytterbium at 100 °C in the absence of solvent (Table 2).

(9) Wang, J.; Cai, T.; Yao, Y.; Zhang, Y.; Shen, Q. *Dalton Trans.* **2007**, 5275.

(10) Zhou, S.; Wang, S.; Yang, G.; Liu, X.; Sheng, E.; Zhang, K.; Cheng, L.; Huang, Z. *Polyhedron* **2003**, *22*, 1019.



(see the mechanism proposed, Scheme 2). All products were characterized by  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR to be amidines via 1,3-H shift and the structure of **3aa** was further confirmed by X-ray structural analysis (Figure 1). The crystal structure



**Figure 1.** Molecular structure of **3aa**, drawn with 30% thermal ellipsoids.

showed a double-bond character for the N(2)–C(1) bond with a distance of 1.290(3) Å.

The reaction mechanism was proposed as shown in Scheme 2. Reaction of an aniline with the lanthanide amide gave the new amido intermediate **A** through an acid–base reaction.<sup>11</sup> A nitrile then was coordinated to the center metal forming a complex **B**. An intramolecular insertion of amide to cyano of nitrile gave the corresponding intermediate **C** as reported previously.<sup>12</sup>

Intermediate **C** underwent protonation by amine to release the product and to generate the lanthanide amide active species **A**. When the reaction of intermediate **C** with additional nitrile is more favorable than that with amine, triazine **E** was produced as the main product.

In summary, we have developed an efficient protocol to the one-step synthesis of monosubstituted amidines in good to excellent yields via a nucleophilic addition of amines to nitriles catalyzed by ytterbium amides under solvent free conditions. The reactions of aromatic nitriles with aromatic primary amines proceed very well to give excellent yields. The present strategy can be viewed as a compensative method for the published intramolecular nucleophilic addition of amine to activated nitrile methods with  $\text{Ln}(\text{OTf})_3$  and/or  $\text{SmI}_2$  as the catalysts,<sup>3d,e</sup> which frequently suffered from difficulty in synthesis of monosubstituted amidines. However, the reaction with aliphatic amines has not yet been successful. Efforts in this direction are ongoing in our laboratory.

**Acknowledgment.** We are grateful to the National Natural Science Foundation of China (Grant No. 20632042) for financial support.

**Supporting Information Available:** Experimental procedures and characterization data; copies of  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL702739C

(11) (a) Schuetz, S. A.; Day, V. W.; Sommer, R. D.; Rheingold, A. L.; Belot, J. A. *Inorg. Chem.* **2001**, *40*, 5292. (b) Bradley, D. C.; Ghotra, J. S.; Hart, F. A. *J. Chem. Soc., Dalton Trans.* **1973**, 1021. (c) Alyea, E. C.; Bradley, D. C.; Copperthwaite, R. G. *J. Chem. Soc., Dalton Trans.* **1972**, 1580.

(12) (a) Bercaw, J. E.; Davies, D. L.; Wolczanski, P. T. *Organometallics* **1986**, *5*, 443. (b) Evans, W. J.; Fujimoto, C. H.; Ziller, J. W. *Organometallics* **2001**, *20*, 4529.