

Diastereoselective Syntheses of Chroman Spiroketal via [4 + 2] Cycloaddition of Enol Ethers and *o*-Quinone Methides


Maurice A. Marsini, Yaodong Huang, Christopher C. Lindsey, Kun-Liang Wu, and Thomas R. R. Pettus*

Department of Chemistry and Biochemistry, University of California at Santa Barbara
Santa Barbara, California 93106-9510

pettus@chem.ucsb.edu

Received February 12, 2008

ABSTRACT



A variety of chroman spiroketals are synthesized via inverse-demand [4 + 2] cycloaddition of enol ethers and *ortho*-quinone methides (*o*-QMs). Low temperature *o*-QM generation in situ allows for the kinetic, diastereoselective construction of these motifs, providing entry to a number of unusual chroman spiroketal natural products.

Aliphatic spiroketals are a common substructure of natural products isolated in a large variety from both marine and terrestrial sources.¹ Studies aimed at understanding the origins of their conformational preference have resulted in the general assumption that the spiroketal carbon corresponds to the thermodynamically most stable isomer as determined by stereoelectronic influences. Hence, the biosynthesis of these fluxional natural products usually results in the construction of the most stable diastereomer or a mixture that reflects energy differences between respective diastereomers.

For some time, we have speculated that a rare subset consisting of chroman spiroketals (**1**,² **2**,³ and **3**,⁴ Figure 1) may possess features that refute this basic tenet and require a kinetic assembly. These uniquely robust chroman spiroketal




Figure 1. Natural products that contain a chroman spiroketal.

scaffolds have caused us to consider new methods for their construction.

Some time ago, we developed a method that enables the controlled, low-temperature generation of *o*-quinone methides (*o*-QMs) (Figure 2, I).⁵ The process, which is driven by the relative stability of sequential anions formed along the cascade, begins by formation of an alkoxide. This species

(1) Perron, F.; Albizzati, K. F. *Chem. Rev.* **1989**, *89*, 1617.

(2) (a) Puder, C.; Loya, S.; Hizi, A.; Zeeck, A. *Eur. J. Org. Chem.* **2000**, 729. (b) Brockmann, H.; Lenk, W.; Schwantje, G.; Zeeck, A. *Tetrahedron Lett.* **1966**, *7*, 3525. (c) Brockmann, H.; Lenk, W.; Schwantje, G.; Zeeck, A. *Chem. Ber.* **1969**, *102*, 126. (d) Brockmann, H.; Zeeck, A. *Chem. Ber.* **1970**, *103*, 1709.

(3) Stierle, A. A.; Stierle, D. B.; Kelly, K. *J. Org. Chem.* **2006**, *71*, 5357.

(4) (a) Namikoshi, M.; Kobayashi, H.; Yoshimoto, T.; Meguro, S. *Chem. Lett.* **2000**, 308. (b) Namikoshi, M.; Kobayashi, H.; Yoshimoto, T.; Meguro, S. *Chem. Pharm. Bull.* **2000**, 1452.




Figure 2. (I) Cascade leading to *o*-QMs. (II) Preferred endo transition state for [4 + 2] cycloaddition with chiral vinyl ether derivatives. (III) Proposal to examine the effects of R¹, R², R³ on the reactions of methylene dihydrofurans.

intercepts a neighboring carbonyl of a phenolic carbonate and thus liberates a phenoxide that subsequently undergoes β -elimination of a carboxylate and thereby forms a reactive *o*-QM species, which captures the first nucleophile that it subsequently encounters.

This is a useful method for *o*-QM generation because, since the reactive *o*-QM intermediate is generated at low temperature in low concentrations, it subsequently participates in very precise, kinetically controlled reactions.⁶ Using this procedure, we demonstrated the first examples of diastereoselective *o*-QM cycloadditions and found that the inverse demand [4 + 2] significantly favors an endo transition state with electron-rich alkenes.⁷ In further experiments, we showed that chiral enol ethers containing remote stereocenters, such as those derived from (1*S*,2*R*)-2-phenylcyclohexanol, will participate in diastereoselective cycloadditions to yield the corresponding chroman acetals with an *R* configuration (Figure 2, II).⁸ On the basis of these findings, we began to speculate that an assortment of chiral 2-methylene tetrahydrofurans with R¹, R², and R³ substituents should also participate in cycloadditions with similar diastereoselectivity (Figure 2, III). Detailed analysis of our proposed transition state allowed us to further speculate that substituents R¹ and R³ would have a pronounced effect on the stereochemical outcome, whereas the R² residue would not, allowing it to be tolerated on the same face as the reaction. Therefore, we began to examine model systems that might

(5) Van de Water, R. W.; Magdziak, D. J.; Chau, J. N.; Pettus, T. R. R. *J. Am. Chem. Soc.* **2000**, *122*, 6502.

(6) Van de Water, R. W.; Pettus, T. R. R. *Tetrahedron* **2002**, *58*, 5367.


(7) Jones, R. M.; Selenski, C.; Pettus, T. R. R. *J. Org. Chem.* **2002**, *67*, 6911.

(8) (a) Selenski, C.; Pettus, T. R. R. *J. Org. Chem.* **2004**, *69*, 9196. (b) Selenski, C.; Mejorado, L. H.; Pettus, T. R. R. *Synlett* **2004**, *6*, 1101.

eventually be developed into strategies leading to the construction of the chroman spiroketal natural products 1–3.

Our investigation began with the preparation of known enol ethers 4,⁹ 5,¹⁰ 6,¹¹ and 7,¹² as well as the unknown enol ether 8, which we prepared from the known acid 11¹³ by sequential acetalization with formaldehyde and methylenation with Tebbe's reagent (Scheme 1).

Scheme 1. Known and Readily Available Five-Membered Ring Enol Ethers Containing Exocyclic Olefins



We also prepared enol ethers 9 and 10 in a short sequence. We began by subjecting the unsaturated lactone 13 (2.4 M in THF) to lithiated acetonitrile 14, which results in exclusive 1,4-addition. Use of the latter as the nucleophile avoids the problem of an undesirable proton transfer during the course of the reaction. Thus, the intermediate enolate undergoes subsequent C-methylation with methyl iodide and affords the *trans*-substituted lactone 15 in 73% yield (>10:1 dr). Freshly prepared Tebbe reagent¹⁴ fails to provide the desired product and returns 15 unchanged. However, if 15 (0.1 M in THF) is refluxed with Petasis' reagent (2.5 equiv, 4 h) and sequentially treated to a methanol/water workup, the enol ether 10 arises whereby the nitrile functionality has also been transformed into a methyl ketone. While investigation of the crude product mixture by ¹H NMR showed compound 10 as the only identifiable product, bulb-to-bulb distillation with base washed glassware provided only a 20–30% yield. To the best of our knowledge, the conversion of a nitrile into a methyl ketone by the Petasis reagent has never before been reported.¹⁵ Doxsee has observed a related transformation with

(9) Okazoe, T.; Takai, K.; Oshima, K.; Utimoto, K. *J. Org. Chem.* **1987**, *52*, 4410.

(10) Pine, S. H.; Zahler, R.; Evans, D. A.; Grubbs, R. H. *J. Am. Chem. Soc.* **1980**, *102*, 3270.

(11) Petasis, N. A.; Lu, S. *J. Am. Chem. Soc.* **1995**, *117*, 6394.


(12) Diaz-Ortiz, A.; Diez-Barra, E.; de la Hoz, A.; Prieto, P. *Synth. Comm.* **1993**, *23*, 1935.

(13) Baltzer, S.; Schoentjes, B.; Van Dorsselaer, V. *Fr. Demande* **2004**, 85.

Tebbe's reagent in the presence of DMAP and trimethyl phosphine.¹⁶ If the amount of THF is limited, however, to the minimal amount required for solubility (toluene/THF, 3:2), the reaction affords the enol ether **9** (4:1 ratio of **9/10**) in a 40–50% yield after flash chromatography and distillation.

With these enol ethers in hand, we began examining their reactivity with *o*-QMs. When the known benzyl alcohol **16**¹⁷ (0.1 M in toluene, –78 °C, 2–4 equiv of the 1,3-dihydroisobenzofuran enol ether **4**) is treated with *t*-BuMgCl, an endo-selective cycloaddition proceeds as the reaction mixture slowly warms to room temperature (Scheme 2). The reaction cleanly produces the chroman spiroketal **17** in 60% isolated

Scheme 2. Some [4 + 2] Cycloadditions of Benzofurans and an *o*-QM



yield (>8:1 dr by crude ¹H NMR). This transformation, therefore, serves as a practical model for an eventual synthetic strategy aimed toward (+)-paecilospirone (**3**).

We next examined a similar cycloaddition using the enol ether **5**. The 2-methylbenzofuran **18** is produced along with a small amount of the desired chroman spiroketal **19** in 10% yield. Presumably, the fragile 2,3-dihydrobenzofuran enol ether **5** succumbs to isomerization under these conditions.¹⁸ Therefore, due to the low yield for this transformation, we sought an alternative strategy for the construction of **1** by investigating *o*-QM cycloadditions with the enol ethers **6–8**.

The starting benzyl alcohol **16** (0.1 M in Et₂O at –78 °C, 2 equiv of **6**) was subjected again to *t*-BuMgCl (Table 1). A cycloaddition occurs as the mixture slowly warms to room temperature, affording a single diastereomer as determined by crude ¹H NMR (45% isolated yield of **21**). On the basis of our extensive experience with these cycloadditions, we

(14) Payack, J. F.; Hughes, D. L.; Cai, D.; Cottrell, I. F.; Verhoeven, T. R. *Org. Synth.* **2002**, *79*, 19.

(15) For reactions of the Petasis reagent with nitriles, see: (a) Petasis, N. A.; Fu, D.-K. *Organometallics* **1993**, *12*, 3776. (b) Barluenga, J.; del Poso Lasada, C.; Olano, B. *Tetrahedron Lett.* **1992**, *33*, 7579. (c) Treatment of phenylacetonitrile with Petasis' reagent smoothly affords 1-phenylacetone in 75% yield. Thus, this mild nitrile-to-ketone transformation appears general and proceeds in better yields with less complicated examples.

(16) (a) Doxsee, K. M.; Farahi, J. B. *J. Am. Chem. Soc.* **1988**, *110*, 7239. (b) Takeda, T. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 195. (c) Doxsee, K. M.; Farahi, J. B. *J. Chem. Soc., Chem. Commun.* **1990**, *20*, 1452.

(17) (a) Marsini, M. A.; Huang, Y.; Van De Water, R. W.; Pettus, T. R. *Org. Lett.* **2007**, *7*, 3229. (b) Lindsey, C. C.; Pettus, T. R. *Tetrahedron Lett.* **2006**, *47*, 201.

(18) (a) Zhou, G.; Zheng, D.; Da, S.; Xie, Z.; Li, Y. *Tetrahedron Lett.* **2006**, *47*, 3349.

(19) Hoarau, C.; Pettus, T. R. *Org. Lett.* **2006**, *8*, 2843.


Table 1. Cycloadditions of Methylene-1,3-dioxolanes

enol ether	benzyl alcohol	product	yield	dr
6	16	21	45%	>20:1
7	20	22	52%	>20:1
8	16	23	45%	>20:1

speculate that the product **21** has the stereochemistry shown, which reflects a reaction proceeding through an endo orientation on the face opposite of phenyl residues at R¹ and R³. Next, we examined the corresponding reaction of enol ether **7** and benzyl alcohol **20**¹⁹ using similar conditions. Again, ¹H NMR revealed formation of **22** as a single diastereomer (>20:1). This result suggested to us that the presence of an R¹ substituent was sufficient to control the diastereoselectivity of the reaction. Thus, we were not altogether surprised that use of the enol ether **8**, which expresses a substituted aryl ring, undergoes a diastereoselective reaction with the *o*-QM derived from **16** in comparable ratio and yield.

With the chroman spiroketal **23** in hand, we paused to investigate its reformulation into the corresponding benzofuran spiro-adduct (Scheme 3). The benzyl ether in **23** (0.5

Scheme 3. Failed Attempts to Effect Trans-Ketalization



M in 95% EtOH) was cleaved in nearly quantitative yield by exposure of the heterogeneous mixture containing 5% Pd/C to a hydrogen atmosphere to provide phenol **24**. Subsequent treatment of **24** with various proton sources, as well as a cadre of Lewis acids, failed to afford the desired spiro-adduct **26** (via **25**). In most instances, the starting material remained unchanged or suffered decomposition.

The resiliency of the chroman spiroketal **24** and related structures is appreciated through an understanding of resonance effects. The lone pair of electrons on the chroman oxygen is delocalized throughout the adjoining aromatic ring. Therefore, it is difficult for the lone pair of electrons to assist

in the departure of the aliphatic ether, which results in the formation of the oxonium **25**. Conversely, the lone pair of electrons belonging to the chroman oxygen is also difficult to protonate, which would lead to the cleavage of the corresponding aryl ether. These facts also account for the unusual stability of MOM-aryl ethers as compared to the corresponding aliphatic variety. On the basis of these results, we abandoned an *o*-QM strategy for **1** and devised an alternative method for the construction of naphthoquinone spiroketals.²⁰

Our attention now turned toward the chroman spiroketal framework found in the natural product **2**. Similar treatment of **27** and **9** with *t*-BuMgCl affords **29** along with its diastereomer in a 61% yield as an inseparable 3:1 ratio (Table 2). We assumed that the R¹ substituent (–Me), which is

Table 2. Cycloadditions of Some Exocyclic Furan Enol Ethers

enol ether		benzyl alcohol	product	yield	dr	
9			29	61%	3:1	
9			30	77%	3.5:1	
9			31	72%	4.5:1	
10	28			32	72%	4.5:1

closer to the site of bond formation, exerted greater stereocontrol than the R² substituent (–CH₂CN), as the latter is further removed from the site of reaction. Nevertheless, the ratio was disappointing and we decided to examine the effects of electronics within the *o*-QM on the stereoselectivity for the reaction. Upon exchange of the –OBn for a more electron deficient –OBoc residue, the reaction of the *o*-QM derived from **20** exhibits a small increase in selectivity (3.5:1) favoring the chroman spiroketal **30**. Unfortunately, the diastereomers again prove inseparable by chromatography (77% combined yield). Introduction of a bromine atom results in a higher yield and better selectivity (4.5:1) for **31**.²¹ Moreover, the bromine atom assists in the subsequent separation of diastereomers by chromatography. Similar findings were observed for the ketone **32** that arises from

(20) (a) Wu, K.-L.; Wilkinson, S.; Reich, N. O.; Pettus, T. R. R. *Org. Lett.* **2007**, *9*, 5537. (b) Lindsey, C. C.; Wu, K.-L.; Pettus, T. R. R. *Org. Lett.* **2006**, *8*, 2365.

(21) Mejorado, L. H.; Hoarau, C.; Pettus, T. R. R. *Org. Lett.* **2004**, *6*, 1535.

the reaction of **28** and the enol ether **10**, which provides further evidence for the mildness of this method and its compatibility with a variety of functional groups. The lower diastereoselectivity compared with that in Table 1 may indicate that the steric bulk of R¹ (Figure 2) plays a substantial role in the stereochemical outcome of the reaction.

While we are confident of our structural assignments based upon analysis of coupling data and NOE effects, we wished to unequivocally establish the relative stereochemistry of adducts **17**, **21–23**, and **29–32**. Unfortunately, our attempts to crystallize these materials failed. We therefore decided to cleave the Boc group from compound **31**, in hope that the phenol **33** would crystallize. This task was accomplished by short exposure to 1 N LiOH. Crystallization from CH₂-Cl₂/MeOH (1:1) provided X-ray quality crystals of **33** (Figure 3). The resulting structure determination proved **31** arises



Figure 3. X-ray crystal structure of **33**.

from reaction of the enol ether in endo orientation on the face opposite the R¹ substituent and on the same side as the R³ substituent.

In conclusion, we provide the first example of kinetically controlled, diastereoselective construction of chiral chroman spiroketals using enol ethers and *o*-QMs. This versatile method, due to its stereocontrol and functional group compatibility, seems applicable for subsequent stereoselective synthetic routes toward chroman spiroketal-containing natural products (Scheme 1). These results will be reported in due course.

Acknowledgment. Support from the University of California, Cancer Research Coordinating Committee for investigations leading to the synthesis of **1** and **2** is greatly appreciated. We applaud Neil Schore (UC Davis) for his oversight of this important program.

Supporting Information Available: Experimental procedures and spectroscopic data (¹H NMR, ¹³C NMR, FT-IR, HRMS) for compounds **8–12**, **15**, **17**, **19**, **21–24**, **27**, and **29–33**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL8003244