

A Pentiptycene-Derived Light-Driven Molecular Brake


Jye-Shane Yang,^{*,†} Yao-Ting Huang,[†] Jinn-Hsuan Ho,[†] Wei-Ting Sun,[†] Hsin-Hau Huang,[†] Ying-Chih Lin,^{*,†} Shing-Jong Huang,[†] Shou-Ling Huang,[†] Hsiu-Feng Lu,[‡] and Ito Chao^{*,‡}

Department of Chemistry and Instrumentation Center, National Taiwan University, Taipei, Taiwan 10617, and Institute of Chemistry, Academia Sinica, Taipei, Taiwan 11529

jsyang@ntu.edu.tw; yclin@ntu.edu.tw; ichao@chem.sinica.edu.tw

Received March 26, 2008

ABSTRACT



A room-temperature light-driven molecular brake (1), consisting of a pentiptycene rotator, a 3,5-dinitrophenyl brake, and a photoisomerizable ethenyl spacer, is reported. The rotation rates of the rotator differ by about 9 orders of magnitude between the brake-on (*cis*-1) and brake-off (*trans*-1) states.

The ability to control specific motions of subunits in molecules or supramolecular ensembles is essential for the development of molecular machines.¹ As photons are cleaner, faster, and longer-range control elements (stimuli) compared to chemicals and electrons,² many efforts have been devoted to constructing prototypes of light-driven or photocontrollable linear and rotary motions.^{1–3} However, the progress in some aspects falls behind the corresponding chemical-controlled systems. One particular example is “molecular brakes”, in which the Brownian rotary motion of a rigid wheel component about a single bond can be slowed down by a brake unit. Since the first chemical-driven molecular brake reported

by Kelly and co-workers in 1994,⁴ several others have also been developed.⁵ The first approach toward a light-driven counterpart was reported by Feringa and co-workers in 1997 using a *cis*–*trans* interconvertible sterically overcrowded alkene.⁶ However, the desired rotary motion is restricted for both the *trans* and *cis* isomers at room temperature because of unexpectedly high and similar rotational barriers (19.7 and 19.0 kcal mol⁻¹, respectively, at 303 K). In a later report on a similar system, the rate of rotation (k_{rot}) was determined to be less than 0.1 s⁻¹ for all the stable and unstable states at 298 K.⁷ To date, an effective room-temperature photocontrollable molecular brake has yet to be demonstrated.

[†] National Taiwan University.

[‡] Academia Sinica.

(1) Balzani, V.; Venturi, M.; Credi, A. *Molecular Devices and Machines—A Journey into the Nano World*; Wiley-VCH: Weinheim, 2003.

(2) (a) Raymo, F. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 5249–5251. (b) Credi, A. *Aust. J. Chem.* **2006**, *59*, 157–169. (c) Saha, S.; Stoddart, J. F. *Chem. Soc. Rev.* **2007**, *36*, 77–92.

(3) (a) Kottas, G. S.; Clarke, L. I.; Horinek, D.; Michl, J. *Chem. Rev.* **2005**, *105*, 1281–1376. (b) Kay, E. R.; Leigh, D. A.; Zerbetto, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 72–191. (c) Feringa, B. L. *J. Org. Chem.* **2007**, *72*, 6635–6652.

(4) Kelly, T. R.; Bowyer, M. C.; Bhaskar, K. V.; Bebbington, D.; Garcia, A.; Lang, F.; Kim, M. H.; Jette, M. P. *J. Am. Chem. Soc.* **1994**, *116*, 3657–3658.

(5) (a) Jog, P. V.; Brown, R. E.; Bates, D. K. *J. Org. Chem.* **2003**, *68*, 8240–8243. (b) Annunziata, R.; Benaglia, M.; Cinquini, M.; Raimondi, L.; Cozzi, F. *J. Phys. Org. Chem.* **2004**, *17*, 749–751. (c) Alfonso, I.; Burguete, M. I.; Luis, S. V. *J. Org. Chem.* **2006**, *71*, 2242–2250.

(6) Schoevaars, A. M.; Kruizinga, W.; Zijlstra, R. W. J.; Veldman, N.; Spek, A. L.; Feringa, B. L. *J. Org. Chem.* **1997**, *62*, 4943–4948.

(7) ter Wiel, M. K. J.; van Delden, R. A.; Meetsma, A.; Feringa, B. L. *Org. Biomol. Chem.* **2005**, *3*, 4071–4076.

We report herein the first example of a room-temperature light-driven molecular brake (**1**) that displays distinct rates of rotation in the brake-on versus brake-off states. The rigid pentiptycene group⁸ in **1** serves as a four-bladed wheel, and the *trans*–*cis* photoisomerizable dinitrostyryl group behaves as a photoresponsive brake component (Figure 1). In the *trans*





Figure 1. Molecular structures of a light-driven molecular brake (**1**) corresponding to the schematic illustrations shown in the Abstract graphic. The *trans* (*trans*-**1**) and *cis* (*cis*-**1**) isomers correspond to the brake-off and brake-on states. The numerical labels for protons and carbons are for discussion of the VT NMR spectra.

form (*trans*-**1**, the brake-off state), the pentiptycene rotator undergoes free rotation ($k_{\text{rot}} \approx 10^9 \text{ s}^{-1}$) due to a small rotational barrier ($\Delta G^\ddagger \approx 4.5 \text{ kcal/mol}$ at 298 K). However, such a rotation is nearly blocked ($k_{\text{rot}} = 3 \text{ s}^{-1}$, $\Delta G^\ddagger \approx 16.4 \text{ kcal mol}^{-1}$ at 298 K) in the *cis* form (*cis*-**1**, the brake-on state) as the result of an intercalation of the dinitrophenyl group into the U-shaped cavities of the rotator. As a result, the rate of rotation in the brake-on and brake-off states differs by nearly 9 orders of magnitude at 298 K.

The synthesis of molecular brake **1** is outlined in Scheme 1. The commercially available 3,5-dinitrobenzyl alcohol **2**



was first converted with thionyl chloride to 3,5-dinitrobenzyl chloride **3**, which then reacted with triphenylphosphine to form the phosphonium salt **4**. The latter reaction was kept

(8) For a recent review on pentiptycene chemistry, see: Yang, J.-S.; Yan, J.-L. *Chem. Commun.* **2008**, 1501–1512.

under 60 °C, because the salt decomposes at higher temperatures. The stilbene backbone was finally constructed by the Wittig reaction of **4** and formylpentiptycene **5**, the synthesis of which has recently been reported.⁹ The resulting *trans* and *cis* isomers of **1** (2: 1) can be readily separated by column chromatography.

Figure 2 shows the aromatic region of ¹H and ¹³C NMR spectra of *trans*-**1** and *cis*-**1** in DMSO-*d*₆ at 298 K. The




Figure 2. Aromatic region of (a) ¹H and (b) ¹³C NMR spectra of *trans*-**1** and *cis*-**1** in DMSO-*d*₆ at 298 K. The solid circles denote peaks due to the dinitrostyryl group and the Arabic numerals denote the corresponding protons and carbons labeled in Figure 1.

presence of only one set of signals for *trans*-**1** suggests that both the pentiptycene rotator and the dinitrophenyl brake can undergo free rotation about the C_{vinyl}–C_{aryl} single bonds. In contrast, the observation of two sets of signals for the pentiptycene methine nuclei but not for the dinitrophenyl nuclei indicates that rotation of the rotator but not the brake component is slower than the NMR time scale. Except for some quaternary carbons, the signals can be unambiguously assigned on the basis of a series of 2D NMR experiments, including COSY, HSQC, NOESY, and ROESY (Supporting Information, Figures S5–S11), and the numerical labels in Figure 2 correspond to the labeled protons and carbons in Figure 1.

The information about rotational barriers and rates for the pentiptycene rotator in *cis*-**1** can be obtained from variable-temperature (VT) NMR studies and spectral simulations. As shown in Figure 3a, some of the proton peaks at 298 K coalesce at higher temperatures. For example, a coalescence temperature (*T*_c) near 348 K is found for protons H₃ and H_{3'}, corresponding to an energy barrier of $\Delta G^\ddagger_{(348\text{K})} = 16.9 \pm 0.2 \text{ kcal mol}^{-1}$. However, the multiplicity of proton signals and the presence of more than one type of spin systems in

(9) Yang, J.-S.; Ko, C.-W. *J. Org. Chem.* **2006**, *71*, 844–847.




Figure 3. Penttiptycene peripheral phenylene (blade) region of the (a) experimental proton and (b) carbon and (c) simulated carbon VT NMR spectra of *cis*-**1** (9 and 60 mM for proton and carbon, respectively, DMSO-*d*₆, 500 MHz). Values of temperature (*T*, K) and interconversion rate (*k*, s⁻¹) between the two isoenergetic states are also given for every trace. Simulated spectra were obtained with the Topspin 2.0 program.¹⁰

the phenylene blades of the penttiptycene rotator prevent a direct spectral simulation. Thus, a VT ¹³C NMR experiment was carried out, and the resulting spectra (Figure 3b) were simulated (Figure 3c).¹⁰ The results suggest that the rotation is nearly blocked at 298 K, and the rate constant (*k*) for interconversion between the two isoenergetic conformers of *cis*-**1** is only 6 s⁻¹. In other words, it takes ca. 0.33 s for a 360° rotation (i.e., *k*_{rot} = *k*/2). The activation energy (*E*_a = 14.8 ± 0.5 kcal mol⁻¹) and enthalpic ($\Delta H^\ddagger = 14.1 \pm 0.5$ kcal mol⁻¹) and entropic ($\Delta S^\ddagger = -7.6 \pm 1.4$ cal K⁻¹ mol⁻¹) contributions to the free energy of activation ($\Delta G^\ddagger = 16.4$ and 16.8 kcal mol⁻¹ at 298 and 348 K, respectively) were obtained by Arrhenius and Eyring plots (Supporting Information, Figures S12 and S13). Since the rotational barrier is mainly due to an enthalpic factor, the transition state is destabilized mainly due to steric effect. It should also be noted that the values of ΔG^\ddagger at 348 K deduced from ¹H and ¹³C VT NMR are essentially the same.

To obtain structural information about the transition structure and the optimized conformation of *cis*-**1** along the penttiptycene rotation coordinates, we have carried out DFT (BMK/6-311+G**//B3LYP/6-31G*) calculations.¹¹ The calculation results are justified by the good agreement of the calculated (16.75 kcal mol⁻¹) and the NMR-determined ΔG^\ddagger value (16.4 kcal mol⁻¹) at 298 K. The brake moiety in the structurally optimized *cis*-**1** (Figure 4a) is, as expected, located at the U-shaped cavities, although the protruding penttiptycene blades create two types of cavities (U- vs V-shaped). The V-shaped cavities are in fact inaccessible to

the dinitrophenyl group as a result of severe steric interactions with H₁, H₁', and the bridgehead hydrogen atom, as revealed by the transition structure (Figure 4b). Strains in bond angles




Figure 4. DFT-derived structures for *cis*-**1**: (a) the optimized conformation and (b) the transition structure along the penttiptycene rotation coordinate.

and bond lengths also account in part for the high energy of the transition state. The structural coordinates are supplied as Supporting Information.

An attempt to evaluate the rotational barriers for the penttiptycene rotator in *trans*-**1** and the dinitrophenyl brake in *cis*-**1** with VT ¹H NMR was hampered by their low energy barriers, as we could not observe decoalescence of the signals even at a temperature as low as 183 K (in CD₂Cl₂). In view of the good agreement of DFT calculations and NMR experiments in the penttiptycene rotation barrier of *cis*-**1**, DFT calculations have been applied to retrieve the corresponding information for that in *trans*-**1** (4.45 kcal mol⁻¹)¹² and the brake rotation in *cis*-**1** (6.85 kcal mol⁻¹). With a calculated ΔG^\ddagger value differing by 12.3 kcal mol⁻¹ for the penttiptycene

(10) All line shape simulations were performed using the Bruker Topspin 2.0 program.

(11) To expedite the calculations, the octyl group was replaced by a methyl group. The BMK functional has been shown to be a good functional in terms of thermochemistry and kinetics for main-group elements. Zhao, Y.; Truhlar, D. G. *Acc. Chem. Res.* **2008**, *41*, 157–167.

rotation in *trans*-**1** versus *cis*-**1** at 298 K, the difference in rotation rate is in the order of 10^9 .¹³

Photoswitching between *trans*-**1** and *cis*-**1** has been investigated in dichloromethane solutions.^{14,15} On the basis of their relative molar absorptivity (ϵ), the wavelengths of 306 ($\epsilon_{trans-1}/\epsilon_{cis-1} = 3.35$) and 254 nm ($\epsilon_{trans-1}/\epsilon_{cis-1} = 1.20$) were adopted for the *trans*-**1** \rightarrow *cis*-**1** and the *cis*-**1** \rightarrow *trans*-**1** conversion, respectively. The corresponding ratio of [*trans*-**1**]/[*cis*-**1**] determined by ¹H NMR is 25/75 and 45/55 in the photostationary states. As monitored by absorption spectra (Figure 5), photoswitching between the two photostationary states is quite robust (Figure 5, inset).

In summary, the pentaptycene-derived stilbene **1** has been prepared and investigated as a photocontrollable molecular brake that functions at room temperature. Both experimental and computational results reveal that at 298 K rotation of the four-bladed pentaptycene (the rotator) is “free” in *trans*-**1** but is nearly blocked in *cis*-**1**. The brake-on (*cis*-**1**) and brake-off (*trans*-**1**) states differ by a rotation rate of $\sim 10^9$ -fold and can be interconverted through the ethylene *trans*–*cis* photoisomerization reactions. Studies on the related systems are in progress in our laboratory in order to gain insights into substituent effects on the brake performance.

(12) A barrier of ~ 1000 cm^{-1} (2.86 kcal mol^{-1}) was found for the rotation of a single phenyl ring in *trans*-stilbene and its 4,4'-disubstituted derivatives. Arp, Z.; Chiang, W.-Y.; Laane, J.; Sakamoto, A.; Tasumi, M. *J. Phys. Chem. A* **2002**, *106*, 3479–3484.

(13) Even with a consideration of 1–2 kcal/mol uncertainty in calculated ΔG^\ddagger value for pentaptycene rotation in *trans*-**1**, the ratio of the on-off rotation rate is still around 10^7 – 10^8 .

(14) The quantum yields for the *trans*-**1** \rightarrow *cis*-**1** and *cis*-**1** \rightarrow *trans*-**1** photoisomerization in dichloromethane are 0.17 and 0.21, respectively.

(15) Thermal isomerization of the double bond in **1** has been evaluated by DFT (UBMK/6-311+G**//UB3LYP/6-31G*) modeling. The calculated free energy barrier was 45.22 kcal mol^{-1} relative to *cis*-**1**, and the free energy difference between *cis*-**1** and *trans*-**1** was 0.76 kcal mol^{-1} . These values are similar to the computational and experimental data reported for the parent stilbene (41–46 kcal mol^{-1} for the thermal isomerization barrier and 2.3–5.7 kcal mol^{-1} for the energy difference between the *cis* and *trans* isomers), see: (a) Meier, H. *Angew. Chem., Int. Ed.* **1992**, *31*, 1399–1420. (b) Han, W.-G.; Lovell, T.; Liu, T.; Noodleman, L. *ChemPhysChem* **2002**, *3*, 167–178.




Figure 5. Absorption spectra of *trans*-**1** (curve a) and *cis*-**1** (curve d) and their photostationary states irradiated with alternating 306-nm (curves c) and 254-nm (curves b) UV light irradiation in dichloromethane. Inset shows the changes in absorbance at 322 nm starting from *trans*-**1** (10 μM) for 7 switching cycles.

Acknowledgment. Financial support for this work was provided by the National Science Council, Academia Sinica, and National Taiwan University. The computing time granted by the National Center for High-Performance Computing and the Computing Center of Academia Sinica is acknowledged. We also thank Mr. G.-J. Huang for measuring the photoisomerization quantum yields.

Supporting Information Available: Experimental methods, synthesis of intermediates **3** and **4**, 1D and 2D NMR spectra and DFT-derived structural coordinates for *trans*-**1** and *cis*-**1**, and the Arrhenius and Eyring plots for *cis*-**1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL800689A