

Multisignal Chemosensor for Cr³⁺ and Its Application in Bioimaging


Kewei Huang, Hong Yang, Zhiguo Zhou, Mengxiao Yu, Fuyou Li,* Xia Gao, Tao Yi, and Chunhui Huang

Department of Chemistry & Laboratory of Advanced Materials, Fudan University, Shanghai 200433, People's Republic of China

fyli@fudan.edu.cn

Received April 11, 2008

ABSTRACT



A multisignal sensor based on rhodamine B with a ferrocene substituent (FD7) has been developed as a highly sensitive chemosensor for Cr³⁺. A prominent fluorescence enhancement was found in the presence of Cr³⁺, which was accompanied by changes in the absorption spectrum and the electrochemical parameters. Furthermore, by means of confocal laser scanning microscopy experiments, it has been demonstrated that FD7 can be used as a fluorescent probe for monitoring Cr³⁺ in living cells.

Trivalent chromium, Cr³⁺, is involved in some biochemical processes at the cellular level.¹ The mechanism by which Cr³⁺ affects human metabolism is based on modulation of the action of insulin through glucose tolerance factors (GTF), thereby activating certain enzymes and stabilizing proteins and nucleic acids.² A deficiency of chromium in the human body would lead to a variety of diseases, including diabetes and cardiovascular disease. Recently, Cr(III) has also been shown to adversely affect cellular structures, although its *in vivo* toxicity observed is much lower than that of Cr(VI).² To date, however, few methods have been developed whereby intracellular Cr(III) levels may be monitored or its distribution in living systems mapped, although organically bound Cr(III) derivatives might be transported across cell membranes by some as yet unknown mechanism.³ Accordingly, it is of necessity to be able to monitor the intracellular Cr³⁺ with the help of some effective sensors.

In this regard, by virtue of its highly sensitive and high-speed spatial analysis of cells, fluorescence bioimaging with

selective chemosensors offers a potentially powerful approach for detecting the transformation and distribution of low molecular-weight analytes (especially metal cations) in living samples.⁴ Such reagents have facilitated the bioimaging Ca²⁺,⁵ Zn²⁺,⁶ Hg²⁺,⁷ Cu⁺,⁸ and Pb²⁺ ions in living cells, but analogous tools for intracellular Cr³⁺ remain underdeveloped owing to the fluorescence quenching of paramagnetic Cr³⁺ and the lack of a selective ligand system. To the best of our knowledge, only two examples of Cr³⁺-

(3) (a) Raspor, P.; Batic, M.; Jamnik, P.; Josic, D.; Milacic, R.; Pas, M.; Recek, M.; Rezac-Dereani, V.; Skrt, M. *Acta Microbiol. Immunol. Hung.* **2000**, *47*, 143–73.

(4) For example, see: (a) Komatsu, H.; Miki, T.; Citterio, D.; Kubota, T.; Shindo, Y.; Kitamura, Y.; Oka, K.; Suzuki, K. *J. Am. Chem. Soc.* **2005**, *127*, 10798–10799. (b) Rosario, M.; Matt, C.; Stifun, M.; Mark, B. *Angew. Chem., Int. Ed.* **2006**, *45*, 5472–5474. (c) Farruggia, G.; Lotti, S.; Montalti, M.; Zaccaroni, N.; Savage, P. B.; Trapani, V.; Sale, P.; Wolf, F. *J. Am. Chem. Soc.* **2006**, *128*, 344–350. (d) Komatsu, H.; Iwasawa, N.; Citterio, D.; Suzuki, Y.; Kubota, T.; Tokuno, K.; Kitamura, Y.; Kotaro, O.; Suzuki, K. *J. Am. Chem. Soc.* **2004**, *126*, 16353–16360. (e) Chang, C. J.; Jaworski, J.; Nolan, E. M.; Sheng, M.; Lippard, S. J. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 1129–1134. (f) Kawabata, E.; Kikuchi, K.; Urano, Y.; Kojima, H.; Odani, A.; Nagano, T. *J. Am. Soc. Chem.* **2005**, *127*, 818–819. (g) Komatsu, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagano, T. *J. Am. Chem. Soc.* **2005**, *127*, 10197–10204. (h) Zhang, M.; Yu, M. X.; Li, F. Y.; Zhu, M. W.; Li, M. Y.; Gao, Y. H.; Li, L.; Liu, Z. Q.; Zhang, J. P.; Zhang, D. Q.; Yi, T.; Huang, C. H. *J. Am. Chem. Soc.* **2007**, *129*, 10322–10323.

(5) Tsien, R. W.; Tsien, R. Y. *Annu. Rev. Cell Biol.* **1990**, *6*, 715–760.

(1) Latva, S.; Jokiniemi, J.; Peraniemi, S.; Ahlgren, M. *J. Anal. At. Spectrom.* **2003**, *18*, 84–85.

(2) Singh, A. K.; Gupta, V. K.; Gupta, B. *Anal. Chim. Acta* **2007**, *585*, 171–178.

selective fluorescent chemosensors have been reported. One was reported by Samanta et al., who used di(2-ethylsulfanyl)ethylamine as a receptor moiety in organic solvents.¹⁰ In the other case, Liu et al. synthesized two fluorescent chemosensors capable of discriminating Fe³⁺ and Cr³⁺.¹¹ However, these Cr³⁺-sensors only functioned in organic solvent and thus needed to be further developed if they are to be applied in bioimaging.

Owing to their excellent spectroscopic properties of visible wavelength excitation and emission profiles, large molar extinction coefficients, and high fluorescence quantum yields, rhodamine derivatives have been utilized as important fluorescent probes. On the basis of the spiroactam (non-fluorescence) to ring-open amide (fluorescence) equilibrium of rhodamine, several rhodamine derivatives have been successfully fabricated as the OFF–ON-type fluorescent chemosensors for Cu²⁺,¹² Pb²⁺,¹³ Fe³⁺,^{14,15} and Hg²⁺.^{16,17} Recently, we have reported that a rhodamine B derivative (**1**, Figure 1) attached an 8-hydroxyquinoline moiety with




Figure 1. Chemical structures of **1** and **FD7**.

δ -bond could be used as a multisignaling chemosensor for Hg²⁺.^{17b} In the present study, an 8-hydroxyquinoline group

(6) (a) Lim, N. C.; Freaake, H. C.; Brückner, C. *Chem. Eur. J.* **2005**, *11*, 38–49. (b) Kikuchi, K.; Komatsu, K.; Nagano, T. *Curr. Opin. Chem. Biol.* **2004**, *8*, 182–191. (c) Burdette, S. C.; Lippard, S. J. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3605–3610.

(7) (a) Yoon, S.; Miller, E. W.; He, Q. W.; Do, P. H.; Chang, C. J. *Angew. Chem., Int. Ed.* **2007**, *46*, 6658–6661. (b) Yoon, S.; Albers, A. E.; Wong, A. P.; Chang, C. J. *J. Am. Chem. Soc.* **2005**, *127*, 16030–16031.

(8) Zeng, L.; Miller, E. W.; Pralle, A.; Isacoff, E. Y.; Chang, C. J. *J. Am. Chem. Soc.* **2006**, *128*, 10–11.

(9) He, Q. W.; Miller, E. W.; Wong, A. P.; Chang, C. J. *J. Am. Chem. Soc.* **2006**, *128*, 9316–9317.

(10) Sarkar, M.; Banthia, S.; Samanta, A. *Tetrahedron Lett.* **2006**, *47*, 7575–7578.

(11) Mao, J.; Wang, L.; Dou, W.; Tang, X.; Yan, Y.; Liu, W. *Org. Lett.* **2007**, *9*, 4567–4570.

(12) (a) Dujols, V.; Ford, F.; Czarnik, A. W. *J. Am. Chem. Soc.* **1997**, *119*, 7386–7387. (b) Xiang, Y.; Tong, A.; Jin, P.; Ju, Y. *Org. Lett.* **2006**, *8*, 2863–2866.

(13) Kwon, J. Y.; Jang, Y. J.; Lee, Y. J.; Kim, K. M.; Seo, M. S.; Nam, W.; Yoon, J. *J. Am. Chem. Soc.* **2005**, *127*, 10107–10111.

(14) Xiang, Y.; Tong, A. *Org. Lett.* **2006**, *8*, 1549–1552.

(15) Zhang, M.; Gao, Y. H.; Yu, M. X.; Li, F. Y.; Li, L.; Zhu, M. W.; Zhang, J. P.; Yi, T.; Huang, C. H. *Tetrahedron Lett.* **2007**, *21*, 3709–3712.

(16) (a) Wu, J. S.; Hwang, I. C.; Kim, K. S.; Kim, J. S. *Org. Lett.* **2007**, *9*, 907–910. (b) Wu, D.; Huang, W.; Duan, C.; Lin, Z.; Meng, Q. *Inorg. Chem.* **2007**, *46*, 1538–1540. (c) Lee, M. H.; Wu, J. S.; Lee, J. W.; Jung, J. H.; Kim, J. S. *Org. Lett.* **2007**, *9*, 2501–2504. (d) Song, K. C.; Kim, J. S.; Park, S. M.; Chung, K. C.; Ahn, S.; Chang, S. K. *Org. Lett.* **2006**, *8*, 3413–3416. (e) Zheng, H.; Qian, Z. H.; Xu, L.; Yuan, F. F.; Lan, L. D.; Xu, J. G. *Org. Lett.* **2006**, *8*, 859–861. (f) Yang, Y. K.; Yook, K. J.; Tae, J. *J. Am. Chem. Soc.* **2005**, *127*, 16760–16761. (g) Wu, D. Y.; Huang, W.; Duan, C. Y.; Lin, Z. H.; Meng, Q. G. *Inorg. Chem.* **2007**, *46*, 1538–1540.

(17) (a) Ko, S. K.; Yang, Y. K.; Tae, J.; Shin, I. *J. Am. Chem. Soc.* **2006**, *128*, 14150–14155. (b) Yang, H.; Zhou, Z. G.; Huang, K. W.; Yu, M. X.; Li, F. Y.; Yi, T.; Huang, C. H. *Org. Lett.* **2007**, *9*, 4729–4732.

as a conjugated moiety was further attached to the rhodamine derivative **FD7** (Figure 1). Interestingly, **FD7** could selectively recognize Cr³⁺ over other metal cations (including Hg²⁺), with suitably large variations in absorption, fluorescence emission, and electrochemical parameters. Further fluorescence bioimaging investigations have indicated that **FD7** may be used as a fluorescent probe for monitoring Cr³⁺ in living cells.

The rhodamine derivative **FD7** (Figure 1) was prepared from rhodamine B hydrazide by a two-step procedure in a yield of 60%. The structures of **FD7** and other intermediate products were confirmed by ¹H NMR, ¹³C NMR, and MS data (Supporting Information).

For practical application, the appropriate pH conditions for successful operation of the sensor were evaluated. Ring opening of the rhodamine occurred under acidic conditions (pH < 5.0) as a result of protonation.¹⁴ At pH > 5.0, no obvious ring opening was observed. However, the addition of Cr³⁺ led to the fluorescence enhancement over a comparatively wide pH range (5.0–10.0), which is attributed to opening of the rhodamine ring (Figure S1, Supporting Information). Consequently, **FD7** may be used to detect Cr³⁺ over a wide pH range since there is a Cr³⁺-induced OFF–ON fluorescence change throughout this range. A similar result was found by examining the UV/vis spectra rather than fluorescence over the same pH range (Figure S2, Supporting Information).

An optimized solvent system of ethanol/H₂O (1:1, v/v, pH 7.4) solution was selected for the spectroscopic investigations. Upon binding with Cr³⁺, the absorption peak of **FD7** (20 μ M) at 565 nm in ethanol/H₂O (1:1, v/v, pH 7.4) solution increased evidently. The colorless solution gradually turned to pink, clearly indicating the ring-opening process of the rhodamine B unit in **FD7**. The association constant for Cr³⁺ was estimated to be $7.5 \times 10^3 \text{ M}^{-1}$ in the ethanol/H₂O (1:1, v/v, pH 7.4) solution on the basis of nonlinear fitting of the titration curve assuming a 1:1 stoichiometry (Figure 2 inset).




Figure 2. Changes in UV–vis absorption spectra of **FD7** (20 μ M) in ethanol/H₂O (1:1, v/v, pH 7.4) solutions with various amounts of Cr³⁺ ions (0–45 equiv). Inset: Absorbance at 565 nm of **FD7** as a function of Cr³⁺ concentration.

As designed, **FD7** shows a distinct current change of the electric current in its reversible ferrocene/ferrocinium redox cycles upon complexation. Differential pulse voltammetry (DPV) curves of **FD7** were recorded in ethanol solution containing 0.1 M *n*-tetrabutylammonium hexafluorophosphate (*n*-Bu₄NPF₆) as supporting electrolyte in the absence and presence of Cr³⁺. As shown in Figure 3, a significant




Figure 3. DPV of **FD7** (100 μM) in ethanol solution in the absence and presence of 1.6 equiv of Cr³⁺ with *n*-Bu₄NPF₆ as supporting electrolyte.

displacement was observed upon addition of Cr³⁺. The peak was cathodically shifted upon complexation ($\Delta E_{1/2} = 140$ mV) in ethanol. This fact indicated that **FD7**, like other chemosensors with ferrocene units,¹⁸ had the potential to be a redox chemosensor of Cr³⁺.

In investigating the fluorescence, the spectrum of **FD7** (20 μM) in ethanol/H₂O (1:1, v/v, pH 7.4) showed a weak emission at around 580 nm ($\lambda_{\text{ex}} = 530$ nm), since the spirocyclic form of rhodamine prevailed.¹⁴ Upon addition of Cr³⁺, an emission band peaked at 587 nm attributable to delocalization in the xantheno moiety of the rhodamine significantly increased in intensity. The solution showed an intense red fluorescence (Figure 4), with a quantum yield of 0.18 and an approximately 15-fold enhancement in the fluorescence intensity at 587 nm. This fact means that **FD7** could be used as an off–on fluorescent chemosensor for Cr³⁺.

To obtain an excellent chemosensor, high selectivity is essential. To validate the selectivity of **FD7** in practice, some other metal ions including alkali, alkaline earth, and transition-metal ions were added into a solution of **FD7** under the same conditions (ethanol/H₂O, 1:1, v/v, pH 7.4). Only Hg²⁺ elicited a slight fluorescence intensity enhancement, while the other metal ions did not cause any discernible changes (Figure 5). Moreover, the enhancement of the fluorescence intensity resulting from the addition of Cr³⁺ was not influenced by subsequent addition of other metal ions. All of these results indicate the high selectivity of **FD7** for Cr³⁺ over other competing cations in ethanol/H₂O (1:1, v/v, pH 7.4) solution.

(18) (a) Caballero, A.; Martínez, R.; Lloveras, V.; Ratera, I.; Vidal-Gancedo, J.; Wurst, K.; Tárraga, A.; Molina, P.; Veciana, J. *J. Am. Chem. Soc.* **2005**, *127*, 15666–15667. (b) Caballero, A.; Lloveras, V.; Tárraga, A.; Espinosa, A.; Velasco, M. D.; Vidal-Gancedo, J.; Rovira, C.; Wurst, K.; Molina, P.; Veciana, J. *Angew. Chem., Int. Ed.* **2005**, *44*, 1977–1981.




Figure 4. Fluorescence emission changes of **FD7** (20 μM) in ethanol/H₂O (1:1, v/v, pH 7.4) solutions upon addition of Cr³⁺ (0–45 equiv). Each spectrum was acquired 1 min after Cr³⁺ addition, $\lambda_{\text{ex}} = 530$ nm. Fluorescence photograph was obtained under irradiation of UV light (365 nm).

The probable complexation mechanism of **FD7** with Cr³⁺ was validated by XPS and ESI-MS. The XPS of **FD7**-Cr indicated the sole presence of Cr³⁺ during this process, that is, no redox process but only complexation, albeit with the rather electron-with rhodamine B hydrazide moiety generally behaving as an electron donor (Figure S8, Supporting Information).¹⁹ Moreover, this binding mode is supported





Figure 5. Fluorescence responses of **FD7** (20 μM) to various metal ions (5 mM for Ca²⁺, Mg²⁺, Na⁺, K⁺, and Zn²⁺; 900 μM for all other cations) in ethanol/H₂O (1:1, v/v, pH 7.4). Bars represent the final integrated fluorescence response (F_f) over the initial integrated emission (F_i). White bars represent the addition of metal ions to a 20 μM solution of **FD7**. Black bars represent the addition of Cr³⁺ (45 equiv) to the solution. ($\lambda_{\text{ex}} = 530$ nm).

by the presence of a peak at m/z 962.0 (calcd 962.7) corresponding to $[\text{FD7} + \text{CrCl}_2]^+$ in the ESI-MS spectrum of a mixture of **FD7** and 200 μM CrCl₃ (Figure S9, Supporting Information). Thus, in accordance with the

(19) Logar, N. Z.; Siljeg, M.; Arcon, I.; Meden, A.; Tusar, N. N.; Stefanovic, S. C.; Kovac, J.; Kaucic, V. *Microporous Mesoporous Mater.* **2006**, *93*, 275–284.

coordination number of 6 of Cr^{3+} , the most likely binding sites for Cr^{3+} are the conjugated moiety including carbonyl O, imino N, and quinoline N and O atoms of **FD7** (Scheme 1).

Scheme 1. Probable Complexation Mechanism of **FD7** with Cr^{3+}



To test the sensitivity of the interaction between **FD7** and Cr^{3+} , we also investigated the time course of the response of **FD7** ($20\ \mu\text{M}$) in the presence of 45 equiv of Cr^{3+} in ethanol/ H_2O (1:1, v/v, pH 7.4). It was found that the obvious spectral change was observed within 1 min (Figure S5 and movie, Supporting Information) upon addition of 45 equiv Cr^{3+} . Therefore, this system is suitably predisposed for the detection of Cr^{3+} .

In view of its favorable spectroscopic properties and the instantaneous interaction with Cr^{3+} , **FD7** should be well-suited for fluorescence imaging in living cells. As determined by laser scanning confocal microscopy, staining HeLa cells with a $20\ \mu\text{M}$ solution of **FD7** in ethanol/PBS (1:99, v/v) buffer for 30 min at $25\ ^\circ\text{C}$ led to very faint intracellular fluorescence (Figure 6a1). The cells were supplemented with 0, 5, 20, and $100\ \mu\text{M}$ CrCl_3 in the growth medium for 2.5 h at $37\ ^\circ\text{C}$ and then loaded with **FD7** under the same conditions, whereupon a significant increase in the fluorescence from the intracellular area was observed (Figure 6a1–d1). Moreover, after incubation with $50\ \mu\text{M}$ CrCl_3 , an obvious increase in the intracellular Cr(III) concentration was observed by ICP-AEC (Supporting Information). Bright-field measurements after treatment with Cr^{3+} and **FD7** confirmed that the cells remained viable throughout the imaging experiments (Figure 6a2–d2). As depicted in Figure 6b3–d3, the overlay of fluorescence and bright-field images revealed that the fluorescence signals were localized in the perinuclear




Figure 6. Confocal fluorescence, brightfield, and overlay images of HeLa cells. The cells were supplemented with $0\ \mu\text{M}$ (a1–a3), $5\ \mu\text{M}$ (b1–b3), $20\ \mu\text{M}$ (c1–c3), or $100\ \mu\text{M}$ (d1–d3) CrCl_3 in the growth media for 2.5 h at $37\ ^\circ\text{C}$ and then were stained with $20\ \mu\text{M}$ **FD7** for 30 min at $37\ ^\circ\text{C}$ ($\lambda_{\text{ex}} = 543\ \text{nm}$). e1–e3 were reference cells without Cr^{3+} .

area of the cytosol, indicating a subcellular distribution of Cr^{3+} . Taken together, the results indicate that the **FD7** is cell-permeable and can respond to changes in intracellular Cr^{3+} concentration within living cells.

In summary, we have utilized the ring-opening of a rhodamine B derivative and the electrochemical characterization of a ferrocenyl group to develop a novel multisignal probe for detecting the transition-metal ion Cr^{3+} in ethanol/ H_2O (1:1, v/v, pH 7.4) solution. The new fluorescent sensor showed an excellent selectivity for Cr^{3+} over other metal ions examined in similar solutions. Confocal laser scanning microscopy experiments have proven that **FD7** can be used to monitor Cr^{3+} in living cells and to map its subcellular distribution.

Acknowledgment. We are thankful for the financial support from NSFC (20490210 and 20775017), NHTPC (2006AA03Z318), SSTC (06QH14002), Shanghai Leading Academic Discipline Project (B108), and the China Post-doctoral Science Foundation.

Supporting Information Available: Synthetic details, NMR spectra, and additional spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL800778A