

Direct Asymmetric Allylic Alkylation of Butenolides with Morita–Baylis–Hillman Carbonates


Hai-Lei Cui,[†] Ji-Rong Huang,[†] Jie Lei,[†] Zhao-Feng Wang,[†] Shi Chen,[†] Li Wu,[†] and Ying-Chun Chen^{*†‡}

Key Laboratory of Drug-Targeting and Drug Delivery System of Education Ministry, Department of Medicinal Chemistry, West China School of Pharmacy, and State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu 610041, China

ycchenhuaxi@yahoo.com.cn

Received December 1, 2009

ABSTRACT



The direct asymmetric allylic alkylation of β,γ -butenolides with MBH carbonates to access γ,γ -disubstituted butenolides containing adjacent quaternary and tertiary chiral centers has been presented in excellent stereoselectivities (86–96% ee, dr >95:5) and moderate to good yield (50–83%). Their synthetic utility has been well demonstrated by the facile construction of bicyclic lactones bearing 4–5 stereogenic centers.

The widespread occurrence of the butenolide unit in natural products and pharmaceutically useful molecules has long stimulated efforts on its synthesis and transformations.¹ Thus, the development of enantioselective protocols to access functionalized butenolides has triggered increasing interest in this regard.² Most literature procedures have been focused on the

Mukaiyama-type additions that were limited to the applications of preformed silyloxyfurans.³ However, in view of atom economy, the direct modification of butenolides represents a more attractive approach, which enables synthetic convenience and efficiency. Shibasaki^{4a} and Trost^{4b} reported their pioneering work in the field of direct enantioselective functionalization of butenolides, although only simple butenolides have been tested. Recently Buchwald and co-workers presented the synthesis of γ,γ -disubstituted butenolides involving the construction of a quaternary carbon center via direct γ -arylation of γ -substituted butenolides.⁵ Nevertheless, to the best of our knowledge, no

[†] West China School of Pharmacy.

[‡] West China Hospital.

(1) For a review on natural products containing the butenolide unit, see: (a) Alali, F. Q.; Liu, X.-X.; McLaughlin, J. L. *J. Nat. Prod.* **1999**, *62*, 504. For reviews on the synthesis of butenolides, see: (b) Rao, Y. S. *Chem. Rev.* **1964**, *64*, 353. (c) Negishi, E.; Kotori, M. *Tetrahedron* **1997**, *53*, 6707. (d) Montagnon, T.; Tofi, M.; Vassilikogiannakis, G. *Acc. Chem. Res.* **2008**, *41*, 1001.

(2) For selected examples, see: (a) Movassaghi, M.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2002**, *124*, 2456. (b) Adrio, J.; Carretero, J. C. *J. Am. Chem. Soc.* **2007**, *129*, 778. (c) He, Y.-T.; Yang, H.-N.; Yao, Z.-J. *Tetrahedron* **2002**, *58*, 8805. (d) Jauch, J. *Eur. J. Org. Chem.* **2001**, 473. (e) Grossmann, G.; Poncioni, M.; Bornand, M.; Jolivet, B.; Neuburger, M.; Séquin, U. *Tetrahedron* **2003**, *59*, 3237. (f) Patil, S. N.; Liu, F. *J. Org. Chem.* **2008**, *73*, 4476.

(3) For reviews, see: (a) Casiraghi, G.; Zanardi, F.; Appendino, G.; Rassu, G. *Chem. Rev.* **2000**, *100*, 1929. (b) Denmark, S. E.; Heemstra, J. R., Jr.; Beutner, G. L. *Angew. Chem., Int. Ed.* **2005**, *44*, 4682.


(4) (a) Yamaguchi, A.; Matsunaga, S.; Shibasaki, M. *Org. Lett.* **2008**, *10*, 2319. (b) Trost, B. M.; Hitce, J. *J. Am. Chem. Soc.* **2009**, *131*, 4572.

(5) Hyde, A. M.; Buchwald, S. L. *Org. Lett.* **2009**, *11*, 2663.

report has yet been published on the direct asymmetric variant of these easily prepared but useful γ -substituted butenolides.⁶

Recently, the allylic alkylation with Morita–Baylis–Hillman (MBH) adducts by the catalysis of metal-free Lewis basic tertiary amines or phosphines has emerged as a powerful strategy to deliver multifunctional compounds.^{7,8} Krische^{9a} reported the first phosphine-catalyzed allylic alkylation of 2-silyloxyfuran with MBH acetates in excellent regio- and diastereoselectivities. Later Shi^{9b} developed the first asymmetric variant to afford γ -substituted butenolides by employing newly designed chiral bifunctional phosphine catalysts. However, limitation of substrate to preformed 2-silyloxyfuran without any substitution incapacitated access to enantio-enriched γ,γ -disubstituted butenolides. We envisioned that, as outlined in Scheme 1, the direct tertiary amine-catalyzed

Scheme 1. Proposed Direct Allylic Alkylation of γ -Substituted Butenolides with MBH Carbonates



asymmetric γ -allylic alkylation of γ -substituted butenolides would be realized via deprotonation by an *in situ* generated *tert*-butoxy anion and subsequent vinylogous addition reaction. We wondered whether the chemo-, regio-, and stereoselectivities could be well accomplished simultaneously, although the rather challenging construction of adjacent quaternary and tertiary chiral centers must be fulfilled.^{10,11}

We began our investigation with the reaction of β,γ -butenolide **1a** bearing a γ -aryl substitution with MBH carbonate


(6) For catalytic asymmetric synthesis of γ,γ -disubstituted butenolides, see: (a) Brown, S. P.; Goodwin, N. C.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2003**, *125*, 1192. (b) Shaw, S. A.; Aleman, P.; Christy, J.; Kampf, J. W.; Va, P.; Vedejs, E. *J. Am. Chem. Soc.* **2006**, *128*, 925.

(7) (a) Kim, J. N.; Lee, H. J.; Gong, J. H. *Tetrahedron Lett.* **2002**, *43*, 9141. (b) Cho, C.-W.; Kong, J.-R.; Krische, M. J. *Org. Lett.* **2004**, *6*, 1337. (c) Du, Y.; Han, X.; Lu, X. *Tetrahedron Lett.* **2004**, *45*, 4967. (d) Zhang, T.-Z.; Dai, L.-X.; Hou, X.-L. *Tetrahedron: Asymmetry* **2007**, *18*, 1990. (e) van Steenis, D. J. V. C.; Marcelli, T.; Lutz, M.; Spek, A. L.; van Maarseveen, J. H.; Hiemstra, H. *Adv. Synth. Catal.* **2007**, *349*, 281. (f) Park, H.; Cho, C.-W.; Krische, M. J. *J. Org. Chem.* **2006**, *71*, 7892. (g) Ma, G.-N.; Cao, S.-H.; Shi, M. *Tetrahedron: Asymmetry* **2009**, *20*, 1086.

(8) For studies from this group, see: (a) Cui, H.-L.; Peng, J.; Feng, X.; Du, W.; Jiang, K.; Chen, Y.-C. *Chem.—Eur. J.* **2009**, *15*, 1574. (b) Jiang, K.; Peng, J.; Cui, H.-L.; Chen, Y.-C. *Chem. Commun.* **2009**, 3955. (c) Cui, H.-L.; Feng, X.; Peng, J.; Jiang, K.; Chen, Y.-C. *Angew. Chem., Int. Ed.* **2009**, *48*, 5737. (d) Feng, X.; Yuan, Y.-Q.; Cui, H.-L.; Jiang, K.; Chen, Y.-C. *Org. Biomol. Chem.* **2009**, *7*, 3660. (e) Zhang, S.-J.; Cui, H.-L.; Jiang, K.; Li, R.; Ding, Z.-Y.; Chen, Y.-C. *Eur. J. Org. Chem.* **2009**, 5804.

2a in the presence of DABCO (20 mol %) in DCE at ambient temperature. To our delight, the reaction proceeded very smoothly, and the regioselective γ,γ -disubstituted butenolide **3a** was obtained with excellent diastereoselectivity (>95:5)¹² in 88% yield after 1 h, although some unidentified byproduct was also observed (Table 1, entry 1). We then examined the

Table 1. Screening Studies of Direct Asymmetric Allylic Alkylation of β,γ -Butenolides **1a** with MBH Carbonate **2a**^a



entry	catalyst ^b	solvent	<i>t</i> (h)	yield ^c (%)	ee ^d (%)
1 ^e	DABCO	DCE	1	88	
2	(DHQD) ₂ AQN	DCE	19	61	83
3	(DHQD) ₂ PHAL	DCE	24	66	81
4	(DHQD) ₂ PYR	DCE	18	90	85
5	(DHQ) ₂ AQN	DCE	24	35	-10
6	(DHQ) ₂ PYR	DCE	25	65	-35
7	(DHQD) ₂ PYR	toluene	32	60	91
8	(DHQD) ₂ PYR	PhCF ₃	22	78	92
9 ^f	(DHQD) ₂ PYR	PhCF ₃	47	62	93
10 ^g	(DHQD) ₂ PYR	PhCF ₃	54	54	92
11 ^h	(DHQD) ₂ PYR	PhCF ₃	10	82	92
12 ^h	(DHQ) ₂ PYR	PhCF ₃	19	45	-46

^a Unless otherwise noted, reactions were performed with 0.1 mmol of **1a**, 0.2 mmol of **2a**, and 10 mol % of catalyst in 1.0 mL of solvent at 50 °C. ^b DABCO: 1,4-diazabicyclo[2.2.2]octane. (DHQD)₂AQN: hydroquinidine (anthraquinone-1,4-diyl) diether. (DHQD)₂PHAL: hydroquinidine 1,4-phthalazinediyl diether. (DHQD)₂PYR: hydroquinidine-2,5-diphenyl-4,6-pyrimidinediyl diether. (DHQ)₂AQN: hydroquinineanthraquinone-1,4-diyl diether. (DHQ)₂PYR: hydroquinine-2,5-diphenyl-4,6-pyrimidinediyl diether. ^c Isolated yield of pure **3a** (dr >95:5).¹² ^d Determined by chiral HPLC analysis. ^e At rt, with 20 mol % of catalyst. ^f At 35 °C. ^g 10 mol % of S-BINOL was added. ^h In 0.5 mL of solvent.

asymmetric catalytic ability of some modified cinchona alkaloids at higher temperature (entries 2–6).¹³ The screening studies revealed that (DHQD)₂PYR was the best choice in terms

(9) (a) Cho, C.-W.; Krische, M. J. *Angew. Chem., Int. Ed.* **2004**, *43*, 6689. (b) Jiang, Y.-Q.; Shi, Y.-L.; Shi, M. *J. Am. Chem. Soc.* **2008**, *130*, 7202.

(10) For reviews on the construction of a quaternary carbon center, see: (a) Christoffers, J.; Baro, A. *Adv. Synth. Catal.* **2005**, *347*, 1473. (b) Cozzi, P. G.; Hilgraf, R.; Zimmermann, N. *Eur. J. Org. Chem.* **2007**, 5969. (c) Bella, M.; Gasperi, T. *Synthesis* **2009**, 1583.

(11) For selected examples of the construction of adjacent quaternary and tertiary chiral centers, see: (a) Austin, J. F.; Kim, S.-G.; Sinz, C. J.; Xiao, W.-J.; MacMillan, D. W. C. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5482. (b) Li, H.; Wang, Y.; Tang, L.; Wu, F.; Liu, X.; Guo, C.; Foxman, B. M.; Deng, L. *Angew. Chem., Int. Ed.* **2005**, *44*, 105. (c) Trost, B. M.; Zhang, Y. *J. Am. Chem. Soc.* **2007**, *129*, 14548. (d) Wilt, J. C.; Pink, M.; Johnston, J. N. *Chem. Commun.* **2008**, 4177. (e) Bui, T.; Syed, S.; Barbas, C. F., III *J. Am. Chem. Soc.* **2009**, *131*, 8758. (f) Kato, Y.; Furutachi, M.; Chen, Z.; Mitsunuma, H.; Matsunaga, S.; Shibasaki, M. *J. Am. Chem. Soc.* **2009**, *131*, 9168.


(12) The diastereomeric ratio of >95:5 indicates that the minor isomer could not be detected by ¹H NMR analysis. Excellent diastereoselectivities (generally >95:5) were reported in the similar reaction of 2-silyloxyfuran by Krische and Shi; see ref 9.

(13) For reviews, see: (a) Tian, S.-K.; Chen, Y.; Hang, J.; Tang, L.; McDaid, P.; Deng, L. *Acc. Chem. Res.* **2004**, *37*, 621. (b) Gaunt, M. J.; Johansson, C. C. C. *Chem. Rev.* **2007**, *107*, 5596.

of enantioselectivity and yield (entry 4). However, much inferior stereocontrol was attained in the presence of tertiary amines derived from DHQ, and the products possessed the opposite configuration (entries 5 and 6). Subsequently, the optimization of other reaction parameters (solvent, temperature, additive, concentration, etc.) was continued with (DHQD)₂PYR (entries 7–11). It was found that better results were gained when PhCF₃ was applied as the solvent (entry 8), but neither lowering the temperature nor adding 10 mol % of *S*-BINOL could improve the data (entries 9 and 10). Finally, the best results with respect to reaction efficacy, yield, and enantioselectivity were achieved by performing the reaction in a higher concentration, affording **3a** in 82% yield and 92% ee in 10 h (entry 11). Nevertheless, the enantioselectivity was still not satisfying when (DHQ)₂PYR was applied (entry 12).

Consequently, more experiments to examine substrate scope and limitations under the optimized conditions were conducted. The allylic alkylation products were generally isolated as a single diastereomer. As summarized in Table 2, excellent

Table 2. Direct Asymmetric Allylic Alkylation of β,γ -Butenolides **1** with MBH Carbonates **2**^a



entry	1	Ar	EWG	<i>t</i> (h)	yield ^b (%)	ee ^c (%)
1	1a	Ph	CO ₂ Me	10	3a , 82	92
2	1b	Ph	CO ₂ Me	24	3b , 60	91
3	1c	Ph	CO ₂ Me	14	3c , 76	92
4	1d	Ph	CO ₂ Me	18	3d , 67	80
5 ^d	1d	Ph	CO ₂ Me	18	3d , 83	88
6 ^{d,e}	1d	Ph	CO ₂ Me	68	3d , 70	88
7	1a	<i>p</i> -ClC ₆ H ₄	CO ₂ Me	19	3e , 66	92
8	1b	<i>m</i> -ClC ₆ H ₄	CO ₂ Me	12	3f , 68	90
9	1a	<i>o</i> -BrC ₆ H ₄	CO ₂ Me	25	3g , 61	94
10	1a	<i>p</i> -MeOC ₆ H ₄	CO ₂ Me	35	3h , 50	94
11	1a	<i>m</i> -MeC ₆ H ₄	CO ₂ Me	23	3i , 62	89 ^f
12	1a	2-thienyl	CO ₂ Me	18	3j , 68	88
13 ^g	1a	Ph	COCH ₃	19	3k , 57	96
14 ^g	1a	<i>p</i> -BrC ₆ H ₄	COCH ₃	26	3l , 67	86
15 ^{d,g}	1d	Ph	COCH ₃	18	3m , 68	93

^a Unless otherwise noted, reactions were performed with 0.1 mmol of **1**, 0.2 mmol of **2**, 10 mol % of (DHQD)₂PYR in 0.5 mL PhCF₃ at 50 °C. ^b Isolated yield of pure **3** (dr >95:5). ^c Based on chiral HPLC analysis. ^d With (DHQD)₂AQN in DCE. ^e At 1.0 mmol scale with 2 mol % of catalyst loading. ^f The absolute configuration of **3i** was determined by X-ray analysis (see Figure 1). The other products were assigned by analogy. ^g At 35 °C.

enantioselectivities and moderate to good yields (60–82%) were obtained for a few β,γ -butenolides **1a–1c** bearing different γ -aryl groups in the reaction with MBH carbonate **2a** (Table 2, entries 1–3). It was pleasing that a simple γ -methyl-substituted butenolide **1d** exhibited good reactivity (entry 4), and higher enantioselectivity and yield were obtained by the catalysis of (DHQD)₂AQN in DCE (entry 5). It was noteworthy that the reaction could be conducted at lower catalyst loading (2 mol %) without affecting the high enantioselectivity (entry

6). On the other hand, excellent enantiocontrol (88–94% ee) was attained for a diversity of MBH carbonates with various electron-deficient or -rich aryl or heteroaryl groups, generally in fair to good yields (50–68%) (entries 7–12). In addition, the MBH carbonates derived from methyl vinyl ketone (MVK) also led to the corresponding γ,γ -disubstituted butenolides in high enantioselectivities (entries 13–15). Unfortunately, a messy reaction was noted for MBH carbonates with a β -alkyl substitution.





Figure 1. ORTEP representation of the X-ray structure of enantiopure **3i** (thermal ellipsoids at 30% probability).

Moreover, we intended to expand the substrate scope to α,β -butenolides. We were pleased to find DABCO still could promote the highly regio- and diastereoselective allylic alkylation of α,β -butenolide **4a** and MBH carbonate **2a** to provide γ -substituted butenolide **5a** in 49% yield at 50 °C, although more side reactions were observed (Table 3, entry 1). Subse-

Table 3. Direct Asymmetric Allylic Alkylation of α,β -Butenolides **4** with MBH Carbonates **2**^a



4a R = Ph, R¹ = Me **4c** R = H, R¹ = Et
4b R = H, R¹ = *i*-Pr **4d** R = H, R¹ = H

entry	catalyst	4	solvent	<i>t</i> (h)	yield ^b (%)	ee ^c (%)
1	DABCO	4a	DCE	24	5a , 49	
2	(DHQD) ₂ PYR	4a	PhCF ₃	94	5a , 44	67
3	(DHQD) ₂ PYR	4a	toluene	43	5a , 18	68
4	(DHQD) ₂ PYR	4a	DCE	43	5a , 45	83
5	(DHQD) ₂ PHAL	4a	DCE	43	5a , 33	88
6	(DHQD) ₂ AQN	4a	DCE	43	5a , 57	90
7	(DHQD) ₂ AQN	4b	DCE	67	5b , 39	89
8	(DHQD) ₂ AQN	4c	DCE	40	5c , 38	90
9 ^d	(DHQD) ₂ AQN	4d	DCE	24	5d , 24	95

^a Unless otherwise noted, reactions were performed with 0.1 mmol of **4**, 0.2 mmol of **2a**, and 10 mol % of catalyst in 0.5 mL solvent. Entries 1–8: EWG = CO₂Me; entry 9: EWG = COMe ^b Isolated yield of pure **5** (dr >95:5). ^c Determined by chiral HPLC analysis. ^d With 0.3 mmol of **4d**, 0.1 mmol of **2a**.

quently, the asymmetric screening experiments were performed (entries 2–6), and high enantioselectivity could be obtained with catalysis by (DHQD)₂AQN in DCE (entry 6). Under the optimized conditions, β -substituted butenolides **4b** and **4c** afforded the allylic products with high ee values in fair yields (entries 7 and 8). The simple α,β -butenolide **4d** could react with MBH carbonate derived from MVK to give the adduct **5d** with an excellent ee value but in poor yield,^{9,14} due to self-dimerization and other complex side reactions (entry 9).^{1b}

We have further explored the transformations of the obtained butenolides with multiple functionalities in order to illustrate their synthetic versatility. As illustrated in Scheme 2, tandem double aza-Michael additions of benzyl-

alcohol **8** as a single isolable isomer with two quaternary chiral centers.¹⁸ We expect that these multifunctional compounds might have potentials in the design and synthesis of biologically active or natural product like materials.


In conclusion, we have developed the first direct asymmetric γ -allylic alkylation of butenolides with MBH carbonates promoted by modified cinchona alkaloids. This methodology exhibits high chemo-, regio-, and diastereoselectivities and allows access to γ,γ -disubstituted butenolides containing adjacent quaternary and tertiary stereogenic centers from β,γ -butenolides in excellent stereoselectivities (86–96% ee, dr >95:5) and moderate to good yields (50–83%). It could be extended to the direct asymmetric allylic alkylation of α,β -butenolides, affording similar adducts with high enantioselectivities, albeit in low to fair yields. The synthetic utility of this strategy has been demonstrated by the facile construction of bicyclic lactones containing 4–5 stereogenic centers. Their further application to the synthesis of biologically important molecules is underway in our laboratory.

Acknowledgment. We are grateful for financial support from the National Natural Science Foundation of China (20772084), PCSIRTC (IRT0846), and National Basic Research Program of China (973 Program) (2010CB833303).

Supporting Information Available: Experimental procedures, structural proofs, NMR spectra and HPLC chromatograms of the products, and CIF file of enantiopure **3i**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL100014M

Scheme 2. Synthetic Transformation of Multifunctional Butenolides



amine to **3d** were easily realized to provide bicyclic piperidine derivative **6** in excellent diastereocontrol (dr >95:5).¹⁵ In addition, the highly diastereoselective tandem Michael additions with nitromethane were also quite successful by the catalysis of tetramethylguanidine (TMG), giving bicyclic lactones **7a** and **7b** with five contiguous chiral centers.¹⁶ We investigated the intramolecular ketyl coupling reaction of **3m** via radical cyclization induced by tributyltin hydride.¹⁷ An interesting reductive Michael addition was observed at the ketone carbonyl carbon, affording bicyclic

(14) The obtained stereochemistry is consistent with that observed when catalysis by triphenylphosphine is employed; see ref 9a.

(15) For examples with related frameworks, see: (a) Lemire, A.; Charette, A. B. *Org. Lett.* **2005**, *7*, 2747. (b) Cassidy, M. P.; Padwa, A. *Org. Lett.* **2004**, *6*, 4029. (c) Wee, A. G. H.; Fan, G.-J. *Org. Lett.* **2008**, *10*, 3869. (d) Heretsch, P.; Rabe, S.; Giannis, A. *Org. Lett.* **2009**, *11*, 5410.

(16) For examples with related frameworks, see: (a) Fuchs, S.; Berl, V.; Lepoittevin, J.-P. *Eur. J. Org. Chem.* **2007**, 1145. (b) Maulide, N.; Markó, I. E. *Chem. Commun.* **2006**, 1200. (c) Zhang, W.; Romo, D. *J. Org. Chem.* **2007**, *72*, 8939. (d) Huber, R. S.; Jones, G. B. *J. Org. Chem.* **1992**, *57*, 5778.

(17) Enholm, E. J.; Kinter, K. S. *J. Org. Chem.* **1995**, *60*, 4850.

(18) For examples with related frameworks, see: (a) Nobitec, E.; Aniol, M.; Wawrzeńczyk, C. *Tetrahedron* **1994**, *50*, 10339. (b) Ghosh, A. K.; Takayama, J. *Tetrahedron Lett.* **2008**, *49*, 3409. (c) Tello-Aburto, R.; Rios, M. Y.; Swenson, D. C.; Olivo, H. F. *Tetrahedron Lett.* **2008**, *49*, 6853. (d) Brandvold, T. A.; Wulff, W. D.; Rheingold, A. L. *J. Am. Chem. Soc.* **1990**, *112*, 1645.