

Discovery and Total Synthesis of a New Estrogen Receptor Heterodimerizing Actinopolymorphol A from *Actinopolymorpha rutilus*


Sheng-Xiong Huang,[†] Emily Powell,[‡] Scott R. Rajski,[†] Li-Xing Zhao,[‡]
Cheng-Lin Jiang,[‡] Yanwen Duan,[#] Wei Xu,[‡] and Ben Shen^{*,†,§,||}

Division of Pharmaceutical Sciences, McArdle Laboratory for Cancer Research, University of Wisconsin National Cooperative Drug Discovery Group, and Department of Chemistry, University of Wisconsin–Madison, Madison, Wisconsin 53705, Yunnan Institute of Microbiology, Yunnan University, Kunming, Yunnan 650091, China, and Hunan Engineering Research Center of Combinatorial Biosynthesis and Natural Product Drug Discovery, Changsha, Hunan 410329, China

bshen@pharmacy.wisc.edu

Received June 11, 2010

ABSTRACT



Estrogen receptor ER α and ER β heterodimerization has been implicated in cancer chemoprevention. The discovery, structural elucidation, and total synthesis of a new natural product, actinopolymorphol A (1), from *Actinopolymorpha rutilus* (YIM45725) that preferentially induces ER α /ER β heterodimerization is reported. Total synthesis of 1 has allowed us to determine its absolute stereochemistry and that of a previously known deacetylated congener, and 1 represents the first member of a new class of natural products not previously recognized to modulate ER function.

The estrogen receptors (ERs) are ligand-inducible transcription factors implicated in a wide array of biological processes. Binding of 17 β -estradiol (E2) and other estrogenic ligands

triggers receptor dimerization and subsequent control of gene transcription. ERs exist in two forms, ER α and ER β , which have opposing roles in regulating estrogen action: ER α promotes, whereas ER β inhibits E2-dependent cell growth.¹ Notably, the coexpression of ER β with ER α leads to reduced proliferation of breast cancer cells and ~60% of all breast tumors express both receptors.² Coexpression of ER β has

[†] Division of Pharmaceutical Sciences, University of Wisconsin–Madison.

[‡] McArdle Laboratory for Cancer Research, University of Wisconsin–Madison.

[‡] Yunnan University.

[#] Hunan Engineering Research Center of Combinatorial Biosynthesis and Natural Product Drug Discovery.

[§] University of Wisconsin National Cooperative Drug Discovery Group, University of Wisconsin–Madison.

^{||} Department of Chemistry, University of Wisconsin–Madison.

(1) Helguero, L. A.; Faulds, M. H.; Gustafsson, J. A.; Haldosen, L. A. *Oncogene* **2005**, *24*, 6605–6616.

(2) Pettersson, K.; Delaunay, F.; Gustafsson, J. A. *Oncogene* **2000**, *19*, 4970–4978.

been correlated with a more favorable prognosis relative to tumors expressing ER α alone, and many data now suggest that ER α/β heterodimers function to regulate distinct E2-responsive genes.³ However, facile homodimerization has obscured a clear understanding of heterodimer function.

We recently reported a bioluminescent resonance energy transfer (BRET) assay to monitor the ER α/β heterodimer and its respective homodimer formation in live cells, setting the stage to screen for homo- and heterodimer selective ER modulators.⁴ We now report the discovery and total synthesis of a new natural product, actinopolymorphol A (**1**), that preferentially induces ER α/β heterodimerization relative to either homodimer. Actinopolymorphol A represents the first member of a new class of natural products not previously recognized to modulate ER function.

During the course of our efforts to identify novel natural products from microorganisms from diverse and unique ecological niches, we first subjected the crude extract from a recently identified *Actinopolymorpha rutilus* (YIM45725)⁵ originating from a soil sample from the Yunnan province of China to the BRET assay. Initial assays clearly indicated the presence of potential ER α/β heterodimerization inducers in the crude extract. Subsequent large-scale fermentation (5 L) and BRET assay-guided fractionation afforded an active fraction, from which **1**, actinopolymorphol B (**2**), and actinopolymorphol C (**3**) were purified. The ¹H and ¹³C NMR spectra of purified **1**, **2**, and **3** were fully assigned on the basis of extensive 1D and 2D NMR (gCOSY, gHSQC, and gHMBC) and APCI-MS and high-resolution ESI-MS analyses (Figure 1B and also see the Supporting Information). Dereplication revealed **1**, **2**, and **3** to be new natural products (Figure 1A) although a relationship to previously reported natural products (Figure 1C)^{6–10} is clearly evident for **1** and **2**.

Second, we determined, using the BRET assay and pure **1**, **2**, and **3**, that **1** was solely responsible for ER α/β heterodimerization induction (EC₅₀ = 19 μ M) and **2** and **3** were devoid of any ER dimerization inducing activity.¹¹ Consequently, further studies focused exclusively on **1**. Significantly, none of the previously reported natural products to which **1** bears a structural resemblance have been noted as ER modulators. Thus, the finding that **1** induces ER α/β




Figure 1. Actinopolymorphol A (**1**), B (**2**), and C (**3**) from ER α/β heterodimerizing fraction of *A. rutilus* crude extract: (A) structures of **1**, **2**, and **3**; (B) COSY and HMBC correlations observed for **1**, **2**, and **3**; and (C) the satabacin, satazolin, and kurasoin classes of natural products bearing molecular scaffolds similar to **1** or **2**.^{6–10}

heterodimerization may have bearing on the scope of previously reported bioactivities for this class of natural products (Figure 1C).^{6,7,9,10}

Compound **1** was isolated as a colorless oil, for which HR-ESIMS established the molecular formula as C₁₅H₂₀O₄ (287.12567 [M + Na]⁺, calcd for C₁₅H₂₀O₄Na, 287.12538) possessing six degrees of unsaturation. Analysis of the NMR spectra revealed chemical shifts indicative of a ketone (δ_C 207.8), acetoxyl (δ_C 170.9, s; δ_C 19.3, q), one 1,4-disubstituted benzene ring [(δ_H 6.68, 2H; δ_C 115.1, d) and (δ_H 7.02, 2H; δ_C 130.3, d)], a secondary alcohol methine (δ_H 5.08, δ_C 79.6, d), one methine (δ_H 2.03, δ_C 43.7, d), two methylenes (δ_C 47.8, t; δ_C 35.7, t), and two secondary methyl groups [(δ_H 0.82, 3H; δ_C 21.7, q) and (δ_H 0.84, 3H; δ_C 21.7, q)] (Table S1, Supporting Information). On the basis of the gHMBC correlations from two methylene protons to corresponding carbons and the gCOSY correlations, the connectivity of every carbon was readily established (Figure 1B).

Third, we established the structure of **1** by total synthesis using the optically pure starting material (*S*)-2-hydroxy-3-(4-hydroxyphenyl)propionic acid (98% purity) (**4**, Figure 2A). Conversion of **4** to methyl ester **5** was effected by using methanolic HCl followed by treatment with TBSCl and imidazole to afford diprotected methyl ester **6**. The methyl ester underwent smooth conversion to the Weinreb amide **7** by treatment with *N,O*-dimethylhydroxylamine hydrochloride in the presence of (CH₃)₂CHMgCl. Grignard reaction of **7** with (CH₃)₂CHCH₂MgCl proceeded extraordinarily slowly over the course of 24 h but ultimately afforded the diprotected isobutyl ketone **8** in 67% yield from the Weinreb amide. Subsequent cleavage of both silyl ethers with tet-

(3) Chang, E. C.; Frasor, J.; Komm, B.; Katzellenbogen, B. S. *Endocrinology* **2006**, *147*, 4831–4842.

(4) Powell, E.; Xu, W. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 19012–19017.

(5) Wang, Y.-X.; Zhang, Y. Q.; Xu, L. H.; Li, W. J. *Int. J. Syst. Evol. Microbiol.* **2008**, *58*, 2443–2446.

(6) Satabacins and satazolin: Lampis, G.; Deidda, D.; Maullu, C.; Madeddu, M. A.; Pompei, R.; Monache, R. D.; Satta, G. *J. Antibiot.* **1995**, *48*, 967–972.

(7) Soraphinol: Li, X.; Yu, T. K.; Kwak, J. H.; Son, B. Y.; Seo, Y.; Zee, O. P.; Ahn, J. W. *J. Microbiol. Biotechnol.* **2008**, *18*, 520–522.

(8) Kurasoins: Andrus, M. B.; Hicken, E. J.; Stephens, J. C.; Bedke, D. K. *J. Org. Chem.* **2006**, *71*, 8651–8654.

(9) Uchida, R.; Shiomi, K.; Inokoshi, J.; Masuma, R.; Kawakubo, T.; Tanaka, H.; Iwai, Y.; Omura, S. *J. Antibiot.* **1996**, *49*, 932–934.

(10) Uchida, R.; Shiomi, K.; Inokoshi, J.; Nishizawa, A.; Hirose, T.; Tanaka, H.; Iwai, Y.; Omura, S. *J. Antibiot.* **1996**, *49*, 886–889.

(11) Powell, E.; Huang, S.-X.; Xu, Y.; Rajski, S. R.; Wang, Y.; Peters, N.; Guo, S.; Xu, H. E.; Hoffmann, F. M.; Shen, B.; Xu, W. *Biochem. Pharmacol.* **2010**, In press.




Figure 2. Synthesis and chiral HPLC analysis of natural vs synthetic **1**: (A) reagents and conditions for total synthesis of **1** from the optically pure starting material **4** [(a) MeOH, HCl, rt, quantitative; (b) TBSCl, imidazole, DMF, rt, 90%; (c) HN(OMe)CH₃-HCl, (CH₃)₂CHMgCl, THF, 0 °C, 81%; (d) (CH₃)₂CHCH₂MgCl, THF, 0 °C-rt, 67%; (e) TBAF, THF, rt, 86%; (f) Ac₂O, pyridine, rt, 72%; (g) pyrrolidine, 1 min, rt, 66%] and (B) chiral HPLC analysis of racemic **1**, **1** from *A. rutilus* (blue), synthetic **1** (red): ●, **1**; and ▼, the *R*-enantiomer of **1**.

rabutylammonium fluoride (TBAF) afforded the corresponding diol **9** in 81% yield. Notably, production of **9** from **4** constitutes a total synthesis of the natural product 4-hydroxysattabacin from *Sorangium cellulosum* (Figure 1C), for which absolute stereochemistry has been unclear.⁶ The optical rotation of **9** was found to be +18.3 (*c* 0.21, CH₃OH), which compares favorably to the number previously published for 4-hydroxysattabacin [+14 (0.3, CHCl₃)],⁶ assigning 4-hydroxysattabacin as the *S*-enantiomer. Diacetylation of **9** was accomplished in 72% yield upon subjection of the diol to excess acetic anhydride and pyridine. Selective deacetylation of the phenol moiety was accomplished by using neat pyrrolidine at room temperature, affording finally the synthetic **1** as the *S*-enantiomer. ¹H and ¹³C NMR spectral data for synthetic **1** were identical with those previously obtained for natural **1** from *A. rutilus* YIM45725. BRET

(12) Jordan, V. C. *J. Med. Chem.* **2003**, *46*, 883–908.

(13) Jordan, V. C. *J. Med. Chem.* **2003**, *46*, 1081–1111.

assay confirmed that natural and synthetic **1** displayed identical (within standard error) abilities to induce ER α / β heterodimerization.¹¹

Finally, the absolute stereochemistry of **1** was confirmed by comparative chiral HPLC and optical rotation analyses. Both the natural and synthetic **1** were found to be identical upon chiral HPLC analyses, possessing the same retention time (Figure 2B) and identical UV-vis spectra, and to have almost identical optical rotations: +20.9 (*c* 0.15, CH₃OH) for natural **1** and +23.5 (*c* 0.28, CH₃OH) for synthetic **1** (the slight difference in optical rotations is attributed to different concentrations). On the basis of these data, we conclude that **1** from *A. rutilus* YIM45725 has the *S*-configuration at C8; this is consistent with the stereochemistries found in the kurasoins and, as unveiled here, 4-hydroxysattabacin.^{6,8}

Selective ER modulators are well established and a number of them are used clinically such as tamoxifen and raloxifene.^{12,13} Of particular significance during the development of such agents has been the realization that diols or other agents, able to form H-bonding networks with the ligand binding domains of ER spaced a distance of ~12 Å apart, can effectively compete for ER binding with E2 and thereby elicit clinically significant activities. On this basis, we postulate that the phenol moiety of **1** is likely critical to ER heterodimerization activity; an understanding of ER-E2 interactions in which the phenol moiety of E2 is involved in H-bonding supports this hypothesis.¹³ The small size of **1** yet the pronounced ability to modulate ER activity, and the relatedness to already established natural products suggests that further study of this class of natural products is clearly warranted. Our ability to produce **1** biosynthetically and via total synthesis provides an outstanding platform to more rigorously elucidate ER heterodimerization by new analogues bearing this privileged molecular scaffold.

Acknowledgment. We thank the Analytical Instrumentation Center of the School of Pharmacy, UW-Madison for support in obtaining MS and NMR data. This work is supported in part by NIH grants CA113297 (B.S.), CA125387 (W.X.), MH089442 (W.X.), and T32CA009135 (E.P.) and the Chinese Ministry of Education 111 Project B08034 (Y.D.).

Supporting Information Available: Full experimental details for strain cultivation, extraction, compound isolation, spectral data, and assignment of all natural products and synthetic intermediates. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL1013526