

The Components of xRNA: Synthesis and Fluorescence of a Full Genetic Set of Size-Expanded Ribonucleosides


Armando R. Hernández and Eric T. Kool*

Department of Chemistry, Stanford University, Stanford, California 94305-5080, United States

kool@stanford.edu

Received December 2, 2010

ABSTRACT



The synthesis and properties of a full set of four benzo-expanded ribonucleosides (xRNA), analogous to A, G, C, and U RNA monomers, are described. The nucleosides are efficient fluorophores with emission maxima of 369–411 nm. The compounds are expected to be useful as RNA pathway probes and as components of an unnatural ribopolymer.

Ribonucleosides and ribonucleotides with unnatural nucleobases have found widespread use as chemical tools to study the functional roles,^{1a,b} hydrogen bonding requirements,^{1c} structure/activity relationships,^{1d} and mechanistic details^{1e} of RNAs in biological systems. In particular, the use of fluorescent ribonucleotides as biological probes and sensors has received much attention.² In an effort to minimally perturb the recognition of RNAs by natural biomolecules, most fluorescent ribonucleotides are designed to retain some of the original canonical Watson–Crick base-pairing recognition groups found in natural RNA.³

(1) (a) Lu, J.; Li, N.-S.; Koo, S. C.; Piccirilli, J. A. *J. Org. Chem.* **2009**, *74*, 8021. (b) Cho, H. D.; Oyelere, A. K.; Strobel, S. A.; Weiner, A. M. *RNA* **2003**, *9*, 970. (c) Somoza, A.; Chelliserrykattil, J.; Kool, E. T. *Angew. Chem., Int. Ed.* **2006**, *45*, 4994. (d) Pokharel, S.; Jayalath, P.; Maydanovych, O.; Goodman, R. A.; Wang, S. C.; Tantillo, D. J.; Beal, P. A. *J. Am. Chem. Soc.* **2009**, *131*, 11882. (e) Araki, L.; Morita, K.; Yamaguchi, M.; Zhao, Z.-Y.; Wilson, T. J.; Lilley, D. M. J.; Harusawa, S. *J. Org. Chem.* **2009**, *74*, 2350.

(2) (a) Stengel, G.; Urban, M.; Purse, B. W.; Kuchta, R. D. *Anal. Chem.* **2010**, *82*, 1082. (b) Srivatsan, S. G.; Tor, Y. *Nat. Protoc.* **2007**, *2*, 1547. (c) Kimoto, M.; Mitsui, T.; Harada, Y.; Sato, A.; Yokoyama, S.; Hirao, I. *Nucleic Acids Res.* **2007**, *35*, 5360.

(3) Zhang, C.-M.; Liu, C.; Christian, T.; Gamper, H.; Rozenski, J.; Pan, D.; Randolph, J. B.; Wickstrom, E.; Cooperman, B. S.; Hou, Y.-M. *RNA* **2008**, *14*, 2245.

Previously, Leonard *et al.* described the synthesis of fluorescent ribonucleoside analogs of adenosine (**1**, Figure 1) and guanosine (**2**) with a benzene ring incorporated within the purine scaffold that expanded their size by 2.4 Å.^{4,5} Among other features, these analogs retained their canonical Watson–Crick base-pairing groups. Nucleoside triphosphate derivatives of **1** and **2** have been prepared





Figure 1. The xRNA genetic set.

enzymatically;⁶ however, technological limitations at the time did not allow for the synthesis of oligoribonucleotides using these novel RNA monomers. Furthermore, analogous benzo-expanded analogs of cytidine (**3**) and uridine (**4**) have never been reported. One example of a size-expanded uridine analog has been reported;⁷ however, the glycosidic bond is at the same N1-position as natural pyrimidine ribonucleosides and is therefore not structurally analogous to **1** and **2** in a size-expanded RNA genetic set.

Our laboratory has previously reported the design and synthesis of a set of four size-expanded DNA analogs (xDNA).^{8,9} After incorporation into oligonucleotides using automated oligonucleotide synthesis, these compounds were studied for their unusual biophysical and biochemical properties in DNA.^{10,11} Herein we describe the synthesis of the complete set of size-expanded RNA analogs (xRNA, Figure 1) and report their photophysical properties. This information will provide a foundation on which to design experiments to use xRNA monomers and polymers in the study of ribonucleotide and RNA pathways in biological systems.


Scheme 1. Synthesis of rxA Triol (**1**)



The syntheses of the expanded purine nucleosides **1** (rxA) and **2** (rxG) were similar to the syntheses of their xDNA counterparts, dxA and dxG,^{8,9} and to previous approaches to the compounds as reported by Leonard and co-workers.^{4,5} The protected xA nucleobase (**5**, Scheme 1) was coupled to 1,2,3,5-tetra-*O*-acetyl- β -D-ribose (*rib*) under Vorbrüggen conditions.¹² This method for forming *N*-nucleosides is a less hazardous alternative to Leonard's

approach of using mercuric cyanide as a coupling agent.⁴ Although both methods are known to form the desired β -anomers of the nucleosides, they also generate a mixture of N3- and undesired N1-regioisomers. After chromatographic purification of the *N*-ribosylated products, an overall yield of 61% was obtained in a ratio of 60:40 (**6a:6b**). The correct regioisomer (**6a**) was identified by 2D-NMR (Supporting Information Figure S1). To obtain additional amounts of the desired N3-regioisomer, **6b** was converted to **6a** via a transisomerization reaction involving catalytic amounts of *p*-toluenesulfonic acid.¹³ Leonard did not report the overall yield and regioisomeric ratios of their *N*-ribosylation reaction, so it is difficult to compare the prior approach to the effectiveness of the present Vorbrüggen approach for generating the rxA precursors. A tandem *O*-deacetylation/nucleophilic aromatic substitution reaction of **6a** with ammonia generated **1** in good yield. Using these improved methods, compound **1** can be synthesized in 5.1% overall yield over 8 steps or 6.0% overall yield in 9 steps (including the transisomerization of **6b**).

Scheme 2. Synthesis of rxG Triol (**2**)



The synthesis of rxG, as shown in Scheme 2, has been previously reported by us as an intermediate in dxG synthesis by reaction of **7** with *rib* under Vorbrüggen conditions.⁹ However, the N3- and N1-regioisomeric mixture (**8a** and **8b**, respectively) could only be separated in later steps. To improve rxG synthesis, we found an eluant mixture for column chromatography that successfully separated **8a** and **8b**. The correct regioisomer (**8a**) was identified using 2D-NMR (Figure S1). As was done in rxA synthesis, we were also able to generate more of the desired **8a** regioisomer by transisomerization of **8b**. Mild deacetylation by aminolysis of **8a** generated **2** in 87% yield. Compound **2** was prepared in 5.0% overall yield over 7 steps or 6.3% overall yield in 8 steps (including the transisomerization of **8b**).

The novel expanded pyrimidines **3** (rxC) and **4** (rxU) were modeled after their xDNA counterparts, dxC and dxT.^{8,9} A major structural difference between the xDNA and xRNA pyrimidine sets is the presence and absence of a methyl group on the C6 position of their nucleobases,

(4) Leonard, N. J.; Sprecker, M. A.; Morrice, A. G. *J. Am. Chem. Soc.* **1976**, *98*, 3987.

(5) Keyser, G. E.; Leonard, N. J. *J. Org. Chem.* **1979**, *44*, 2989.

(6) (a) Leonard, N. J.; Keyser, G. E. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, *76*, 4262. (b) Leonard, N. J.; Scopes, D. I. C.; Van Der Lijn, P.; Barrio, J. R. *Biochemistry* **1978**, *17*, 3677.

(7) Xie, Y.; Maxson, T.; Tor, Y. *J. Am. Chem. Soc.* **2010**, *132*, 11896.

(8) Liu, H.; Gao, J.; Maynard, L.; Saito, Y. D.; Kool, E. T. *J. Am. Chem. Soc.* **2004**, *126*, 1102.


(9) Liu, H.; Gao, J.; Kool, E. T. *J. Org. Chem.* **2005**, *70*, 639.

(10) Gao, J.; Liu, H.; Kool, E. T. *J. Am. Chem. Soc.* **2004**, *126*, 11831.

(11) Krueger, A. T.; Kool, E. T. *J. Am. Chem. Soc.* **2008**, *130*, 3989.

(12) Vorbrüggen, H.; Höfle, G. *Chem. Ber.* **1981**, *114*, 1256.

(13) Boryski, J.; Golankiewicz, B. *Nucleosides Nucleotides* **1989**, *8*, 529.


Scheme 3. Synthesis of rxU Triol (**4**)

respectively. This difference required a substantial change of synthetic approach. Our strategy for the synthesis of xRNA *C*-nucleosides involved metallation of an aryl halide nucleobase followed by nucleophilic addition to 2,3,5-tri-*O*-benzyl- β -D-ribo-1,4,-lactone (*rIac*) and subsequent reduction at C-1', as was done previously in our laboratory.¹⁴ For the synthesis of rxU, we envisioned the use of 8-bromo-2,4-dimethoxyquinazoline (**12**, Scheme 3) as an aryl halide. After reduction at C-1', the nucleoside would serve as a common intermediate between rxU and rxC syntheses.

To the best of our knowledge, the synthesis of the brominated heterocycle intermediate **12** has not been previously reported. We investigated various synthetic approaches to generate a sizable amount of **12**. In the end, we found that this could be best accomplished beginning with the synthesis of 2-amino-3-bromobenzonitrile, **9**. Jiang and Lai previously reported the synthesis of 2-amino-3-chlorobenzonitrile from 2,6-dichloroaniline using copper(I) cyanide in NMP in 31% yield.¹⁵ Replacement of the starting material with commercially available 2,6-dibromoaniline and improving reaction conditions afforded **9** in 71% yield. Cyclization of **9** using carbon dioxide gas in the presence of DBU produced **10** in 91% yield, as was reported for the cyclization of various

2-aminobenzonitriles.¹⁶ Following this, compounds **11** and **12** were synthesized in excellent yield employing methods previously used by our laboratory.⁸

Various metallating agents were screened in the *C*-ribosylation reaction of **12** with the ribonolactone. We found that 1.1 equiv of *tert*-butyllithium best generated the activated species for the nucleophilic addition reaction. Following reduction at C-1' using $\text{BF}_3 \cdot \text{OEt}_2$ and triethylsilane, a single product, **13**, was formed in 65% yield. gHMBC and ROESY studies on **13** revealed that the xU nucleobase was attached to the sugar in the correct C8-position and that the nucleoside was the desired β -anomer (Figure S2).

Scheme 4. Synthesis of rxC Triol (**3**)

At this point, we were interested in using **13** as the branching point between rxU and rxC syntheses. It is known that boron tribromide is an effective debenzylating reagent.¹⁷ BBr_3 also serves as a demethylating agent for aryl methyl ethers.¹⁸ We found that by controlling the amount of BBr_3 and reaction conditions on **13**, we could generate two different and useful products. Using 10 equiv of BBr_3 beginning at -78 °C and warming to room temperature while stirring overnight generated **4** in 75% yield. On the other hand, using milder conditions (3 equiv of BBr_3 kept at -78 °C with stirring for 3 h) generated the 2,4-dimethoxyquinazoline triol nucleoside, **14** (Scheme 4), in 79% yield. To obtain the desired rxC precursor, **15**, we performed an aromatic nucleophilic substitution reaction on **14** using ammonia in methanol. We found that only one monoamino substituted product was formed in very good yield. ^1H NMR data showed the presence of a methoxy and amino group in the compound. ROESY analysis of this product (**15**) revealed that the methoxy and amino groups were in the desired 2- and 4-positions,

Table 1. Photophysical Data for xRNA Triols in Methanol

xRNA triol	λ_{max} abs. ^a (nm)	λ_{max} ex. (nm)	λ_{max} em. (nm)	ϵ_{260} (L/mol·cm)	ϵ_{max} (L/mol·cm)	Φ^b	brightness ^c (L/mol·cm)	τ (ns)
rxA	331	331	373	18 500	10 500	0.44	4600	3.6
rxG	326	337	411	4900	4500	0.40	1800	7.3
rxC	322	322	379	5300	4500	0.48	2200	4.2
rxU	311	311	369	1000	3900	0.27	1050	2.9

^a Long λ maximum. ^b Error $\leq \pm 0.03$. ^c Brightness calculated as $\epsilon_{\text{max}} \times \Phi$.




Figure 2. Fluorescence spectra of rxA (A), rxG (B), rxC (C), and rxU (D) triols in methanol. Excitation spectra were monitored at the emission maxima.

respectively (Figure S2). Demethylation of **15** using sodium iodide in acetic acid generated **3** in excellent yield. Compound **3** was synthesized in 19.2% overall yield over 8 steps, and compound **4** was prepared in 22.6% yield in 6 steps.

To evaluate the utility of the xRNA nucleosides to be employed as fluorescence probes, we measured the photophysical properties of the entire genetic set (**1–4**) in methanol (Table 1) and water at pH 7.0 (Table S1). Previously, we reported the photophysical properties of the xDNA monomers in methanol.¹¹ In addition, partial photophysical data for rxA and rxG triols have been reported by Leonard.^{4,5,6a}

Our experiments reveal that all four nucleosides have high quantum yields (Table 1). The data show that rxA is the brightest of the four, owing to its higher molar absorptivity. In water, the data for rxA are in good agreement with those of Leonard.⁴ In methanol, the fluorescence spectra of rxA (Figure 2A) and those of dxA are similar, with the exception that the emission of dxA is red-shifted by 20 nm.¹¹

The rxG nucleoside emits furthest to the red (411 nm) of the set. Our data for rxG in water seem to be in general agreement with previous limited data for rxG triol in aqueous buffer.^{5,6a} We also find that rxG in methanol (Figure 2B) has very similar photophysical properties to those of dxG in the same solvent.¹¹ Interestingly, the emission and excitation wavelengths of rxG are all blue-shifted (by *ca.* 20 nm) in water as compared to data taken in methanol.

We also measured the photophysical properties of the previously unknown xRNA pyrimidines (rxC and rxU). In comparison to the xDNA pyrimidines,¹¹ there is a small blue-shifting of absorption, excitation, and emission maxima by ~ 10 nm (in each case) in the xRNA pyrimidine nucleosides. This difference is likely due to the absence of the C6 methyl group in rxC and rxU. We find that the two have virtually the same properties in methanol (Figure 2C and D) or water.

In summary, we have synthesized a complete four-letter benzo-expanded RNA genetic set, in good yield and purity. We have also demonstrated the photophysical properties of the xRNA nucleosides, which we find to be efficient fluorophores. Future work will be directed toward the use of these compounds as nucleotide probes and in unnatural RNA polymers.

Acknowledgment. This work was supported by the National Institutes of Health (GM063587). We also acknowledge the NIH for support of A.H. as a Ruth L. Kirschstein NRSA predoctoral fellow (GM084680). We thank Dr. Urvashi Sahni (Stanford University) for technical assistance and Dr. Stephen R. Lynch (Stanford University) for assistance with NMR experiments.

Supporting Information Available. Experimental procedures, NMR data, fluorescence experiment details, UV–vis and fluorescence spectra, and fluorescence lifetime data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

- (14) Silverman, A. P.; Kool, E. T. *J. Am. Chem. Soc.* **2007**, *129*, 10626.
- (15) Jiang, J.; Lai, Y.-H. *J. Am. Chem. Soc.* **2003**, *125*, 14296.
- (16) Mizuno, T.; Ishino, Y. *Tetrahedron* **2002**, *58*, 3155.
- (17) Kundu, N. G.; Hertzberg, R. P.; Hannon, S. J. *Tetrahedron Lett.* **1980**, *21*, 1109.
- (18) Sakamoto, T.; Kondo, Y.; Sato, S.; Yamanaka, H. *Tetrahedron Lett.* **1994**, *35*, 2919.