

Stereospecific Ring Expansion of Chiral Vinyl Aziridines


Matthew Brichacek,[†] Mauricio Navarro Villalobos,[†] Alexandra Plichta,[†] and Jon T. Njardarson^{*,†,‡}

Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853, United States, and Department of Chemistry and Biochemistry, University of Arizona, 1306 East University Blvd., Tucson, Arizona 85721, United States

njardars@email.arizona.edu

Received December 26, 2010

ABSTRACT



In this report, it is demonstrated that chiral vinyl aziridines can be stereospecifically ring expanded. This synthetic approach allows controlled access to chiral 2,5-*cis*- or 2,5-*trans*-3-pyrroline products from starting materials with the appropriate aziridine geometry. Twenty three ring expansion examples, most of which feature a stereospecific cyclization, are presented.

A cursory review of the structural motifs of the top 200 top selling drugs¹ reveals that around 90% contain at least one nitrogen atom and approximately 65% are decorated with a heterocycle. Not surprisingly, the majority of these heterocycles are nitrogenous, with pyrrolidines a commonly occurring heterocyclic scaffold. Given the success of chiral pyrrolidines as important pharmaceutical building blocks it follows that a range of practical synthetic methods are needed² to provide access to any targeted structural and stereochemical pyrrolidine pattern.

We have chosen to tackle the challenge of developing useful pyrrolidine forming methods by revisiting the ring expansion of vinyl aziridines, first reported by Atkinson.³ Surprisingly, despite the potential usefulness of converting a vinylaziridine into a 3-pyrroline, there had only been a single study focused on using metal catalysts to aid the

rearrangement prior to our contribution to this field.⁴ Oshima and co-workers found that tosylated diene-aziridines could be ring expanded in the presence of a palladium catalyst to the 3-pyrrolines. Both the *N*-tosyl group and the diene moiety were reported to be essential for the success of this rearrangement. Simple nondienic vinyl aziridines did not ring expand, furnishing instead a complex mixture of products. In our recent report, we demonstrated that this significant substrate limitation could be solved by using Cu(hfacac)₂ as a catalyst.⁵ The substrate scope of this new transformation, which we demonstrated for a range of tosyl (Ts) and *N*-phthalimide (NPhth) protected vinyl aziridines, was shown to be quite broad. In this report we expand these investigations further and focus our attention on stereospecific vinyl aziridine ring expansions and application of this new methods toward accessing chiral pyrroline products.

To maximize the synthetic potential of our method for preparing chiral pyrroline products, it is essential that reliable, asymmetric, convergent, and scalable routes be available to access the requisite starting materials (chiral

[†] Cornell University.

[‡] University of Arizona.

(1) The structures of the top 200 selling drugs can be downloaded from <http://cbc.arizona.edu/njardarson/group>. McGrath, N. A.; Brichacek, M.; Njardarson, J. T. *J. Chem. Educ.* **2010**, *87*, 1348–1349.


(2) *Asymmetric Synthesis of Nitrogen Heterocycles*; Royer, J., Ed.; Wiley-VCH: Weinheim, Germany, 2009.

(3) (a) Atkinson, R. S.; Rees, C. W. *Chem. Commun.* **1967**, 1232–1232. (b) Hudlicky, T.; Frazier, J. O.; Kwart, L. D. *Tetrahedron Lett.* **1985**, *26*, 3523–3526. (c) Pearson, W. H. *Tetrahedron Lett.* **1985**, *26*, 3527–3530. (d) Hudlicky, T.; Reed, J. W. *Angew. Chem., Int. Ed.* **2010**, *49*, 4864–4876. (e) Scheiner, P. *J. Org. Chem.* **1967**, *32*, 2628–2630. (f) Hirner, S.; Somfai, P. *Synlett* **2005**, 3099–3102.

(4) (a) Fugami, K.; Morizawa, Y.; Oshima, K.; Nozaki, H. *Tetrahedron Lett.* **1985**, *26*, 857–860. (b) Fugami, K.; Miura, K.; Morizawa, Y.; Oshima, K.; Utimoto, K.; Nozaki, H. *Tetrahedron* **1989**, *45*, 3089–3098.

(5) (a) Brichacek, M.; Lee, D.; Njardarson, J. T. *Org. Lett.* **2008**, *10*, 5023–5026. (b) For a review of 3-pyrroline synthetic approaches: Brichacek, M.; Njardarson, J. T. *Org. Biomol. Chem.* **2009**, *7*, 1761–1770.

Scheme 1. Retrosynthetic Analysis for Chiral Pyrrolines



vinyl aziridines).⁶ The union of an imine and suitably activated nucleophile quickly emerged as the optimal approach (Scheme 1). The imine-based retrosynthetic analysis is highlighted for chiral pyrroline **3**, which we envisioned would originate from the copper-catalyzed ring expansion of a *trans*- or *cis*-vinyl aziridine (**4** and **5**). These isomeric vinyl aziridines could be accessed by treating imines **7** and **8** with nucleophiles **6** and **9**, respectively. An attractive feature of this disconnection approach is that either the imine or the nucleophile could serve the role of chiral auxiliary.

Prior to assembling the requisite chiral aziridines it was of critical importance to learn what other nitrogen protecting groups (**P**) besides Ts and Phth might also be suitable for this reaction (Table 1). We realized that aryl- and acyl-substituted aziridines would be particularly challenging substrates, given their known tendency to undergo competing Claisen rearrangements or intramolecular displacement reactions.⁷ This prediction turned out to be true as in the case of *N*-benzyl (entry d) a known hydride shift occurred instead,⁸ while for *N*-phenyl (entry c) the expected Claisen rearrangement was observed.⁹ Boc-protected aziridine **10** (entry e) did ring expand to the desired 3-pyrroline,¹⁰ while in contrast when a benzoyl group (entry j) was present, the ring expansion failed and instead a mixture of five- and seven-membered heterocycles was formed.¹¹ When vinyl aziridine **10** was not protected (NH, entry i) it rapidly oligomerized when subjected to the reaction conditions. Curiously, the *tert*-butyl sulfinamide-substituted vinyl aziridine (entry f) formed 1-phenylbutadiene in high yield rather than the expected 3-pyrroline product (**11**).¹² We were delighted to find that 4-nitrobenzylsulfonamide (Ns) and Bus *tert*-butylsulfonamide (Bus) groups, in

Table 1. Ring Expansion: Nitrogen Substituent Tolerance^h

entry	P =	time (h)	yield of 11 (%)
a	Ts	5	92
b	NPhth ^{a,b}	2	97
c	Ph ^a	2	0 ^c
d	Bn ^a	2	0 ^d
e	Boc	5	60
f	S(O) ^f Bu	1	0 ^e
g	Bus	5	86
h	Ns	5	94
i	H	2	0 ^f
j	Bz	5	0 ^g

^a Racemic aziridine used. ^b Cis/trans mixture. ^c Mixture of azepine and 2-pyrroline products. ^d (*Z*)-*N*-Benzylidene-1-phenylbut-2-en-1-amine isolated in 86% yield. ^e (*E*)-Buta-1,3-dien-1-ylbenzene formed in 69% yield. ^f Polymer formed. ^g Five and seven membered ring products observed. ^h Conditions: 5 mol % Cu(hfacac)₂, 150 °C.


addition to *p*-toluenesulfonamide (entry a), were compatible with the ring expansion conditions. These studies suggest that sulfonamides are especially well suited for our ring expansion reaction.

Armed with insights into what aziridine protecting groups are compatible with our reaction conditions we turned our attention to the design of a scalable asymmetric route to vinyl aziridine substrates. The use of proline derivatives as organocatalysts¹³ and as key building blocks of many pharmaceuticals and natural products¹⁴ ensures its status as a privileged structural motif. Chiral 2,3-dehydroproline (3-pyrroline) products commonly originate from natural 3-hydroxy proline,¹⁵ and this neglected family of chiral proline products emerged as an ideal target for our new methodology. Retrosynthetic analysis for 2,5-dihydropyrrolidine **15** (Scheme 2) suggests that the chiral vinyl aziridine ring expansion substrate (**14**) should originate from a Darzens reaction¹⁶ between a bromo acetate (**12**) and a chiral conjugated Ellman type imine¹⁷ (**13**). These chiral imines are attractive substrates because they are trivial to make, very stable, available in both enantiomeric forms, and easy to handle. Since our nitrogen substituent ring expansion compatibility study (Table 1) had

(6) Pellisier, H. *Tetrahedron* **2010**, *66*, 1509–1555.
 (7) Pommelet, J. C.; Chuche, J. *Can. J. Chem.* **1976**, *54*, 1571–1581.
 (8) Somfai, P.; Ahman, J. *Tetrahedron Lett.* **1995**, *36*, 1953–1956.
 (9) Fantauzzi, S.; Gallo, E.; Caselli, A.; Piangiolino, C.; Ragaini, F.; Re, N.; Cenini, S. *Chem.—Eur. J.* **2009**, *15*, 1241–1251.
 (10) Mishra, A.; Rice, S. N.; Lwowski, W. *J. Org. Chem.* **1968**, *33*, 481–486.
 (11) Heine, H. W.; Fetter, M. E.; Nicholson, E. M. *J. Am. Chem. Soc.* **1959**, *81*, 2202–2204.
 (12) We have not been able to find other examples of this type of deamination reaction for sulfinamide protected aziridines. Furthermore, we have tested this on several other substrates and observed the same deamination behavior. Aziridine deaminations are not without precedence, see: (a) Clard, R. D.; Helmkamp, G. K. *J. Org. Chem.* **1964**, *29*, 1316–1320. (b) Muller, R. K.; Felix, D.; Schreiber, J.; Eschenmosher, A. *Helv. Chim. Acta* **1970**, *53*, 1479–1484.

(13) (a) Kotsuki, H.; Ikishima, H.; Okuyama, A. *Heterocycles* **2008**, *75*, 493–529. (b) Kotsuki, H.; Ikishima, H.; Okuyama, A. *Heterocycles* **2008**, *75*, 757–797.
 (14) Mauger, A. M. *J. Nat. Prod.* **1996**, *59*, 1205–1211.
 (15) (a) Meyer, S. C.; Ramanjulu, J.; Vera, M. D.; Pfizenmayer, A. J.; Jollie, M. M. *J. Org. Chem.* **1994**, *59*, 5192–5205. (b) Greenwood, E. S.; Hitchcock, P. B.; Parsons, P. J. *Tetrahedron* **2003**, *59*, 3307–3314.
 (16) Davis, F. A.; Liu, H.; Zhou, P.; Fang, T.; Reddy, G. V.; Zhang, Y. *J. Org. Chem.* **1999**, *64*, 7559–7567.
 (17) (a) Liu, G.; Cogan, D. A.; Ellman, J. A. *J. Am. Chem. Soc.* **1997**, *119*, 9913–9914. (b) Ellman, J. A.; Owens, T. D.; Tang, T. P. *Acc. Chem. Res.* **2002**, *35*, 984–995. (c) Davis, F. A.; Reddy, R. T.; Reddy, R. E. *J. Org. Chem.* **1992**, *57*, 6387–6389. (d) Davis, F. A. *J. Org. Chem.* **2006**, *71*, 8993–9003.

Scheme 2. Darzen Reaction Route to Chiral Proline Products



unfortunately revealed that the immediate products of this union, *N*-sulfonamide protected vinyl aziridines, did not ring expand we needed to resort to oxidizing the Darzen reaction products prior to ring expansion in order to have a

Table 2. Ring Expansion of Bus-Protected Vinyl Aziridines^a


entry	substrate (14)	product (15)	yield
a			72%
b			77%
c			87%
d			86%
e			90%
f			90%
g			81%
h			83%
i			67%
j			87%
k			59%
l			50%

^a Conditions: 5 mol % Cu(hfacac)₂, toluene, 5 h, 150 °C. Cy = cyclohexyl. R = Bus (14a–l) R = S(O)^tBu (16a–l).

compatible substrate (Bus-protected aziridines). This was readily accomplished by using *m*-CPBA. Interestingly, we have learned that this oxidation can also be accomplished with ammonium molybdate tetrahydrate in excellent yield.¹⁸

Twelve chiral aziridine substrates shown in Table 2 (R = Bus) were prepared by using the strategy detailed above. The Darzen reaction step was in general high yielding and the *cis/trans*-aziridine ratios were in the modest 2:1–4:1 range. To better understand the scope of the ring expansion, the resulting *cis*- and *trans*-vinyl aziridines readily ring expand to 3-pyrrolines in excellent yields in the presence of catalytic amounts of Cu(hfacac)₂. Entries e–h highlight the methodologies ability to offer two suitable retrosynthetic choices for accessing a particular enantiomer (entries e and f give the (*S*)-enantiomer and g and h the (*R*)-enantiomer). Entries i–k demonstrate how control of a desired 2,5-pyrroline substitution pattern can be achieved simply by starting with the appropriate *cis*- or *trans*-aziridine precursor.¹⁹ Deprotection of Bus-protected cyclic dialkylamines containing an adjacent electron-withdrawing group like the 3,4-dehydropyrroline products in Table 2 is well documented to proceed cleanly without any epimerization.²⁰

Scheme 3. Asymmetric Synthesis of Tosyl Protected Aziridines



A complementary convergent asymmetric approach to chiral vinyl aziridines en route to 3-pyrroline products employs a chiral sulfide in place of a chiral imine (Scheme 3). Although the use of chiral nucleophiles versus chiral electrophiles for accessing aziridines is currently far less developed, this approach would avoid the oxidation step prior to ring expansion. Aggarwal²¹ and co-workers have shown that chiral sulfonium nucleophiles can be used to synthesize aziridines in a highly selective manner. We chose to employ the limonene-based chiral auxiliary (19)²² for

(18) This cheap, practical, and simple oxidation protocol, which is commonly employed for accessing Julia olefination substrates, has not been reported previously for converting sulfonamides to sulfonamides: Blakemore, P. R. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2563–2585.

(19) Entry l is an exception as this highly congested system scrambled prior to ring expansion.

(20) (a) Koep, S.; Gais, H.-J.; Raabe, G. *J. Am. Chem. Soc.* **2003**, *125*, 13243–13251. (b) Gunter, M.; Gais, H.-J. *J. Org. Chem.* **2003**, *68*, 8037–8041. (c) Tiwari, S. K.; Gais, H.-J.; Lindenmaier, A.; Babu, G. S.; Raabe, G.; Reddy, L. R.; Kohler, F.; Gunter, M.; Koep, S.; Iska, V. B. R. *J. Am. Chem. Soc.* **2006**, *128*, 7360–7373.

(21) Aggarwal, V. K.; Vasse, J.-L. *Org. Lett.* **2003**, *5*, 3987–3990.

Table 3. Ring Expansion of Tosyl-Protected Vinyl Aziridines^a

entry	Substrate (20)	product (21)	t (h)	yield
a	R ₁ = H R ₂ = Ph R ₃ = 4-Cl-C ₆ H ₄		2	84%
b	R ₁ = H R ₂ = Ph R ₃ = 4-F-C ₆ H ₄		2	91%
c	R ₁ = H R ₂ = Ph R ₃ = 4-Br-C ₆ H ₄		2	92%
d	R ₁ = H R ₂ = Ph R ₃ = 4-I-C ₆ H ₄		2	87%
e	R ₁ = H R ₂ = Ph R ₃ = 2-OMe-C ₆ H ₄		2	81%
f	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = 2-Br-C ₆ H ₄		2	97%
g	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = C ₆ H ₅		2	92%
h	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = 4-CF ₃ -C ₆ H ₄		2	93%
i	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = 4-NO ₂ -C ₆ H ₄		5	96%
j	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = 4-CF ₃ -C ₆ H ₄		4	92%
k	R _{1, R₂} = -(CH ₂) ₄ - R ₃ = 4-NO ₂ -C ₆ H ₄		5	71%

^a Conditions: 5 mol % Cu(hfacac)₂, toluene, 150 °C.

additions to sulfonamide imines (**18**). The Aggarwal chiral auxiliary is easily prepared in either enantiomeric form. This route provided us with the chiral ring expansion precursors (**20**) in a single step. Yields for this step were mostly high and the *cis/trans*-aziridine product ratios were in the 6:1–20:1 range. Interestingly, the bis-aryl vinyl aziridine products (**20a–e**) were obtained as a single

(22) Illa, O.; Arshad, M.; Ros, A.; McGarrigle, E. M.; Aggarwal, V. K. *J. Am. Chem. Soc.* **2010**, *132*, 1828–1830.

trans-isomer (dr > 20:1) while the cyclohexyl-substituted aziridines (**20f–k**) were obtained with slightly lower selectivity (dr = 6:1–11:1).

Chiral 2-aryl-substituted pyrrolidines are important structures and have received increased attention in recent years.²³ We were delighted to find that all of the chiral vinyl aziridine substrates **20a–k** could be ring expanded stereospecifically using Cu(hfacac)₂ as catalyst in uniformly excellent yields to corresponding 2-aryl-3-pyrrolines **21a–k** (Table 3). A single stereoisomer was obtained in each case. Both electron-donating and -withdrawing substituents are well-tolerated, including aryl bromides, chlorides, iodides, and fluorides. The broad tolerance of aryl substituents should serve those researchers interested in applying this new methodology to complex natural products, pharmaceuticals, or organocatalytic architectures well. Impressively, the corresponding chiral *cis*-vinyl aziridines stereospecifically afford the *trans*-fused 3-pyrroline products as single stereoisomeric products (entries j and k).

In summary, we have demonstrated two new practical complementary routes to access valuable chiral 3-pyrroline products from readily accessible vinyl aziridine precursors. This was accomplished by coupling an imine with an appropriate nucleophile and then catalytically ring expanding the resulting chiral vinyl aziridine with Cu(hfacac)₂. The substrate scope and functional group tolerance of this new synthesis is very broad. The excellent chirality transfer demonstrated by the examples presented also serves to underscore the mechanistic uniqueness of this new catalytic reaction and advantage over all other aziridine ring expansion routes aiming to form 3-pyrrolines. It is important to note that an added advantage to both of our asymmetric 3-pyrroline routes is that they provide a double bond handle for further functionalization, which can be readily converted to more complex pyrrolidine products in a substrate-controlled manner. As better asymmetric aziridine methods are developed more opportunities will arise for applications of this new method.

Acknowledgment. We would like to thank NSF (CHE-0848324) and the NIH (training grant, T32GM008500, to M.B.) for financial support. Special thanks go to Dr. Emil Lobkovsky for obtaining all X-ray crystal structures.

Supporting Information Available. Full experimental details for all new compounds reported in this article including X-ray data for compounds **15d**, **15j**, **15l**, **21d**, and **21f**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(23) For recent approaches to chiral 2-pyrrolines consult: (a) Campos, K. R.; Klapars, A.; Waldman, J. H.; Dormer, P. G.; Chen, C. *J. Am. Chem. Soc.* **2006**, *128*, 3538–3539. (b) Davis, F. A.; Song, M.; Augustine, A. *J. Org. Chem.* **2006**, *71*, 2779–2786. (c) Fang, Y.-Q.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2008**, *130*, 5660–5661.